Index to Authors*

A

Abdel-Latif A (*see* Banchs et al). 2006;19:1401.e1-e3 (Case rep.)
Abe H (*see* Iwahashi et al). 2006;19:1238-44
Abe Y (*see* Pulerwitz et al). 2006;19:540-5
Abergel E (*see* Cohen et al). 2006;19:580 (Letter)
Abraham TP (*see* Bos et al). 2006;19:1033-7
Abraham TP (*see* Wang et al). 2006;19:1102-11
Ackerman MJ (*see* Sorajja et al). 2006;19:788-95
Adamopoulos S (*see* Paraskevaidis et al). 2006;19:529-35
Aeschlimann SE (*see* Gepner et al). 2006;19:1170-4
Aeschlimann SE (*see* Gepner et al). 2006;19:223-8
Aeschlimann SE (*see* Tzou et al). 2006;19:1286-92
Aggarwal K (*see* Gosavi et al). 2006;19:1529.e1-e3 (Case rep.)
Agler DA (*see* Song et al). 2006;19:1129-35
Agmon Y (*see* Dabbah et al). 2006;19:733-43
Agrawal N (*see* Banchs et al). 2006;19:1401.e1-e3 (Case rep.)
Agresti C (*see* Palmieri et al). 2006;19:483-90
Agricola E (*see* Picano et al). 2006;19:356-63 (Review article)
Agricola E, Picano E, Oppizzi M, Pisani M, Meris A, Fragasso G, Margonato A. Assessment of stress-induced pulmonary interstitial edema by chest ultrasound during exercise echocardiography and its correlation with left ventricular function. 2006;19:457-63
Ahmadi-Kashani M (*see* Movahed et al). 2006;19:911-3
Ahmed I, Katz DH, Crooke GA, Li M, Doddamani S, Haramati LB, Ostfeld RJ, Gordon GM, Spevack DM. Biventricular mural vegetations in a patient without valvular pathology. 2006;19:938 (Case rep.)
Ahn J-A (*see* Choi et al). 2006;19:1211-9
Ahn J-A (*see* Ha et al). 2006;19:1314-9
Aiazian E (*see* Dijkmans et al). 2006;19:285-93
Akasaka T (*see* Akiyama et al). 2006;19:55-63
Akasaka T (*see* Katayama et al). 2006;19:335-40
Akasaka T (*see* Kume et al). 2006;19:914-8
Akasaka T (*see* Watanabe et al). 2006;19:71-5
Akiyama M, Akasaka T, Fujimoto K, Koyama Y, Watanabe N, Neishi Y, Tomita J, Toyoda T, Kaji S, Yoshida K. Heterogeneity of myocardial perfusion in distal coronary embolism with different particle sizes. 2006;19:55-63
Akiyama S (*see* Hirano et al). 2006;19:536-9
Aletras AH (*see* Sachdev et al). 2006;19:34-9
Allman KC (*see* Chung et al). 2006;19:347-53
Alper AT (*see* Uslu et al). 2006;19:341-6
Alshaher MM (*see* Silverstein et al). 2006;19:1529.e5-e8 (Case rep.)
Altunkeser BB (*see* Can et al). 2006;19:1074.e1-e2 (Case rep.)
Ammer M (*see* Nesser et al). 2006;19:1286-92 (Case rep.)
Anavekar NS (*see* Thune et al). 2006;19:1462-5
Anavekar NS, Mirza A, Skali H, Plappert T, St John Sutton M, Pfeffer MA, Solomon SD. Risk assessment in patients with depressed left ventricular function after myocardial infarction using the myocardial performance index-survival and ventricular enlargement (SAVE) experience. 2006;19:28-33
Angel Garcia-Fernandez M (*see* Perez de Isla et al). 2006;19:1338-44
Anner H (*see* Khamaisi et al). 2006;19:835.e5-e8 (Case rep.)
Aqel R, Dobbs J, Lau Y, Lloyd S, Gupta H, Zoghbi GJ. Transjugular biopsy of a right atrial mass under intracardiac echocardiographic guidance. 2006;19:1072.e1-e3 (Case rep.)
Arai AE (*see* Sachdev et al). 2006;19:34-9
Aras D (*see* Ozeke et al). 2006;19:1245-50
Aras D (*see* Tufekcioglu et al). 2006;19:1320-5
Aravot D (*see* Shiran et al). 2006;19:578.e1-e4 (Case rep.)
Arezzi E (*see* Palmieri et al). 2006;19:491-8
Arita T (*see* Pohlel et al). 2006;19:229 (Case rep.)
Arnold JMO (*see* Thune et al). 2006;19:1462-5
Aronson D (*see* Dabbah et al). 2006;19:733-43
Aronson S (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)
Arsenault M (*see* Plante et al). 2006;19:1520-5
Asada K (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Asada-Kamiguchi J, Jones M, Greenberg NL, Popovic ZB, Tsujino H, Zetts AD, Qin JX, Garcia MJ, Thomas JD, Shiota T. Intraventricular pressure gradients in left ventricular aneurysms determined by color M-mode Doppler method: An animal study. 2006;19:1112-8
Asanuma T (*see* Okuda et al). 2006;19:781-7
Asanuma T (*see* Otani et al). 2006;19:973-81
Ask P (*see* Ressner et al). 2006;19:154-64
Askew JW (*see* Lester et al). 2006;19:919-23
Aslanian N (*see* Kobal et al). 2006;19:1012-6
Atar S (*see* Kobal et al). 2006;19:1012-6
Avigni N (*see* Mele et al). 2006;19:1373-81
Aydin M (*see* Dogan et al). 2006;19:243-8
Aylward PE (*see* Perry et al). 2006;19:1257-63

B

Badano LP (*see* Tosoratti et al). 2006;19:1190.e5-e8 (Case rep.)
Bahlmann E, Schneider C, Vitali Serdoz L, Hoffmann-Riem M, Broemel T, Kuck K-H. Spontaneous retraction of an intramyocardial dissecting hemorrhage and multiple left ventricular thrombus formations in subacute myocardial infarction and antiphospholipid syndrome: A case report with long-term follow-up. 2006;19:578.e5-e8 (Case rep.)
Bailey KR (*see* Bergeron et al). 2006;19:772-6
Balbarini A (*see* Di Bello et al). 2006;19:1063-71
Ballo P (*see* Mondillo et al). 2006;19:1085-91
Banchs JE, Dawn B, Abdel-Latif A, Qureshi A, Agrawal N, Bouvette M, Stoddard MF. Acquired aortic cusp fusion after chronic left ventricular assist device support. 2006;19:1401.e1-e3 (Case rep.)
Bangalore S, Petre L, Herweg B, Sichrovsky T, Vragel S, Steinberg JS, Chaudhry FA. Cardioversion in patients with left ventricular thrombus is not associated with increased thromboembolic risk. 2006;19:438-40
Bara C, Ghodsizad A, Niehaus M, Makoui M, Piechaczek C, Martin U, Warnecke G, Karck M, Gams E, Klein HM, Haverich A, Ruhparwar A. In vivo echocardiographic imaging of transplanted human adult stem cells in the myocardium labeled with clinically applicable clinimacs nanoparticles. 2006;19:563-8
Bardo D (*see* Corsi et al). 2006;19:886-93
Barrea C (*see* Roman et al). 2006;19:1382-92
Barutcu I (*see* Karakaya et al). 2006;19:434-7
Basic D, Siu SC, Skyba DM, Sloggett C, Jamorski M, Iwanochko RM, Yu EHC. Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested or known ischemic heart disease. 2006;19:1203-10
Basmadjian AJ (*see* Gisbert et al). 2006;19:140-6
Battagliese A (*see* De Luca et al). 2006;19:172-7
Bauer WR (*see* Weidemann et al). 2006;19:839-47
Baumann G (*see* Eddicks et al). 2006;19:1073.e7-e9 (Case rep.)
Baysal E (*see* Cay et al). 2006;19:1286-92 (Case rep.)
Baysan O (*see* Genc et al). 2006;19:940 (Letter)
Bednarz JE (*see* Sugeng et al). 2006;19:413-21

Indexes to the Seventeenth Annual Scientific Sessions abstracts are located in the May issue.

*January, pp. 1-112; February, pp. 121-242; March, pp. 243-364; April, pp. 365-470; May, pp. 471-732; June, pp. 733-838; July, pp. 839-942; August, pp. 943-1076; September, pp. 1077-1202; October, pp. 1203-1302; November, pp. 1303-1412; December, pp. 1413-1580.

Belcik T (*see* Kalvaitis et al). 2006;19:1488-93
Belcik TA (*see* Micari et al). 2006;19:379-85
Belham M, Kruger A, Pritchard C. The Tei index identifies a differential effect on left and right ventricular function with low-dose anthracycline chemotherapy. 2006;19:206-10
Belohlavek M (*see* Kjaergaard et al). 2006;19:322-8
Belohlavek M (*see* Wang et al). 2006;19:1102-11
Benali T (*see* Meimoun et al). 2006;19:1220-8
Benedetti G (*see* De Luca et al). 2006;19:172-7
Benson DW (*see* Markham et al). 2006;19:865-71
Benvenuti LA (*see* Salemi et al). 2006;19:354 (Case rep.)
Beppu S (*see* Okuda et al). 2006;19:781-7
Beppu S (*see* Otani et al). 2006;19:973-81
Bercow NR (*see* Mihalatos et al). 2006;19:1158-64
Bergeron S, Møller JE, Bailey KR, Chen HH, Burnett JC, Pellikka PA. Exertional changes in circulating cardiac natriuretic peptides in patients with suggested coronary artery disease. 2006;19:772-6
Berkay SH (*see* Mihalatos et al). 2006;19:1158-64
Bertagna X (*see* Meune et al). 2006;19:1466-70
Bertherat J (*see* Meune et al). 2006;19:1466-70
Bezerra Lira-Filho E, Campos O, Lazaro Andrade J, Henrique Fischer C, Godoy Nunes C, Cavalcanti Lins A, Hata F, Carvalho AC. Thoracic aorta evaluation in patients with Takayasu's arteritis by transesophageal echocardiography. 2006;19:829-34
Biadi O (*see* Di Bello et al). 2006;19:1063-71
Biagini E (*see* Pedone et al). 2006;19:982-6
Biclar L (*see* Nair et al). 2006;19:924-31
Bierig SM, Ehler D, Knoll ML, Waggoner AD. American Society of Echocardiography Minimum Standards for the Cardiac Sonographer: A Position Paper. 2006;19:471-4 (AMERICAN SOCIETY OF ECHOCARDIOGRAPHY POSITION PAPER)
Bijnens B (*see* Weidemann et al). 2006;19:839-47
Bitigen A (*see* Karakaya et al). 2006;19:434-7
Bitterman H (*see* Shiran et al). 2006;19:578.e1-e4 (Case rep.)
Blanchard DG (*see* Malouf et al). 2006;19:464-7
Bogers AJJC (*see* van den Bosch et al). 2006;19:7-13
Bogers AJJC (*see* van den Bosch et al). 2006;19:815-21
Bogot N (*see* Khamaisi et al). 2006;19:835.e5-e8 (Case rep.)
Bolson EL (*see* Jung et al). 2006;19:802-10
Border WL (*see* Markham et al). 2006;19:865-71
Borges AC (*see* Eddicks et al). 2006;19:1073.e7-e9 (Case rep.)
Bos JM, Hagler DJ, Silvilairat S, Cabalka A, O'Leary P, Daniels O, Miller FA, Abraham TP. Right ventricular function in asymptomatic individuals with a systemic right ventricle. 2006;19:1033-7
Boshoff D (*see* Eyskens et al). 2006;19:994-1000
Bouchard D (*see* Gisbert et al). 2006;19:140-6
Bouchard T (*see* Mor-Avi et al). 2006;19:48-54
Bourgoun M (*see* Thune et al). 2006;19:1462-5
Bouvette M (*see* Banchs et al). 2006;19:1401.e1-e3 (Case rep.)
Bouzas A (*see* Peteiro et al). 2006;19:1229-37
Bouzas A (*see* Peteiro et al). 2006;19:894-901
Bouzas B (*see* Peteiro et al). 2006;19:894-901
Bowry R (*see* David et al). 2006;19:329-34
Bracke F (*see* Jansen et al). 2006;19:185-91
Brandão CM (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Bratos JL (*see* Pérez-Sanz et al). 2006;19:1402.e5-e8 (Case rep.)
Brecker SJD (*see* Sharma et al). 2006;19:266-73
Breithardt G (*see* Bruch et al). 2006;19:522-8
Breithardt G (*see* Bruch et al). 2006;19:95-101
Breitwieser C (*see* Eddicks et al). 2006;19:1073.e7-e9 (Case rep.)
Brenneman CL (*see* Sachdev et al). 2006;19:34-9
Brodin L (*see* Lind et al). 2006;19:255-65
Brodin L (*see* Yumi Hayashi et al). 2006;19:1092-101
Brodin L-A (*see* Nilsson et al). 2006;19:499-506
Brodin L-A (*see* Ressner et al). 2006;19:154-64
Broemel T (*see* Bahlmann et al). 2006;19:578.e5-e8 (Case rep.)
Brook MM (*see* Lai et al).
Brown AL (*see* Dong et al). 2006;19:1017-25
Bruch C, Rothenburger M, Gotzmann M, Sindermann J, Scheld HH, Breithardt G, Wichter T. Risk stratification in chronic heart failure: Independent and incremental prognostic value of echocardiography and brain natriuretic peptide and its N-terminal fragment. 2006;19:522-8
Bruch C, Stypmann J, Grude M, Gradaus R, Breithardt G, Wichter T. Left bundle branch block in chronic heart failure-impact on diastolic function, filling pressures, and B-type natriuretic peptide levels. 2006;19:95-101
Bruschi G (*see* Moreo et al). 2006;19:938 (Case rep.)
Buchbinder M (*see* Naqvi et al). 2006;19:147-53
Bulut M (*see* Karakaya et al). 2006;19:434-7
Burden L (*see* Lim et al). 2006;19:280-4
Burnett JC (*see* Bergeron et al). 2006;19:772-6
Burwash IG (*see* Peters et al). 2006;19:1530.e7-e11 (Case rep.)
Busse KL (*see* Gepner et al). 2006;19:1170-4

C

Cabalka A (*see* Bos et al). 2006;19:1033-7
Cabalka AK (*see* Vyas and Cabalka). 2006;19:1286-92 (Case rep.)
Caiani EG (*see* Corsi et al). 2006;19:886-93
Calabrò R (*see* Caputo et al). 2006;19:836.e1-e4 (Case rep.)
Calabrò R (*see* Dialetto et al). 2006;19:836.e5-e8 (Case rep.)
Calabrò R (*see* Pisacane et al). 2006;19:107 (Case rep.)
Caldi F (*see* Palmieri et al). 2006;19:483-90
Campbell DB (*see* Yang et al). 2006;19:1051-5
Campian ME (*see* Hardziyenka et al). 2006;19:1272-9
Campos Machado LJ (*see* Freire et al). 2006;19:1251-6
Campos O (*see* Bezerra Lira-Filho et al). 2006;19:829-34
Can I, Altunkeser BB, Yavas O, Duzenli A, Ozdemir K, Gok H. Transit thrombus entrapped in patent foramen ovale resolved without clinical embolic events. 2006;19:1074.e1-e2 (Case rep.)
Capaldo B (*see* Palmieri et al). 2006;19:848-56
Caputo S, Santoro G, Morelli C, Sarubbi B, D'Alto M, Russo MG, Calabrò R. Subtotal anomalous pulmonary venous connection and left cor triatriatum: A rare diagnosis in adulthood. 2006;19:836.e1-e4 (Case rep.)
Carasso S (*see* Kuperstein et al). 2006;19:811-4
Cardello FP, Yoon DA, Halligan RE Jr., Richter H. The falciform ligament in the echocardiographic diagnosis of ascites. 2006;19:1074.e3-e4 (Case rep.)
Carnesecchi C (*see* Mumoli et al). 2006;19:1286-92 (Case rep.)
Carrier M (*see* Gisbert et al). 2006;19:140-6
Carvalho AC (*see* Bezerra Lira-Filho et al). 2006;19:829-34
Caso P (*see* Pisacane et al). 2006;19:107 (Case rep.)
Castro-Beiras A (*see* Peteiro et al). 2006;19:1229-37
Castro-Beiras A (*see* Peteiro et al). 2006;19:894-901
Cavalcanti Lins A (*see* Bezerra Lira-Filho et al). 2006;19:829-34
Cay S, Tufekcioglu O, Ozturk S, Ulupinar H, Tuna F, Baysal E, Pac A. Left ventricular diverticulum with contractile function in an unusual site. 2006;19:1286-92 (Case rep.)
Cei M (*see* Mumoli et al). 2006;19:1286-92 (Case rep.)
Celenk MK (*see* Ozeke et al). 2006;19:1245-50
Celentano A (*see* Palmieri et al). 2006;19:491-8
Celentano A (*see* Palmieri et al). 2006;19:848-56
Chakravarti S (*see* Katz et al). 2006;19:1446-8
Chaliki HP (*see* Lester et al). 2006;19:919-23
Chambers CE (*see* Yang et al). 2006;19:1051-5
Chan J (*see* Jenkins et al). 2006;19:1119-28
Chan J, Jenkins C, Khafagi F, Du L, Marwick TH. What is the optimal clinical technique for measurement of left ventricular volume after myocardial infarction? A comparative study of 3-dimensional echocardiography, single photon emission computed tomography, and cardiac magnetic resonance imaging. 2006;19:192-201
Chandrasekaran K (*see* Yang et al). 2006;19:1136-43
Chang S-M (*see* Hsiao et al). 2006;19:507-14
Chang S-M (*see* Hsiao et al). 2006;19:822-8
Chaudhry FA (*see* Bangalore et al). 2006;19:438-40
Chaudhry FA (*see* Shah et al). 2006;19:939 (Case rep.)
Chauvel C (*see* Cohen et al). 2006;19:580 (Letter)

Chen HH (*see* Bergeron et al). 2006;19:772-6
Cheng S-M (*see* Tsai et al). 2006;19:579.e1-e3 (Case rep.)
Chew DP (*see* Perry et al). 2006;19:1257-63
Cho E-J (*see* Youn et al). 2006;19:373-8
Cho S-Y (*see* Ha et al). 2006;19:1314-9
Chohan R (*see* Rovner et al). 2006;19:872-9
Choi B-W (*see* Choi et al). 2006;19:1211-9
Choi D (*see* Choi et al). 2006;19:1211-9
Choi E-Y (*see* Ha et al). 2006;19:1314-9
Choi E-Y, Seo H-S, Park S, Kim H-J, Ahn J-A, Ko Y-G, Choi B-W, Kang S-M, Choi D, Ha J-W, Rim S-J, Jang Y, Chung N. Prediction of transmural extent of infarction with contrast echocardiographically derived index of myocardial blood flow and myocardial blood volume fraction: Comparison with contrast-enhanced magnetic resonance imaging. 2006;19:1211-9
Choi K-B (*see* Youn et al). 2006;19:373-8
Choi K-J (*see* Kang et al). 2006;19:178-84
Choi Y-S (*see* Sohn et al). 2006;19:777-80
Chu C-S (*see* Su et al). 2006;19:274-9
Chu K-M (*see* Tsai et al). 2006;19:579.e1-e3 (Case rep.)
Chung N (*see* Choi et al). 2006;19:1211-9
Chung N (*see* Ha et al). 2006;19:1314-9
Chung T, Emmett L, Khoury V, Lau GT, Elsik M, Foo F, Allman KC, Kritharides L. Atrial and ventricular echocardiographic correlates of the extent of pulmonary embolism in the elderly. 2006;19:347-53
Chung W-S (*see* Youn et al). 2006;19:373-8
Citro R, Galderisi M, Maione A, Innelli P, Provenza G, Gregorio G. Sequential transthoracic ultrasound assessment of coronary flow reserve in a patient with tako-tsubo syndrome. 2006;19:1402.e5-e8 (Case rep.)
Claus P (*see* Eyskens et al). 2006;19:994-1000
Clayson S (*see* Horton et al). 2006;19:835.e1-e5 (Case rep.)
Coggins TR (*see* Main et al). 2006;19:1038-44
Coggins TR (*see* Main et al). 2006;19:64-70
Coglitore S (*see* de Gregorio et al). 2006;19:1190.e1-e4 (Case rep.)
Cohen A, Chauvel C, Abergel E, Raffoul H, Diebold B. Pulmonary regurgitant flow and detection of dip plateau. 2006;19:580 (Letter)
Cohen MS (*see* Seliem et al). 2006;19:121-9
Cokkinos P (*see* Paraskevaidis et al). 2006;19:249-54
Cokkinos P (*see* Paraskevaidis et al). 2006;19:529-35
Coles J (*see* Roman et al). 2006;19:1382-92
Collinson PO (*see* Sharma et al). 2006;19:266-73
Comenale Pinto S (*see* Pisacane et al). 2006;19:107 (Case rep.)
Coon P (*see* Corsi et al). 2006;19:886-93
Coon P (*see* Sugeng et al). 2006;19:413-21
Coon PD, Lang RM. Improved visualization of sinus venosus atrial septal defects in adults from the transthoracic approach. 2006;19:1072.e1-e3 (Case rep.)
Cordovil A, Fischer CH, Rodrigues ACT, Lira Filho EB, Vieira MLC, Cury AF, Naccarato GAF, Valente C, Brandão CM, Pommerantzeff PM, Morhy SS. Papillary muscle rupture after blunt chest trauma. 2006;19:464-7 (Case rep.)
Corsi C, Coon P, Goonewardena S, Weinert L, Sugeng L, Polonsky TS, Veronesi F, Caiani EG, Lamberti C, Bardo D, Lang RM, Mor-Avi V. Quantification of regional left ventricular wall motion from real-time 3-dimensional echocardiography in patients with poor acoustic windows: Effects of contrast enhancement tested against cardiac magnetic resonance. 2006;19:886-93
Coselli J (*see* Ramasubbu et al). 2006;19:464-7 (Case rep.)
Costa SP (*see* Olson et al). 2006;19:83-7
Cotrufo M (*see* Dialetto et al). 2006;19:836.e5-e8 (Case rep.)
Couet J (*see* Plante et al). 2006;19:1520-5
Couture P (*see* Gisbert et al). 2006;19:140-6
Covino FE (*see* Dialetto et al). 2006;19:836.e5-e8 (Case rep.)
Cripe LH (*see* Markham et al). 2006;19:865-71
Crone RA (*see* Mitchell and Crone). 2006;19:1190.e5-e8 (Case rep.)
Crooke GA (*see* Ahmed et al). 2006;19:938 (Case rep.)
Cui W (*see* Roberson et al). 2006;19:1175-81
Cui W, Roberson DA. Left ventricular TEI index in children: Comparison of tissue Doppler imaging, pulsed wave Doppler, and M-Mode echocardiography normal values. 2006;19:1438-45
Cuneo BF (*see* Roberson et al). 2006;19:1175-81
Cury AF (*see* Cordovil et al). 2006;19:464-7 (Case rep.)

D

Dabbah S, Reisner SA, Aronson D, Agmon Y. Left ventricular filling hemodynamics in patients with pulmonary edema and preserved versus reduced left ventricular ejection fraction: A prospective Doppler echocardiographic study. 2006;19:733-43
Daimon M (*see* Song et al). 2006;19:1129-35
Dalamaga EG (*see* Karvounis et al). 2006;19:1150-7
Dalirrooyfard M (*see* Sadeghpour et al). 2006;19:1073.e1-e5 (Case rep.)
Dall'Armellina E (*see* Tosoratti et al). 2006;19:1190.e5-e8 (Case rep.)
D'Alto M (*see* Caputo et al). 2006;19:836.e1-e4 (Case rep.)
Damus P (*see* Mihalatos et al). 2006;19:1158-64
Daniels BK (*see* Gilman et al). 2006;19:1526-8
Daniels O (*see* Bos et al). 2006;19:1033-7
Danik S (*see* Jang et al). 2006;19:1073.e7-e9 (Case rep.)
Dark JH (*see* Zaglavara et al). 2006;19:300-6
Das B (*see* Dyer et al). 2006;19:21-7
Das B (*see* Dyer et al). 2006;19:403-12
David J-S, Tousignant CP, Bowry R. Tricuspid annular velocity in patients undergoing cardiac operation using transesophageal echocardiography. 2006;19:329-34
Davidson WR Jr., (*see* Yang et al). 2006;19:1051-5
Davies RA (*see* Peters et al). 2006;19:1530.e7-e11 (Case rep.)
Dávila-Román VG (*see* Dong et al). 2006;19:1017-25
Dávila-Román VG (*see* Peterson et al). 2006;19:749-55
Dávila-Román VG (*see* Rovner et al). 2006;19:872-9
Dávila-Román VG (*see* Waggoner et al). 2006;19.307-13
Davis R (*see* McCulloch et al). 2006;19:744-8
Davogg K (*see* Nesser et al). 2006;19:1286-92 (Case rep.)
Dawn B (*see* Banchs et al). 2006;19:1401.e1-e3 (Case rep.)
Dawn B (*see* Siddiqui et al). 2006;19:1144-9
De Chiara B (*see* Moreo et al). 2006;19:938 (Case rep.)
de Gregorio C, Recupero A, Grimaldi P, Coglitore S. Can transthoracic live 3-dimensional echocardiography improve the recognition of midventricular obliteration in hypertrophic obstructive cardiomyopathy?. 2006;19:1190.e1-e4 (Case rep.)
De Gregorio P (*see* Pisacane et al). 2006;19:107 (Case rep.)
de las Fuentes L (*see* Dong et al). 2006;19:1017-25
de las Fuentes L (*see* Rovner et al). 2006;19:872-9
de las Fuentes L (*see* Waggoner et al). 2006;19:307-13
De Luca L, Sardella G, Proietti P, Battagliese A, Benedetti G, Di Roma A, Fedele F. Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: A Doppler echocardiographic study. 2006;19:172-7
De Pasquale CG (*see* Perry et al). 2006;19:1257-63
Debska-Slizien A (*see* Fijalkowski et al). 2006;19:1359-64
DeCastro S (*see* Maddukuri et al). 2006;19:1026-32
DeGroff C. Response to: "Echocardiographic assessment of isolated pulmonary valve stenosis: Which outpatient Doppler gradient has the most clinical validity?" 2006;19:364 (Letter)
Della Corte A (*see* Dialetto et al). 2006;19:836.e5-e8 (Case rep.)
Delle Donne MG (*see* Di Bello et al). 2006;19:1063-71
DeMaria AN (*see* Malouf et al). 2006;19:464-7
Demkow M (*see* Janus et al). 2006;19:1191.e5-e7 (Case rep.)
Denault AY (*see* Gisbert et al). 2006;19:140-6
Deveci B (*see* Ozeke et al). 2006;19:1245-50
Di Bello V, Santini F, Di Cori A, Pucci A, Palagi C, Delle Donne MG, Fierabracci P, Marsili A, Talini E, Giannetti M, Biadi O, Balbarini A, Mariani M, Pinchera A. Obesity cardiomyopathy: Is

it a reality? An ultrasonic tissue characterization study. 2006; 19:1063-71
Di Cori A (*see* Di Bello et al). 2006;19:1063-71
Di Minno G (*see* Palmieri et al). 2006;19:848-56
Di Roma A (*see* De Luca et al). 2006;19:172-7
Di Segni E (*see* Kuperstein et al). 2006;19:811-4
Di Tullio MR (*see* Pulerwitz et al). 2006;19:540-5
Di Valentino M, Menafoglio A, Mazzucchelli L, Siclari F, Gallino A. Rapid-growing left intraventricular cardiac hemangioma. 2006; 19:939 (Case rep.)
Dialetto G, Covino FE, Scognamiglio G, Della Corte A, Manduca S, Iacono C, Calabrò R, Cotrufo M. A rare complication of atrial septal occluders: Diagnosis by transthoracic echocardiography. 2006;19:836.e5-e8 (Case rep.)
Diebold B (*see* Cohen et al). 2006;19:580 (Letter)
Dijkmans PA, Knaapen P, Sieswerda GTJ, Aiazian E, Visser CA, Lammertsma AA, Visser FC, Kamp O. Quantification of myocardial perfusion using intravenous myocardial contrast echocardiography in healthy volunteers: Comparison with positron emission tomography. 2006;19:285-93
do Carmo Pereira Nunes M (*see* Freire et al). 2006;19:1251-6
Dobbs J (*see* Aqel et al). 2006;19:1072.e1-e3 (Case rep.)
Doddamani S (*see* Ahmed et al). 2006;19:938 (Case rep.)
Doddamani S (*see* Vennalaganti et al). 2006;19:1286-92 (Case rep.)
Dodouras T (*see* Paraskevaidis et al). 2006;19:529-35
Dogan S, Aydin M, Gursurer M, Dursun A, Onuk T, Madak H. Prediction of subclinical left ventricular dysfunction with strain rate imaging in patients with mild to moderate rheumatic mitral stenosis. 2006;19:243-8
Dohi K (*see* Yamanaka-Funabiki et al). 2006;19:1332-7
Dohi K, Pinsky MR, Kanzaki H, Severyn D, Gorcsan J III,. Effects of radial left ventricular dyssynchrony on cardiac performance using quantitative tissue Doppler radial strain imaging. 2006; 19:475-82
Dokainish H (*see* McCulloch and Dokainish). 2006;19:1411 (Letter)
Dokainish H (*see* McCulloch et al). 2006;19:744-8
Domanski MJ, Nanda N. Echocardiographic determination of left ventricular function. 2006;19:941 (Editorial)
Dong S-J, de las Fuentes L, Brown AL, Waggoner AD, Ewald GA, Dávila-Román VG. N-terminal pro B-type natriuretic peptide levels: Correlation with echocardiographically determined left ventricular diastolic function in an ambulatory cohort. 2006; 19:1017-25
Douglas PS (*see* Neilan et al). 2006;19:1280-5
Dousset B (*see* Meune et al). 2006;19:1466-70
Dowling RD (*see* Silverstein et al). 2006;19:1529.e5-e8 (Case rep.)
Drant S (*see* Peterson et al). 2006;19:1393-400
Drolet M-C (*see* Plante et al). 2006;19:1520-5
Dror Z (*see* Kuperstein et al). 2006;19:811-4
Du L (*see* Chan et al). 2006;19:192-201
Duboc D (*see* Meune et al). 2006;19:1466-70
Ducharme A (*see* Gisbert et al). 2006;19:140-6
Dursun A (*see* Dogan et al). 2006;19:243-8
Duzenli A (*see* Can et al). 2006;19:1074.e1-e2 (Case rep.)
Dwivedi G (*see* Lim et al). 2006;19:280-4
Dwivedi G, Janardhanan R, Hayat S, Senior R. Detection of residual infarct-related coronary artery stenosis and multivessel disease after thrombolysis: Comparison between myocardial contrast echocardiography and single photon emission computed tomography. 2006;19:546-51
Dyer K, Lanning C, Das B, Lee P-F, Ivy DD, Valdes-Cruz L, Shandas R. Noninvasive Doppler tissue measurement of pulmonary artery compliance in children with pulmonary hypertension. 2006;19:403-12
Dyer KL, Pauliks LB, Das B, Shandas R, Ivy D, Shaffer EM, Valdes-Cruz LM. Use of myocardial performance index in pediatric patients with idiopathic pulmonary arterial hypertension. 2006;19:21-7

E

Ebihara A (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Eddicks S, Kivelitz D, Breitwieser C, Knebel F, Theres H, Baumann G, Borges AC. Right ventricular metastasis caused by a renal cell carcinoma (Grawitz tumor): Case report. 2006;19: 1073.e7-e9 (Case rep.)
Edibam C, Playford D, Texler M, Edwards M. Isolated left atrial amyloidosis: Acute premitral stenosis secondary to spontaneous intramural left atrial hemorrhagic dissection. 2006;19: 938.e1-e4 (Case rep.)
Edwards M (*see* Edibam et al). 2006;19:938.e1-e4 (Case rep.)
Ehler D (*see* Bierig et al). 2006;19:471-4 (AMERICAN SOCIETY OF ECHOCARDIOGRAPHY POSITION PAPER)
Ehlgen A (*see* Main et al). 2006;19:1038-44
Eidem BW (*see* Havemann et al). 2006;19:355 (Case rep.)
El-Chami MF, Martin RP, Lerakis S. Esophageal dissection complicating transesophageal echocardiogram—the lesson to be learned: Do not force the issue. 2006;19:579.e5-e7 (Case rep.)
Elhassan HHM (*see* Ozeke et al). 2006;19:1245-50
Elhendy A, Tsutsui JM, O'Leary EL, Xie F, Porter TR. Noninvasive diagnosis of coronary artery bypass graft disease by dobutamine stress real-time myocardial contrast perfusion imaging. 2006;19:1482-7
Elliott PM (*see* Sharma et al). 2006;19:354 (Case rep.)
Elnoamany M (*see* Mahfouz Badran and Elnoamany). 2006;19: 1471-81
Elsik M (*see* Chung et al). 2006;19:347-53
Emmett L (*see* Chung et al). 2006;19:347-53
Engel DJ, Muratore R, Hirata K, Otsuka R, Fujikura K, Sugioka K, Marboe C, Lizzi FL, Homma S. Myocardial lesion formation using high-intensity focused ultrasound. 2006;19:932-7
Enseleit F, Reho I, Largiadèr T, Jenni R. Continuous wave Doppler signal: A mystery. 2006;19:1190.e5-e8 (Case rep.)
Eren M (*see* Uslu et al). 2006;19:341-6
Ergun K (*see* Ozeke et al). 2006;19:1245-50
Eriksson M (*see* Lind et al). 2006;19:255-65
Ertl G (*see* Weidemann et al). 2006;19:839-47
Esen AM (*see* Karakaya et al). 2006;19:434-7
Esmailzadeh M (*see* Sadeghpour et al). 2006;19:1073.e1-e5 (Case rep.)
Espinola-Zavaleta N (*see* Muñóz-Castellanos et al). 2006;19:1182-9
Ewald GA (*see* Dong et al). 2006;19:1017-25
Ewert GD (*see* Kean et al). 2006;19:202-5
Ewing S (*see* Seliem et al). 2006;19:121-9
Eyskens B, Ganame J, Claus P, Boshoff D, Gewillig M, Mertens L. Ultrasonic strain rate and strain imaging of the right ventricle in children before and after percutaneous closure of an atrial septal defect. 2006;19:994-1000

F

Faddis MN (*see* Waggoner et al). 2006;19:307-13
Fahmy Elnoamany M, Mahfouz Badran H, Helmy Abo Elazm T, Shawky Abdelaziz E. Asynchrony of left ventricular systolic performance after the first acute myocardial infarction in patients with narrow QRS complexes: Doppler tissue imaging study. 2006;19:1449-57
Fallon J (*see* Webber et al). 2006;19:1530.e1-e3 (Case rep.)
Farrell PE Jr., (*see* Seliem et al). 2006;19:121-9
Fedec A (*see* Mercer-Rosa et al). 2006;19:1511-9
Fedec A (*see* Seliem et al). 2006;19:121-9
Fedele F (*see* De Luca et al). 2006;19:172-7
Federici L (*see* Mele et al). 2006;19:1373-81
Feigenbaum H. Editorial. 2006;19:1075 (Editorial)
Feinberg MS (*see* Kuperstein et al). 2006;19:811-4
Feinstein JA (*see* Friedberg et al). 2006;19:559-62
Ferrari R (*see* Mele et al). 2006;19:1373-81
Fidler F (*see* Weidemann et al). 2006;19:839-47
Fierabracci P (*see* Di Bello et al). 2006;19:1063-71
Fijalkowski M, Koprowski A, Gruchala M, Galaska R, Debska-Slizien A, Rogowski J, Rutkowski B, Rynkiewicz A. Effect of preload reduction by hemodialysis on myocardial ultrasonic

characterization, left atrial volume, and Doppler tissue imaging in patients with end-stage renal disease. 2006;19:1359-64
Finsterer, *Wien, Austria J* (*see* Stöllberger and Finsterer, *Wien, Austria). 2006;19:109-10 (Letter)*
Fioretti PM (*see* Tosoratti et al). 2006;19:1190.e5-e8 (Case rep.)
Fischer CH (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Florit J (*see* Perez de Isla et al). 2006;19:1338-44
Foo F (*see* Chung et al). 2006;19:347-53
Fragasso G (*see* Agricola et al). 2006;19:457-63
Francis J (*see* Sunil Roy et al). 2006;19:1402.e1-e4 (Case rep.)
Fraser CD (*see* Havemann et al). 2006;19:355 (Case rep.)
Frassani R (*see* Tosoratti et al). 2006;19:1190.e5-e8 (Case rep.)
Frassi F (*see* Picano et al). 2006;19:356-63 (Review article)
Freire CMV, do Carmo Pereira Nunes M, Melo Barbosa M, Ribeiro de Oliveira Longo J, Impeliziere Nogueira A, Santos Assreuy Diniz S, Campos Machado LJ, Ribeiro de Oliveira A Jr.,. Gestational diabetes: A condition of early diastolic abnormalities in young women. 2006;19:1251-6
Friedberg MK, Feinstein JA, Rosenthal DN. A novel echocardiographic Doppler method for estimation of pulmonary arterial pressures. 2006;19:559-62
Friedberg MK, Silverman NH. The systolic to diastolic duration ratio in children with heart failure secondary to restrictive cardiomyopathy. 2006;19:1326-31
Frigstad S (*see* Malm et al). 2006;19:1494-501
Frigstad S (*see* Malm et al). 2006;19:365-72
Frigstad S (*see* Malm et al). 2006;19:40-7
Frommelt PC (*see* Lai et al).
Fuentes L (*see* Peterson et al). 2006;19:749-55
Fujii Y (*see* Okamoto et al). 2006;19:211-4
Fujikura K (*see* Engel et al). 2006;19:932-7
Fujimoto K (*see* Akiyama et al). 2006;19:55-63
Fujimoto K (*see* Hirata et al). 2006;19:165-71
Fukuda S (*see* Shin et al). 2006;19:796-801
Fukuda S (*see* Song et al). 2006;19:1129-35
Fulquet E (*see* Pérez-Sanz et al). 2006;19:1402.e5-e8 (Case rep.)
Fuyuki H (*see* Okura et al). 2006;19:552-8
Fye WB (*see* Gilman et al). 2006;19:1526-8

G

Gackowski A (*see* Janus et al). 2006;19:1191.e5-e7 (Case rep.)
Gaibazzi, *Piacenza, Italy N.* Asynchrony detection and Doppler tissue imaging. 2006;19:111 (Letter)
Galaska R (*see* Fijalkowski et al). 2006;19:1359-64
Galderisi M (*see* Citro et al). 2006;19:1402.e5-e8 (Case rep.)
Galderisi M (*see* Mondillo et al). 2006;19:1085-91
Galema TW (*see* Pedone et al). 2006;19:982-6
Gallino A (*see* Di Valentino et al). 2006;19:939 (Case rep.)
Gams E (*see* Bara et al). 2006;19:563-8
Ganame J (*see* Eyskens et al). 2006;19:994-1000
Ganame J (*see* Havemann et al). 2006;19:355 (Case rep.)
Garcia MJ (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Gardin JM (*see* Gerhard-Herman et al). 2006;19:955-72 (American Society of Echocardiography Report)
Gardin JM (*see* Roman et al). 2006;19:943-54 (American Society of Echocardiography Report)
Gargani L (*see* Picano et al). 2006;19:356-63 (Review article)
Gaynor JW (*see* Mercer-Rosa et al). 2006;19:1511-9
Gaynor JW (*see* Seliem et al). 2006;19:121-9
Gaze DC (*see* Sharma et al). 2006;19:266-73
Ge S (*see* Pemberton et al). 2006;19:1403-10 (Review article)
Geleijnse ML, Nemes A, Vletter WB. Response to: "Contrast-enhanced real-time 3-dimensional dobutamine stress echocardiography." 2006;19:1076 (Letter)
Genc C, Uzun M, Baysan O. Reply to "Angle correction for proximal isovelocity surface area method: Is it spheric cap or lune?" 2006;19:940 (Letter)
Gepner AD, Keevil JG, Wyman RA, Korcarz CE, Aeschlimann SE, Busse KL, Stein JH. Use of carotid intima-media thickness and vascular age to modify cardiovascular risk prediction. 2006;19:1170-4
Gepner AD, Korcarz CE, Aeschlimann SE, LeCaire TJ, Palta M, Tzou WS, Stein JH. Validation of a carotid intima-media thickness border detection program for use in an office setting. 2006;19:223-8
Gerhard-Herman M (*see* Roman et al). 2006;19:943-54 (American Society of Echocardiography Report)
Gerhard-Herman M, Gardin JM, Jaff M, Mohler E, Roman M, Naqvi T. Guidelines for noninvasive vascular laboratory testing: A report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology. 2006;19:955-72 (American Society of Echocardiography Report)
Gersh BJ (*see* Sorajja et al). 2006;19:788-95
Geva T (*see* Lai et al).
Geva T (*see* Wald et al). 2006;19:939.e4-36 (Case rep.)
Gewillig M (*see* Eyskens et al). 2006;19:994-1000
Ghali JK (*see* Thune et al). 2006;19:1462-5
Ghodsizad A (*see* Bara et al). 2006;19:563-8
Giannetti M (*see* Di Bello et al). 2006;19:1063-71
Gilman G, Lutzi CA, Daniels BK, Springer RF, Fye WB. The architecture of a mobile outreach echocardiography service. 2006;19:1526-8
Gilman S (*see* Kuperstein et al). 2006;19:811-4
Gin P-L (*see* Hsiao et al). 2006;19:902-10
Gin P-L, Wang W-C, Yang S-H, Hsiao S-H, Tseng J-C. Right heart function in systemic lupus erythematosus: Insights from myocardial Doppler tissue imaging. 2006;19:441-9
Gisbert A, Soulière V, Denault AY, Bouchard D, Couture P, Pellerin M, Carrier M, Levesque S, Ducharme A, Basmadjian AJ. Dynamic quantitative echocardiographic evaluation of mitral regurgitation in the operating department. 2006;19:140-6
Glas K (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)
Gleva MJ (*see* Waggoner et al). 2006;19:307-13
Gligorova S (*see* Picano et al). 2006;19:356-63 (Review article)
Godoy Nunes C (*see* Bezerra Lira-Filho et al). 2006;19:829-34
Gok H (*see* Can et al). 2006;19:1074.e1-e2 (Case rep.)
Golbasi Z (*see* Ozeke et al). 2006;19:1245-50
Goldman JH (*see* Main et al). 2006;19:1038-44
Goldman M (*see* Jang et al). 2006;19:1073.e7-e9 (Case rep.)
Gomberg-Maitland M (*see* Jacobs et al). 2006;19:464-7 (Case rep.)
Gonzales M (*see* Horton et al). 2006;19:835.e1-e5 (Case rep.)
Goonewardena S (*see* Corsi et al). 2006;19:886-93
Goonewardena S (*see* Jacobs et al). 2006;19:464-7 (Case rep.)
Gopal AS (*see* Mihalatos et al). 2006;19:1158-64
Gopal AS (*see* Mihalatos et al). 2006;19:76-82
Gopal AS (*see* Premkumar et al). 2006;19:107 (Case rep.)
Gorcsan J III, (*see* Dohi et al). 2006;19:475-82
Gordon GM (*see* Ahmed et al). 2006;19:938 (Case rep.)
Gordon GM (*see* Vennalaganti et al). 2006;19:1286-92 (Case rep.)
Gorgulu S (*see* Uslu et al). 2006;19:341-6
Gosavi S, Khaja A, Aggarwal K. Syncope with a twist. 2006;19:1529.e1-e3 (Case rep.)
Goto K, Mikami T, Onozuka H, Kaga S, Inoue M, Komatsu H, Komuro K, Yamada S, Tsutsui H, Kitabatake A. Role of left ventricular regional diastolic abnormalities for global diastolic dysfunction in patients with hypertrophic cardiomyopathy. 2006;19:857-64
Gotzmann M (*see* Bruch et al). 2006;19:522-8
Gradaus R (*see* Bruch et al). 2006;19:95-101
Greenberg NL (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Gregorio G (*see* Citro et al). 2006;19:1402.e5-e8 (Case rep.)
Gregson H (*see* Sharma et al). 2006;19:266-73
Grimaldi P (*see* de Gregorio et al). 2006;19:1190.e1-e4 (Case rep.)
Grimson R (*see* Mihalatos et al). 2006;19:1158-64
Grimson R (*see* Mihalatos et al). 2006;19:76-82
Gropler RJ (*see* Peterson et al). 2006;19:749-55
Gruchala M (*see* Fijalkowski et al). 2006;19:1359-64
Grude M (*see* Bruch et al). 2006;19:95-101
Guerra V (*see* Nii et al). 2006;19:391-402
Guerra V (*see* Takahashi et al). 2006;19:1502-10
Gupta A (*see* Leeper et al). 2006;19:1191.e5-e7 (Case rep.)
Gupta H (*see* Aqel et al). 2006;19:1072.e1-e3 (Case rep.)

Gursurer M (*see* Dogan et al). 2006;19:243-8
Gurudevan S (*see* Malouf et al). 2006;19:464-7

H

Ha J-W (*see* Choi et al). 2006;19:1211-9
Ha J-W, Ahn J-A, Kim J-M, Choi E-Y, Kang S-M, Rim S-J, Jang Y, Shim W-H, Cho S-Y, Oh JK, Chung N. Abnormal longitudinal myocardial functional reserve assessed by exercise tissue Doppler echocardiography in patients with hypertrophic cardiomyopathy. 2006;19:1314-9
Haaverstad R (*see* Zaglavara et al). 2006;19:300-6
Hagen ME (*see* Hillis et al). 2006;19:450-6
Hagler DJ (*see* Bos et al). 2006;19:1033-7
Halligan RE Jr., (*see* Cardello et al). 2006;19:1074.e3-e4 (Case rep.)
Halpern EF (*see* Neilan et al). 2006;19:1280-5
Hamilton JRL (*see* Zaglavara et al). 2006;19:300-6
Hammond E (*see* Horton et al). 2006;19:835.e1-e5 (Case rep.)
Han Y, Yeon S. Midventricular hypokinesis as a cardiac manifestation of anaphylaxis: A case report. 2006;19:1529.e9-e11 (Case rep.)
Hanekom L (*see* Jenkins et al). 2006;19:1119-28
Hannen MN (*see* Main et al). 2006;19:64-70
Haramati LB (*see* Ahmed et al). 2006;19:938 (Case rep.)
Hardziyenka M, Campian ME, Rianne de Bruin-Bon HACM, Michel MC, Tan HL. Sequence of echocardiographic changes during development of right ventricular failure in rat. 2006;19:1272-9
Hasan A (*see* Zaglavara et al). 2006;19:300-6
Hasegawa T (*see* Iwahashi et al). 2006;19:1238-44
Hashimoto M (*see* Okamoto et al). 2006;19:211-4
Hassager C (*see* Kjaergaard et al). 2006;19:1264-71
Hassager C (*see* Kjaergaard et al). 2006;19:1301 (Letter)
Hassager C (*see* Kjaergaard et al). 2006;19:1365-72
Hassager C (*see* Kjaergaard et al). 2006;19:322-8
Hassager C (*see* Kjaergaard et al). 2006;19:515-21
Hata F (*see* Bezerra Lira-Filho et al). 2006;19:829-34
Havemann L, McMahon CJ, Ganame J, Price J, Fraser CD, Eidem BW, Pignatelli RH. Rapid ventricular remodeling with left ventricular unloading postventricular assist device placement: New insights with strain imaging. 2006;19:355 (Case rep.)
Haverich A (*see* Bara et al). 2006;19:563-8
Hayat S (*see* Dwivedi et al). 2006;19:546-51
Hedberg P, Jonason T, Lönnberg I, Nilsson G, Pehrsson K, Ringqvist I. Mitral annulus motion as a predictor of mortality in a community-based sample of 75-year-old men and women. 2006;19:88-94
Helbing WA (*see* van den Bosch et al). 2006;19:1-6
Helmy Abo Elazm T (*see* Fahmy Elnoamany et al). 2006;19:1449-57
Henningsson R (*see* Nilsson et al). 2006;19:499-506
Henrique Fischer C (*see* Bezerra Lira-Filho et al). 2006;19:829-34
Hentz JG (*see* Lester et al). 2006;19:919-23
Herweg B (*see* Bangalore et al). 2006;19:438-40
Herz S (*see* Pulerwitz et al). 2006;19:540-5
Hesse B (*see* Ulrich et al). 2006;19:1165-9
Heywood JT (*see* Varadarajan et al). 2006;19:1458-61
Hillis GS, Ujino K, Mulvagh SL, Hagen ME, Oh JK. Echocardiographic indices of increased left ventricular filling pressure and dilation after acute myocardial infarction. 2006;19:450-6
Hirano T (*see* Okuda et al). 2006;19:781-7
Hirano Y, Uehara H, Nakamura H, Ikuta S, Nakano M, Akiyama S, Ishikawa K. Efficacy of ultrasound-assisted stress testing using a hand-carried ultrasound device for diagnosis of coronary artery disease. 2006;19:536-9
Hirata K (*see* Engel et al). 2006;19:932-7
Hirata K (*see* Pulerwitz et al). 2006;19:540-5
Hirata K, Watanabe H, Otsuka R, Fujimoto K, Tokai K, Yamagishi H, Yoshiyama M, Yoshikawa J. Noninvasive diagnosis of restenosis by transthoracic Doppler echocardiography after percutaneous coronary intervention: Comparison with exercise Tl-SPECT. 2006;19:165-71
Hoff L (*see* Ressner et al). 2006;19:154-64
Hoffmann-Riem M (*see* Bahlmann et al). 2006;19:578.e5-e8 (Case rep.)
Homma S (*see* Engel et al). 2006;19:932-7
Homma S (*see* LaRocca et al). 2006;19:464-7 (Case rep.)
Homma S (*see* Pulerwitz et al). 2006;19:540-5
Hong G-J (*see* Tsai et al). 2006;19:579.e1-e3 (Case rep.)
Hong S-J (*see* Youn et al). 2006;19:373-8
Horton SC, Nay S, Gonzales M, McIntosh M, Hammond E, Clayson S. Smooth muscle neoplasm presenting as intrapericardial myxoma. 2006;19:835.e1-e5 (Case rep.)
Hozumi T (*see* Takagi et al). 2006;19:1001-11
Hsiao S-H (*see* Gin et al). 2006;19:441-9
Hsiao S-H, Lee C-Y, Chang S-M, Lin S-K, Liu C-P. Right heart function in scleroderma: Insights from myocardial Doppler tissue imaging. 2006;19:507-14
Hsiao S-H, Lee C-Y, Chang S-M, Yang S-H, Lin S-K, Huang W-C. Pulmonary embolism and right heart function: Insights from myocardial Doppler tissue imaging. 2006;19:822-8
Hsiao S-H, Lin S-K, Wang W-C, Yang S-H, Gin P-L, Liu C-P. Severe tricuspid regurgitation shows significant impact in the relationship among peak systolic tricuspid annular velocity, tricuspid annular plane systolic excursion, and right ventricular ejection fraction. 2006;19:902-10
Huang W-C (*see* Hsiao et al). 2006;19:822-8
Humes RA (*see* Lai et al).
Hurst RT (*see* Lester et al). 2006;19:919-23
Huynh HK (*see* Naqvi and Huynh). 2006;19:569-77

I

Iaccarino M (*see* Palmieri et al). 2006;19:848-56
Iacono C (*see* Dialetto et al). 2006;19:836.e5-e8 (Case rep.)
Ibuki M (*see* Katayama et al). 2006;19:335-40
Ihm S-H (*see* Youn et al). 2006;19:373-8
Ikuta S (*see* Hirano et al). 2006;19:536-9
Iliodromitis EK (*see* Paraskevaidis et al). 2006;19:249-54
Impeliziere Nogueira A (*see* Freire et al). 2006;19:1251-6
Innelli P (*see* Citro et al). 2006;19:1402.e5-e8 (Case rep.)
Innocenti F (*see* Palmieri et al). 2006;19:483-90
Inoue M (*see* Goto et al). 2006;19:857-64
Ishikawa K (*see* Hirano et al). 2006;19:536-9
Ishikura F (*see* Okuda et al). 2006;19:781-7
Ishikura F (*see* Otani et al). 2006;19:973-81
Ito M (*see* Yamanaka-Funabiki et al). 2006;19:1332-7
Ito T, Suwa M, Tonari S, Okuda N, Kitaura Y. Regional postsystolic shortening in patients with hypertrophic cardiomyopathy: Its incidence and characteristics assessed by strain imaging. 2006;19:987-93
Ivy D (*see* Dyer et al). 2006;19:21-7
Ivy DD (*see* Dyer et al). 2006;19:403-12
Iwahashi N, Nakatani S, Kanzaki H, Hasegawa T, Abe H, Kitakaze M. Acute improvement in myocardial function assessed by myocardial strain and strain rate after aortic valve replacement for aortic stenosis. 2006;19:1238-44
Iwamoto A (*see* Okamoto et al). 2006;19:211-4
Iwanochko RM (*see* Basic et al). 2006;19:1203-10
Iwata K (*see* Okura et al). 2006;19:552-8

J

Jaber WA (*see* Ulrich et al). 2006;19:1165-9
Jacobs A, Weinert LC, Goonewardena S, Gomberg-Maitland M, Lang RM. Three-Dimensional transthoracic echocardiography to evaluate cor triatriatum in the adult. 2006;19:464-7 (Case rep.)
Jacobs LD (*see* Mor-Avi et al). 2006;19:48-54
Jaeggi ET (*see* Nii et al). 2006;19:1431-7
Jaeggi ET (*see* Nii et al). 2006;19:314-21
Jaff M (*see* Gerhard-Herman et al). 2006;19:955-72 (American Society of Echocardiography Report)
Jaff M (*see* Roman et al). 2006;19:943-54 (American Society of Echocardiography Report)

Jamorski M (*see* Basic et al). 2006;19:1203-10
Janardhanan R (*see* Dwivedi et al). 2006;19:546-51
Janardhanan R (*see* Lim et al). 2006;19:280-4
Janerot-Sjoberg B (*see* Ressner et al). 2006;19:154-64
Jang JJ, Danik S, Goldman M. Primary cardiac lymphoma: Diagnosis and treatment guided by transesophageal echocardiogram perfusion imaging. 2006;19:1073.e7-e9 (Case rep.)
Jang Y (*see* Choi et al). 2006;19:1211-9
Jang Y (*see* Ha et al). 2006;19:1314-9
Jansen AHM, Bracke F, van Dantzig JM, Meijer A, Korsten EHM, Peels KH, van Hemel NM. Optimization of pulsed wave tissue Doppler to predict left ventricular reverse remodeling after cardiac resynchronization therapy. 2006;19:185-91
Jansson T (*see* Ressner et al). 2006;19:154-64
Janus B, Krol-Jawien W, Demkow M, Gackowski A, Klimeczek P, Moczulski Z. Pulmonary artery dissection: A rare complication of pulmonary balloon valvuloplasty diagnosed 11 years after the procedure. 2006;19:1191.e5-e7 (Case rep.)
Jassal DS (*see* Neilan et al). 2006;19:1280-5
Jassal DS, Qureshi A, Neilan TG, Nomura C, Wittram C, King MEE. Pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia: An echocardiographic perspective. 2006;19:229 (Case rep.)
Javois AJ (*see* Roberson et al). 2006;19:1175-81
Jenkins C (*see* Chan et al). 2006;19:192-201
Jenkins C, Chan J, Hanekom L, Marwick TH. Accuracy and feasibility of online 3-dimensional echocardiography for measurement of left ventricular parameters. 2006;19:1119-28
Jenni R (*see* Enseleit et al). 2006;19:1190.e5-e8 (Case rep.)
Jeon H-K (*see* Youn et al). 2006;19:373-8
Jerosch-Herold M (*see* Pemberton et al). 2006;19:1403-10 (Review article)
Jin Z (*see* Pulerwitz et al). 2006;19:540-5
Johnson BD (*see* Kjaergaard et al). 2006;19:515-21
Jolly N (*see* Sugeng et al). 2006;19:413-21
Jonason T (*see* Hedberg et al). 2006;19:88-94
Jones M (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Jones PG (*see* Main et al). 2006;19:64-70
Joseph MX (*see* Perry et al). 2006;19:1257-63
Joseph S (*see* Mihalatos et al). 2006;19:76-82
Joslin NB (*see* Khumri et al). 2006;19:1072.e1-e3 (Case rep.)
Jude N (*see* Meune et al). 2006;19:1466-70
Jung H-O (*see* Youn et al). 2006;19:373-8
Jung HO, Sheehan FH, Bolson EL, Waiss M-P, Otto CM. Evaluation of midwall systolic function in left ventricular hypertrophy: A comparison of 3-dimensional versus 2-dimensional echocardiographic indices. 2006;19:802-10
Juraszek AL (*see* Wald et al). 2006;19:939.e4-36 (Case rep.)

K

Kaga S (*see* Goto et al). 2006;19:857-64
Kaji S (*see* Akiyama et al). 2006;19:55-63
Kajihara K (*see* Okamoto et al). 2006;19:211-4
Kalvaitis S, Kaul S, Tong KL, Rinkevich D, Belcik T, Wei K. Effect of time delay on the diagnostic use of contrast echocardiography in patients presenting to the emergency department with chest pain and no S-T Segment elevation. 2006;19:1488-93
Kamp O (*see* Dijkmans et al). 2006;19:285-93
Kamp O (*see* Klein et al). 2006;19:763-71
Kang D-H (*see* Kang et al). 2006;19:178-84
Kang S-J, Song J-K, Song J-M, Kang D-H, Lee EY, Kim J, Nam G-B, Choi K-J, Kim J-J, Kim Y-H. Usefulness of ventricular longitudinal contractility assessed by Doppler tissue imaging in the prediction of reverse remodeling in patients with severe left ventricular systolic dysfunction. 2006;19:178-84
Kang S-M (*see* Choi et al). 2006;19:1211-9
Kang S-M (*see* Ha et al). 2006;19:1314-9
Kanzaki H (*see* Dohi et al). 2006;19:475-82
Kanzaki H (*see* Iwahashi et al). 2006;19:1238-44
Karakaya O, Turkmen M, Bitigen A, Saglam M, Barutcu I, Esen AM, Bulut M, Turkyilmaz E, Kirma C. Effect of percutaneous mitral balloon valvuloplasty on left atrial appendage function: A Doppler tissue study. 2006;19:434-7
Karamitsos TD (*see* Karvounis et al). 2006;19:1150-7
Karavolias G (*see* Paraskevaidis et al). 2006;19:529-35
Karck M (*see* Bara et al). 2006;19:563-8
Karkabi B (*see* Shiran et al). 2006;19:578.e1-e4 (Case rep.)
Karp K (*see* Lindqvist et al). 2006;19:1345-9
Karvounis HI (*see* Zaglavara et al). 2006;19:300-6
Karvounis HI, Dalamaga EG, Papadopoulos CE, Karamitsos TD, Vassilikos V, Paraskevaidis S, Styliadis IH, Parharidis GE, Louridas GE. Improved papillary muscle function attenuates functional mitral regurgitation in patients with dilated cardiomyopathy after cardiac resynchronization therapy. 2006;19:1150-7
Kasravi B (*see* Movahed et al). 2006;19:911-3
Kataoka T (*see* Tanaka et al). 2006;19:756-62
Katayama M, Yamamuro A, Ueda Y, Tamita K, Yamabe K, Ibuki M, Takagi T, Yagi T, Akasaka T, Morioka S. Coronary flow velocity pattern assessed noninvasively by transthoracic color Doppler echocardiography serves as a predictor of adverse cardiac events and left ventricular remodeling in patients with acute myocardial infarction. 2006;19:335-40
Kates R (*see* Mihalatos et al). 2006;19:1158-64
Katz DH (*see* Ahmed et al). 2006;19:938 (Case rep.)
Katz JC, Chakravarti S, Ko HH, Lytrivi ID, Srivastava S, Lai WW, Parness IA, Nguyen K, Nielsen JC. Common origin of the innominate and carotid arteries: Prevalence, nomenclature, and surgical implications. 2006;19:1446-8
Kaul S (*see* Kalvaitis et al). 2006;19:1488-93
Kaul S (*see* Micari et al). 2006;19:379-85
Kawai H (*see* Tanaka et al). 2006;19:756-62
Kawamoto T (*see* Kume et al). 2006;19:914-8
Kawamoto T (*see* Watanabe et al). 2006;19:71-5
Kean AJ, McCloskey VR, Seghatol FF, Ewert GD, Mehlman DJ, McPherson DD, Rigolin VH. Preservation of ventricular function in amateur athletes after completion of a marathon. 2006;19:202-5
Keevil JG (*see* Gepner et al). 2006;19:1170-4
Keirns C (*see* Muñóz-Castellanos et al). 2006;19:1182-9
Kenny A (*see* Zaglavara et al). 2006;19:300-6
Khafagi F (*see* Chan et al). 2006;19:192-201
Khaja A (*see* Gosavi et al). 2006;19:1529.e1-e3 (Case rep.)
Khamaisi M, Anner H, Marincheva G, Leitersdorf E, Bogot N. Stent graft in treatment of penetrating thoracic aortic ulcer. 2006;19:835.e5-e8 (Case rep.)
Khandheria BK (*see* Nesser et al). 2006;19:1286-92 (Case rep.)
Khandheria BK (*see* Yang et al). 2006;19:1136-43
Khoury PR (*see* Markham et al). 2006;19:865-71
Khoury V (*see* Chung et al). 2006;19:347-53
Khumri TM, Joslin NB, Nayyar S, Main ML. Transesophageal echocardiographic diagnosis of *Aspergillus fumigatus* aortitis after percutaneous coronary intervention. 2006;19:1072.e1-e3 (Case rep.)
Kim J-H (*see* Youn et al). 2006;19:373-8
Kim H-J (*see* Choi et al). 2006;19:1211-9
Kim H-K (*see* Sohn et al). 2006;19:777-80
Kim H-K, Sohn D-W, Lee S-E, Kim Y-J, Oh B-H, Park Y-B. Coronary blood flow after cold exposure and the effect of warm-up exercise. 2006;19:386-90
Kim H-S (*see* Sohn et al). 2006;19:777-80
Kim J (*see* Kang et al). 2006;19:178-84
Kim J-J (*see* Kang et al). 2006;19:178-84
Kim J-M (*see* Ha et al). 2006;19:1314-9
Kim K-Y (*see* Sohn et al). 2006;19:777-80
Kim Y-H (*see* Kang et al). 2006;19:178-84
Kim Y-J (*see* Kim et al). 2006;19:386-90
Kim Y-J (*see* Sohn et al). 2006;19:777-80
King MEE (*see* Jassal et al). 2006;19:229 (Case rep.)
Kingdom J (*see* Nii et al). 2006;19:1431-7
Kirma C (*see* Karakaya et al). 2006;19:434-7
Kisslo J (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)
Kitabatake A (*see* Goto et al). 2006;19:857-64

Kitakaze M (*see* Iwahashi et al). 2006;19:1238-44
Kitaura Y (*see* Ito et al). 2006;19:987-93
Kivelitz D (*see* Eddicks et al). 2006;19:1073.e7-e9 (Case rep.)
Kjaergaard J, Hassager C, Sogaard P. Response to comments on our article "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet". 2006;19:1301 (Letter)
Kjaergaard J, Korinek J, Belohlavek M, Oh JK, Sogaard P, Hassager C. Accuracy, reproducibility, and comparability of Doppler tissue imaging by two high-end ultrasound systems. 2006;19:322-8
Kjaergaard J, Krogsgaard Schaadt B, Lund JO, Hassager C. Quantitative measures of right ventricular dysfunction by echocardiography in the diagnosis of acute nonmassive pulmonary embolism. 2006;19:1264-71
Kjaergaard J, Snyder EM, Hassager C, Oh JK, Johnson BD. Impact of preload and afterload on global and regional right ventricular function and pressure: A quantitative echocardiography study. 2006;19:515-21
Kjaergaard J, Sogaard P, Hassager C. Quantitative echocardiographic analysis of the right ventricle in healthy individuals. 2006;19:1365-72
Klein HM (*see* Bara et al). 2006;19:563-8
Klein LJ, van Campen L, Sieswerda GT, Kamp O, Visser CA, Visser FC. Glucose-insulin-potassium echocardiography detects improved segmental myocardial function and viable tissue shortly after acute myocardial infarction. 2006;19:763-71
Klimeczek P (*see* Janus et al). 2006;19:1191.e5-e7 (Case rep.)
Knaapen P (*see* Dijkmans et al). 2006;19:285-93
Knebel F (*see* Eddicks et al). 2006;19:1073.e7-e9 (Case rep.)
Knoll ML (*see* Bierig et al). 2006;19:471-4 (AMERICAN SOCIETY OF ECHOCARDIOGRAPHY POSITION PAPER)
Ko HH (*see* Katz et al). 2006;19:1446-8
Ko Y-G (*see* Choi et al). 2006;19:1211-9
Kobal SL, Pollick C, Atar S, Miyamoto T, Aslanian N, Neuman Y, Tolstrup K, Naqvi TZ, Luo H, Macrum B, Siegel RJ. Stress echocardiography in octogenarians: Transesophageal atrial pacing is accurate, safe, and well tolerated. 2006;19:1012-6
Køber L (*see* Thune et al). 2006;19:1462-5
Koh L-P (*see* Seow et al). 2006;19:102-6
Kokumai M (*see* Nakai et al). 2006;19:880-5
Kokumai M (*see* Takeuchi et al). 2006;19:1077-84
Komatsu H (*see* Goto et al). 2006;19:857-64
Komuro K (*see* Goto et al). 2006;19:857-64
Komuro T (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Konstantinov I (*see* Nii et al). 2006;19:314-21
Koo B-K (*see* Sohn et al). 2006;19:777-80
Koprowski A (*see* Fijalkowski et al). 2006;19:1359-64
Korcarz CE (*see* Gepner et al). 2006;19:1170-4
Korcarz CE (*see* Gepner et al). 2006;19:223-8
Korcarz CE (*see* Tzou et al). 2006;19:1286-92
Korinek J (*see* Kjaergaard et al). 2006;19:322-8
Korinek J (*see* Wang et al). 2006;19:1102-11
Korsten EHM (*see* Jansen et al). 2006;19:185-91
Kort S. Intracardiac echocardiography: Evolution, recent advances, and current applications. 2006;19:1192-201 (Review article)
Kort S. Real-time 3-dimensional echocardiography for prosthetic valve endocarditis: Initial experience. 2006;19:130-9
Koyama Y (*see* Akiyama et al). 2006;19:55-63
Kremastinos DT (*see* Paraskevaidis et al). 2006;19:249-54
Krenning BJ (*see* van den Bosch et al). 2006;19:1-6
Krishnan MN (*see* Sunil Roy et al). 2006;19:1402.e1-e4 (Case rep.)
Kritharides L (*see* Chung et al). 2006;19:347-53
Krogsgaard Schaadt B (*see* Kjaergaard et al). 2006;19:1264-71
Krol-Jawien W (*see* Janus et al). 2006;19:1191.e5-e7 (Case rep.)
Kruger A (*see* Belham et al). 2006;19:206-10
Krumholz HM (*see* Mena et al). 2006;19:1295-300 (Review article)
Kubo T (*see* Okura et al). 2006;19:552-8
Kuck K-H (*see* Bahlmann et al). 2006;19:578.e5-e8 (Case rep.)
Kukar A (*see* Shah et al). 2006;19:939 (Case rep.)
Kukuzke JA (*see* Yang et al). 2006;19:1136-43
Kum LCC (*see* Zhang et al). 2006;19:422-8
Kume T, Kawamoto T, Akasaka T, Watanabe N, Toyota E, Neishi Y, Wada N, Okahashi N, Yoshida K. Rate of progression of valvular aortic stenosis in patients undergoing dialysis. 2006;19:914-8
Kuperstein R, Feinberg MS, Carasso S, Gilman S, Dror Z, Di Segni E. The added value of real-time 3-dimensional echocardiography in the diagnosis of isolated cleft mitral valve in adults. 2006;19:811-4
Kuri-Nivón M (*see* Muñóz-Castellanos et al). 2006;19:1182-9
Kusnetzky LL (*see* Main et al). 2006;19:64-70
Kuvin JT (*see* Maddukuri et al). 2006;19:1026-32
Kyrzopoulos S (*see* Paraskevaidis et al). 2006;19:249-54

L

Lachance D (*see* Plante et al). 2006;19:1520-5
Lagrand WK (*see* Nemes et al). 2006;19:355 (Case rep.)
Lai W-T (*see* Su et al). 2006;19:274-9
Lai WW (*see* Katz et al). 2006;19:1446-8
Lai WW (*see* Peterson et al). 2006;19:1393-400
Lai WW, Geva T, Shirali GS, Frommelt PC, Humes RA, Brook MM, Pignatelli RH, Rychik J. Guidelines and standards for performance of a pediatric echocardiogram: A report from the task force of the pediatric council of the american society of echocardiography.
Lam Y-Y (*see* Zhang et al). 2006;19:422-8
Lamberti C (*see* Corsi et al). 2006;19:886-93
Lamendola C (*see* Mihalatos et al). 2006;19:1158-64
Lammertin G (*see* Sugeng et al). 2006;19:413-21
Lammertsma AA (*see* Dijkmans et al). 2006;19:285-93
Lang RM (*see* Coon and Lang). 2006;19:1072.e1-e3 (Case rep.)
Lang RM (*see* Corsi et al). 2006;19:886-93
Lang RM (*see* Jacobs et al). 2006;19:464-7 (Case rep.)
Lang RM (*see* Mor-Avi et al). 2006;19:48-54
Lang RM (*see* Nakai et al). 2006;19:880-5
Lang RM (*see* Sugeng et al). 2006;19:413-21
Lang RM (*see* Takeuchi et al). 2006;19:1077-84
Lang RM (*see* Takeuchi et al). 2006;19:294-9
Lanning C (*see* Dyer et al). 2006;19:403-12
Lanza P (*see* Main et al). 2006;19:1038-44
Lanza P (*see* Main et al). 2006;19:64-70
Largiadèr T (*see* Enseleit et al). 2006;19:1190.e5-e8 (Case rep.)
LaRocca GM, Shimbo D, Rodriguez CJ, Stewart A, Naka Y, Weinberger J, Homma S, Pizzarello R. The impella recover LP 5.0 left ventricular assist device: A bridge to coronary artery bypass grafting and cardiac transplantation. 2006;19:464-7 (Case rep.)
Lau GT (*see* Chung et al). 2006;19:347-53
Lau Y (*see* Aqel et al). 2006;19:1072.e1-e3 (Case rep.)
Lavi N, Zelinger A, Silver M. Thrombotic "ghosts": Echocardiographic appearance of thrombi with hollow cores and implications regarding mechanism of spontaneous clot lysis. 2006;19:1530.e5-e6 (Case rep.)
Lavine JA (*see* Lavine and Lavine). 2006;19:1350-8
Lavine SJ, Lavine JA. The effect of acute hypertension on left ventricular diastolic pressures in a canine model of left ventricular dysfunction with a preserved ejection fraction and elevated left ventricular filling pressures. 2006;19:1350-8
Lazaro Andrade J (*see* Bezerra Lira-Filho et al). 2006;19:829-34
LeCaire TJ (*see* Gepner et al). 2006;19:223-8
Lee J-M (*see* Youn et al). 2006;19:373-8
Lee P-F (*see* Dyer et al). 2006;19:403-12
Lee C-Y (*see* Hsiao et al). 2006;19:507-14
Lee C-Y (*see* Hsiao et al). 2006;19:822-8
Lee EY (*see* Kang et al). 2006;19:178-84
Lee K-T (*see* Su et al). 2006;19:274-9
Lee P-W (*see* Zhang et al). 2006;19:422-8
Lee S-E (*see* Kim et al). 2006;19:386-90
Leeper NJ, Gupta A, Murphy DJ, Wu JC. Right coronary cameral

fistula resulting from surgery of double chamber right ventricle. 2006;19:1191.e5-e7 (Case rep.)
Leitersdorf E (*see* Khamaisi et al). 2006;19:835.e5-e8 (Case rep.)
Lemos P (*see* Salemi et al). 2006;19:354 (Case rep.)
Lerakis S (*see* El-Chami et al). 2006;19:579.e5-e7 (Case rep.)
Lerakis S (*see* Pohlel et al). 2006;19:229 (Case rep.)
Lester SJ, Askew JW, Hurst RT, Moreno CA, Hentz JG, Wood LA, Simpson RA, Schneck SL, Tyler R, Chaliki HP, Lynch JJ. Contrast echocardiography: Experience in a clinical echocardiography laboratory. 2006;19:919-23
Lever HM (*see* Song et al). 2006;19:1129-35
Levesque S (*see* Gisbert et al). 2006;19:140-6
Levi DS (*see* Peterson et al). 2006;19:1393-400
Levine RA (*see* Mele et al). 2006;19:1373-81
Levy F (*see* Meimoun et al). 2006;19:1220-8
Lewis BS (*see* Shiran et al). 2006;19:578.e1-e4 (Case rep.)
Li J (*see* Nii et al). 2006;19:314-21
Li M (*see* Ahmed et al). 2006;19:938 (Case rep.)
Lie GR (*see* Sachdev et al). 2006;19:34-9
Lim TK, Burden L, Janardhanan R, Dwivedi G, Ping C, Moon J, Pennell DJ, Senior R. Contrast echocardiography versus gated single photon emission computed tomography for the assessment of parameters of left ventricular remodeling after acute myocardial infarction. 2006;19:280-4
Lin H (*see* Zhang et al). 2006;19:422-8
Lin S-K (*see* Hsiao et al). 2006;19:507-14
Lin S-K (*see* Hsiao et al). 2006;19:822-8
Lin S-K (*see* Hsiao et al). 2006;19:902-10
Lin T-H (*see* Su et al). 2006;19:274-9
Lind B (*see* Yumi Hayashi et al). 2006;19:1092-101
Lind B, Eriksson M, Roumina S, Nowak J, Brodin L. Longitudinal isovolumic displacement of the left ventricular myocardium assessed by tissue velocity echocardiography in healthy individuals. 2006;19:255-65
Lindholm B (*see* Yumi Hayashi et al). 2006;19:1092-101
Lindner JR (*see* Micari et al). 2006;19:379-85
Lindqvist P, Mörner S, Karp K, Waldenström A. New aspects of septal function by using 1-Dimensional strain and strain rate imaging. 2006;19:1345-9
Lira Filho EB (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Liu C-P (*see* Hsiao et al). 2006;19:507-14
Liu C-P (*see* Hsiao et al). 2006;19:902-10
Livi U (*see* Tosoratti et al). 2006;19:1190.e5-e8 (Case rep.)
Lizzi FL (*see* Engel et al). 2006;19:932-7
Lloyd S (*see* Aqel et al). 2006;19:1072.e1-e3 (Case rep.)
Logan J (*see* Naqvi et al). 2006;19:147-53
Lönnberg I (*see* Hedberg et al). 2006;19:88-94
Louridas GE (*see* Karvounis et al). 2006;19:1150-7
Louridas GE (*see* Zaglavara et al). 2006;19:300-6
Lund JO (*see* Kjaergaard et al). 2006;19:1264-71
Luo H (*see* Kobal et al). 2006;19:1012-6
Lutzi CA (*see* Gilman et al). 2006;19:1526-8
Luycx-Bore A (*see* Meimoun et al). 2006;19:1220-8
Lynch JJ (*see* Lester et al). 2006;19:919-23
Lytrivi ID (*see* Katz et al). 2006;19:1446-8

M

Macgowan CK (*see* Nii et al). 2006;19:391-402
Macgowan CK (*see* Roman et al). 2006;19:1382-92
Macrum B (*see* Kobal et al). 2006;19:1012-6
Madak H (*see* Dogan et al). 2006;19:243-8
Madani M (*see* Malouf et al). 2006;19:464-7
Maddukuri PV, Vieira MLC, DeCastro S, Maron MS, Kuvin JT, Patel AR, Pandian NG. What is the best approach for the assessment of left atrial size? Comparison of various unidimensional and two-dimensional parameters with three-dimensional echocardiographically determined left atrial volume. 2006;19:1026-32
Maden O (*see* Ozeke et al). 2006;19:1245-50
Maden O (*see* Tufekcioglu et al). 2006;19:1320-5
Madronero LF (*see* Roberson et al). 2006;19:1175-81
Mady C (*see* Salemi et al). 2006;19:354 (Case rep.)
Magalski A (*see* Main et al). 2006;19:64-70
Mahapatra S, Nishimura RA, Oh JK, McGoon MD. The prognostic value of pulmonary vascular capacitance determined by Doppler echocardiography in patients with pulmonary arterial hypertension. 2006;19:1045-50
Mahfouz Badran H (*see* Fahmy Elnoamany et al). 2006;19:1449-57
Mahfouz Badran H, Elnoamany M. Impact of type 2 diabetes mellitus on aortic elastic properties in normotensive diabetes: Doppler tissue imaging study. 2006;19:1471-81
Main ML (*see* Khumri et al). 2006;19:1072.e1-e3 (Case rep.)
Main ML, Ehlgen A, Coggins TR, Morris BA, Lanza P, Tremblay TM, Schiller NB, Goldman JH. Pulmonary hemodynamic effects of dipyridamole infusion in patients with normal and elevated pulmonary artery systolic pressure receiving PB127. 2006;19:1038-44
Main ML, Hannen MN, Kusnetzky LL, Martin JL, Coggins TR, Lanza P, Morris BA, Magalski A, Jones PG. Myocardial contrast echocardiographic estimates of infarct size predict likelihood of left ventricular remodeling after acute anterior wall myocardial infarction. 2006;19:64-70
Maione A (*see* Citro et al). 2006;19:1402.e5-e8 (Case rep.)
Makita Y (*see* Okamoto et al). 2006;19:211-4
Makoui M (*see* Bara et al). 2006;19:563-8
Maleki M (*see* Sadeghpour et al). 2006;19:1073.e1-e5 (Case rep.)
Malhotra D (*see* Vennalaganti et al). 2006;19:1286-92 (Case rep.)
Malm S, Frigstad S, Sagberg E, Steen PA, Skjarpe T. Real-time simultaneous triplane contrast echocardiography gives rapid, accurate, and reproducible assessment of left ventricular volumes and ejection fraction: A comparison with magnetic resonance imaging. 2006;19:1494-501
Malm S, Frigstad S, Stoylen A, Torp H, Sagberg E, Skjarpe T. Effects of ultrasound contrast during tissue velocity imaging on regional left ventricular velocity, strain, and strain rate measurements. 2006;19:40-7
Malm S, Frigstad S, Torp H, Wiseth R, Skjarpe T. Quantitative adenosine real-time myocardial contrast echocardiography for detection of angiographically significant coronary artery disease. 2006;19:365-72
Malouf PJ, Madani M, Gurudevan S, Waltman TJ, Raisinghani AB, DeMaria AN, Blanchard DG. Assessment of diastolic function with Doppler tissue imaging after cardiac surgery: Effect of the "postoperative septum" in on-pump and off-pump procedures. 2006;19:464-7
Manduca S (*see* Dialetto et al). 2006;19:836.e5-e8 (Case rep.)
Manning WJ (*see* Subramaniam et al). 2006;19:429-33
Mar de la Torre M (*see* Pérez-Sanz et al). 2006;19:1402.e5-e8 (Case rep.)
Marboe C (*see* Engel et al). 2006;19:932-7
Margonato A (*see* Agricola et al). 2006;19:457-63
Mariani M (*see* Di Bello et al). 2006;19:1063-71
Mariñas J (*see* Peteiro et al). 2006;19:1229-37
Mariñas J (*see* Peteiro et al). 2006;19:894-901
Marincheva G (*see* Khamaisi et al). 2006;19:835.e5-e8 (Case rep.)
Marino, on behalf of the Study Group of Echocardio PN, graphy of the Italian Society of Cardiology (*see* Mondillo et al). 2006;19:1085-91
Markham LW, Michelfelder EC, Border WL, Khoury PR, Spicer RL, Wong BL, Benson DW, Cripe LH. Abnormalities of diastolic function precede dilated cardiomyopathy associated with Duchenne muscular dystrophy. 2006;19:865-71
Maron MS (*see* Maddukuri et al). 2006;19:1026-32
Marshall JE (*see* Neilan et al). 2006;19:1280-5
Marsili A (*see* Di Bello et al). 2006;19:1063-71
Martin JL (*see* Main et al). 2006;19:64-70
Martin RP (*see* El-Chami et al). 2006;19:579.e5-e7 (Case rep.)
Martin RP (*see* Pohlel et al). 2006;19:229 (Case rep.)
Martin U (*see* Bara et al). 2006;19:563-8
Martinez MW, Singh CP, Mookadam F. Transcatheter closure of a postinfarction ventricular septal rupture. 2006;19:1401.e5-e7 (Case rep.)
Martino S (*see* Palmieri et al). 2006;19:491-8

Marwick TH (*see* Chan et al). 2006;19:192-201
Marwick TH (*see* Jenkins et al). 2006;19:1119-28
Masotti G (*see* Palmieri et al). 2006;19:483-90
Masuda K (*see* Okuda et al). 2006;19:781-7
Masuda K (*see* Otani et al). 2006;19:973-81
Mathew JP, Glas K, Troianos CA, Sears-Rogan P, Savage R, Shanewise J, Kisslo J, Aronson S, Shernan S. American society of echocardiography/society of cardiovascular anesthesiologists recommendations and guidelines for continuous quality improvement in perioperative echocardiography. 2006;19:1303-13 (Letter to the editor)
Mathew ST (*see* Mihalatos et al). 2006;19:76-82
Mauri F (*see* Moreo et al). 2006;19:938 (Case rep.)
Mazza do Nascimento M (*see* Yumi Hayashi et al). 2006;19:1092-101
Mazzaro E (*see* Tosoratti et al). 2006;19:1190.e5-e8 (Case rep.)
Mazzucchelli L (*see* Di Valentino et al). 2006;19:939 (Case rep.)
McCloskey VR (*see* Kean et al). 2006;19:202-5
McCulloch M, Dokainish H. Response to: "Color and spectral modes of tissue Doppler imaging have similar diagnostic utility but different numerical values". 2006;19:1411 (Letter)
McCulloch M, Zoghbi WA, Davis R, Thomas C, Dokainish H. Color tissue Doppler myocardial velocities consistently underestimate spectral tissue Doppler velocities: Impact on calculation peak transmitral pulsed Doppler velocity/early diastolic tissue Doppler velocity (E/Ea). 2006;19:744-8
McCully RB (*see* Yang et al). 2006;19:1136-43
McGhie JS (*see* Nemes et al). 2006;19:108 (Case rep.)
McGhie JS (*see* Nemes et al). 2006;19:355 (Case rep.)
McGhie JS (*see* van den Bosch et al). 2006;19:1-6
McGhie JS (*see* van den Bosch et al). 2006;19:7-13
McGhie JS (*see* van den Bosch et al). 2006;19:815-21
McGill JB (*see* Peterson et al). 2006;19:749-55
McGoon MD (*see* Mahapatra et al). 2006;19:1045-50
McIntosh M (*see* Horton et al). 2006;19:835.e1-e5 (Case rep.)
Mckenna WJ (*see* Sharma et al). 2006;19:354 (Case rep.)
McMahon CJ (*see* Havemann et al). 2006;19:355 (Case rep.)
McMahon EM (*see* Wang et al). 2006;19:1102-11
McNamara RL (*see* Mena et al). 2006;19:1295-300 (Review article)
McPherson DD (*see* Kean et al). 2006;19:202-5
Mehlman DJ (*see* Kean et al). 2006;19:202-5
Mehta RL (*see* Sharma et al). 2006;19:266-73
Meijboom FJ (*see* van den Bosch et al). 2006;19:1-6
Meijboom FJ (*see* van den Bosch et al). 2006;19:7-13
Meijboom FJ (*see* van den Bosch et al). 2006;19:815-21
Meijer A (*see* Jansen et al). 2006;19:185-91
Meimoun P, Sayah S, Tcheuffa JC, Benali T, Luycx-Bore A, Levy F, Tribouilloy C. Transthoracic coronary flow velocity reserve assessment: Comparison between adenosine and dobutamine. 2006;19:1220-8
Mele D, Soukhomovskaia O, Pacchioni E, Merli E, Avigni N, Federici L, Levine RA, Ferrari R. Improved detection of left ventricular thrombi and spontaneous echocontrast by tissue harmonic imaging in patients with myocardial infarction. 2006;19:1373-81
Melo Barbosa M (*see* Freire et al). 2006;19:1251-6
Memon N (*see* Rovner et al). 2006;19:872-9
Mena C, Wencker D, Krumholz HM, McNamara RL. Detection of heart transplant rejection in adults by echocardiographic diastolic indices: A systematic review of the literature. 2006;19:1295-300 (Review article)
Menafoglio A (*see* Di Valentino et al). 2006;19:939 (Case rep.)
Mercer-Rosa L, Seliem MA, Fedec A, Rome J, Rychik J, Gaynor JW. Illustration of the additional value of real-time 3-dimensional echocardiography to conventional transthoracic and transesophageal 2-dimensional echocardiography in imaging muscular ventricular septal defects: Does this have any impact on individual patient treatment?. 2006;19:1511-9
Meris A (*see* Agricola et al). 2006;19:457-63
Merli E (*see* Mele et al). 2006;19:1373-81
Mertens L (*see* Eyskens et al). 2006;19:994-1000
Meune C, Bertherat J, Dousset B, Jude N, Bertagna X, Duboc D, Weber S. Reduced myocardial contractility assessed by tissue Doppler echocardiography is associated with increased risk during adrenal surgery of patients with pheochromocytoma: Report of a preliminary study. 2006;19:1466-70
Micari A, Sklenar J, Belcik TA, Kaul S, Lindner JR. Automated quantification of the spatial extent of perfusion defects and viability on myocardial contrast echocardiography. 2006;19:379-85
Michel MC (*see* Hardziyenka et al). 2006;19:1272-9
Michelfelder EC (*see* Markham et al). 2006;19:865-71
Migliaresi P (*see* Palmieri et al). 2006;19:491-8
Mihalatos DG, Gopal AS, Kates R, Toole RS, Bercow NR, Lamendola C, Berkay SH, Damus P, Robinson N, Grimson R, Shen K, Reichek N. Intraoperative assessment of mitral regurgitation: Role of phenylephrine challenge. 2006;19:1158-64
Mihalatos DG, Mathew ST, Gopal AS, Joseph S, Grimson R, Reichek N. Relationship of mitral annular remodeling to severity of chronic mitral regurgitation. 2006;19:76-82
Mikami T (*see* Goto et al). 2006;19:857-64
Miller FA (*see* Bos et al). 2006;19:1033-7
Minichiello S (*see* Palmieri et al). 2006;19:491-8
Mirza A (*see* Anavekar et al). 2006;19:28-33
Mitchell SA, Crone RA. Takotsubo cardiomyopathy: A case report. 2006;19:1190.e5-e8 (Case rep.)
Miyamoto T (*see* Kobal et al). 2006;19:1012-6
Miyazaki C (*see* Takeuchi et al). 2006;19:1056-62
Mizoguchi T (*see* Tanaka et al). 2006;19:756-62
Moczulski Z (*see* Janus et al). 2006;19:1191.e5-e7 (Case rep.)
Mohler E (*see* Gerhard-Herman et al). 2006;19:955-72 (American Society of Echocardiography Report)
Mohler E (*see* Roman et al). 2006;19:943-54 (American Society of Echocardiography Report)
Møller JE (*see* Bergeron et al). 2006;19:772-6
Molloy MD (*see* Naqvi et al). 2006;19:147-53
Monda V (*see* Pisacane et al). 2006;19:107 (Case rep.)
Mondillo S, Galderisi M, Ballo P, Marino, on behalf of the Study Group of Echocardio PN, graphy of the Italian Society of Cardiology. Left ventricular systolic longitudinal function: Comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals. 2006;19:1085-91
Monserrat L (*see* Peteiro et al). 2006;19:894-901
Monserrrat L (*see* Peteiro et al). 2006;19:1229-37
Mookadam F (*see* Martinez et al). 2006;19:1401.e5-e7 (Case rep.)
Moon J (*see* Lim et al). 2006;19:280-4
Moore PA, Segar DS. Twenty-five-year-old man with a giant left ventricular mass. 2006;19:1286-92 (Case rep.)
Mor-Avi V (*see* Corsi et al). 2006;19:886-93
Mor-Avi V, Jacobs LD, Weiss RJ, Sugeng L, Weinert L, Bouchard T, Spencer KT, Lang RM. Color encoding of endocardial motion improves the interpretation of contrast-enhanced echocardiographic stress tests by less-experienced readers. 2006;19:48-54
Morelli C (*see* Caputo et al). 2006;19:836.e1-e4 (Case rep.)
Moreno CA (*see* Lester et al). 2006;19:919-23
Moreo A, De Chiara B, Bruschi G, Tarelli G, Parolini M, Mauri F. An unusual cause of pulmonary edema: Acute rupture of noncoronary sinus of valsalva aneurysm into the left atrium. 2006;19:938 (Case rep.)
Morhy SS (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Morioka S (*see* Katayama et al). 2006;19:335-40
Mörner S (*see* Lindqvist et al). 2006;19:1345-9
Morris BA (*see* Main et al). 2006;19:1038-44
Morris BA (*see* Main et al). 2006;19:64-70
Motomura N (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Mottola G (*see* Picano et al). 2006;19:356-63 (Review article)
Movahed M-R, Ahmadi-Kashani M, Kasravi B, Saito Y. Increased prevalence of mitral stenosis in women. 2006;19:911-3
Muangmingsuk S (*see* Roberson et al). 2006;19:1175-81
Mulvagh SL (*see* Hillis et al). 2006;19:450-6

Mumoli N, Cei M, Pasquinelli P, Carnesecchi C, Verzuri MS. Floating heart metastasis. 2006;19:1286-92 (Case rep.)
Muñóz-Castellanos L, Espinola-Zavaleta N, Kuri-Nivón M, Ruíz JF, Keirns C. Atrial septal defect: Anatomoechocardiographic correlation. 2006;19:1182-9
Muratore R (*see* Engel et al). 2006;19:932-7
Murphy DJ (*see* Leeper et al). 2006;19:1191.e5-e7 (Case rep.)

N

Naccarato GAF (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Nagai R (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Nair P, Siu SC, Sloggett CE, Biclar L, Sidhu RS, Yu EHC. The assessment of technical and interpretative proficiency in echocardiography. 2006;19:924-31
Naka Y (*see* LaRocca et al). 2006;19:464-7 (Case rep.)
Nakai H (*see* Takeuchi et al). 2006;19:1077-84
Nakai H, Takeuchi M, Nishikage T, Kokumai M, Otani S, Lang RM. Effect of aging on twist-displacement loop by 2-dimensional speckle tracking imaging. 2006;19:880-5
Nakamura H (*see* Hirano et al). 2006;19:536-9
Nakano M (*see* Hirano et al). 2006;19:536-9
Nakano T (*see* Yamanaka-Funabiki et al). 2006;19:1332-7
Nakatani S (*see* Iwahashi et al). 2006;19:1238-44
Nam G-B (*see* Kang et al). 2006;19:178-84
Nanda N (*see* Domanski and Nanda). 2006;19:941 (Editorial)
Naqvi T (*see* Gerhard-Herman et al). 2006;19:955-72 (American Society of Echocardiography Report)
Naqvi TZ (*see* Kobal et al). 2006;19:1012-6
Naqvi TZ (*see* Roman et al). 2006;19:943-54 (American Society of Echocardiography Report)
Naqvi TZ, Huynh HK. A new window of opportunity in echocardiography. 2006;19:569-77
Naqvi TZ, Zarbatany D, Molloy MD, Logan J, Buchbinder M. Intracardiac echocardiography for percutaneous mitral valve repair in a swine model. 2006;19:147-53
Nay S (*see* Horton et al). 2006;19:835.e1-e5 (Case rep.)
Nayyar S (*see* Khumri et al). 2006;19:1072.e1-e3 (Case rep.)
Neilan TG (*see* Jassal et al). 2006;19:229 (Case rep.)
Neilan TG (*see* Pérez-Sanz et al). 2006;19:1402.e5-e8 (Case rep.)
Neilan TG, Ton-Nu T-T, Jassal DS, Popovic ZB, Douglas PS, Halpern EF, Marshall JE, Thomas JD, Picard MH, Yoerger DM, Wood MJ. Myocardial adaptation to short-term high-intensity exercise in highly trained athletes. 2006;19:1280-5
Neishi Y (*see* Akiyama et al). 2006;19:55-63
Neishi Y (*see* Kume et al). 2006;19:914-8
Nemes A (*see* Geleijnse et al). 2006;19:1076 (Letter)
Nemes A, Lagrand WK, McGhie JS, ten Cate FJ. Three-dimensional transesophageal echocardiography in the evaluation of aortic valve destruction by endocarditis. 2006;19:355 (Case rep.)
Nemes A, McGhie JS, ten Cate FJ. Real-time 3-dimensional echocardiographic evaluation of aortic dissection. 2006;19:108 (Case rep.)
Nesser HJ, Davogg K, Winter S, Ammer M, Khandheria BK. Right ventricular metastasis of uterine squamous cell carcinoma. 2006;19:1286-92 (Case rep.)
Neuman Y (*see* Kobal et al). 2006;19:1012-6
Nguyen K (*see* Katz et al). 2006;19:1446-8
Niehaus M (*see* Bara et al). 2006;19:563-8
Nielsen JC (*see* Katz et al). 2006;19:1446-8
Nii M (*see* Roman et al). 2006;19:1382-92
Nii M (*see* Takahashi et al). 2006;19:1502-10
Nii M, Guerra V, Roman KS, Macgowan CK, Smallhorn JF. Three-dimensional tricuspid annular function provides insight into the mechanisms of tricuspid valve regurgitation in classic hypoplastic left heart syndrome. 2006;19:391-402
Nii M, Roman KS, Kingdom J, Redington AN, Jaeggi ET. Assessment of the evolution of normal fetal diastolic function during mid and late gestation by spectral Doppler tissue echocardiography. 2006;19:1431-7
Nii M, Shimizu M, Roman KS, Konstantinov I, Li J, Redington AN, Jaeggi ET. Doppler tissue imaging in the assessment of atrioventricular conduction time: Validation of a novel technique and comparison with electrophysiologic and pulsed wave Doppler-derived equivalents in an animal model. 2006;19:314-21
Nilsson B, Henningsson R, Brodin L-Å, Wandt B. Transesophageal echocardiographic acquisition of mitral annulus motion during monitoring of left ventricular function in anesthetized patients. 2006;19:499-506
Nilsson G (*see* Hedberg et al). 2006;19:88-94
Nishikage T (*see* Nakai et al). 2006;19:880-5
Nishikage T (*see* Takeuchi et al). 2006;19:1077-84
Nishimura RA (*see* Mahapatra et al). 2006;19:1045-50
Nishimura RA (*see* Sorajja et al). 2006;19:788-95
Nobori T (*see* Yamanaka-Funabiki et al). 2006;19:1332-7
Nomura C (*see* Jassal et al). 2006;19:229 (Case rep.)
Noohi F (*see* Sadeghpour et al). 2006;19:1073.e1-e5 (Case rep.)
Nose T (*see* Tanaka et al). 2006;19:756-62
Nowak J (*see* Lind et al). 2006;19:255-65
Nurkalem Z (*see* Uslu et al). 2006;19:341-6

O

Ogasawara Y (*see* Watanabe et al). 2006;19:71-5
Ogunyankin KO. Color and spectral modes of tissue Doppler imaging have similar diagnostic utility but different numerical values. 2006;19:1411 (Letter)
Oh B-H (*see* Kim et al). 2006;19:386-90
Oh B-H (*see* Sohn et al). 2006;19:777-80
Oh JK (*see* Ha et al). 2006;19:1314-9
Oh JK (*see* Hillis et al). 2006;19:450-6
Oh JK (*see* Kjaergaard et al). 2006;19:322-8
Oh JK (*see* Kjaergaard et al). 2006;19:515-21
Oh JK (*see* Mahapatra et al). 2006;19:1045-50
Oh JK (*see* Yang et al). 2006;19:1136-43
Oh Y-S (*see* Youn et al). 2006;19:373-8
Ohte N (*see* Yamanaka-Funabiki et al). 2006;19:1332-7
Ojaghi M (*see* Sadeghpour et al). 2006;19:1073.e1-e5 (Case rep.)
Okahashi N (*see* Kume et al). 2006;19:914-8
Okajima K (*see* Pulerwitz et al). 2006;19:540-5
Okamoto M, Fujii Y, Makita Y, Kajihara K, Yamasaki S, Iwamoto A, Hashimoto M, Sueda T, Yoshida N. Left atrial appendage function in patients with systemic embolism in spite of in sinus rhythm. 2006;19:211-4
Okuda K, Asanuma T, Hirano T, Masuda K, Otani K, Ishikura F, Beppu S. Impact of the coronary flow reduction at rest on myocardial perfusion and functional indices derived from myocardial contrast and strain echocardiography. 2006;19:781-7
Okuda N (*see* Ito et al). 2006;19:987-93
Okura H, Fuyuki H, Kubo T, Iwata K, Taguchi H, Toda I, Yoshikawa J. Noninvasive diagnosis of ischemic and nonischemic cardiomyopathy using coronary flow velocity measurements of the left anterior descending coronary artery by transthoracic Doppler echocardiography. 2006;19:552-8
O'Leary EL (*see* Elhendy et al). 2006;19:1482-7
O'Leary P (*see* Bos et al). 2006;19:1033-7
Olson JJ, Costa SP, Young CE, Palac RT. Early mitral filling/diastolic mitral annular velocity ratio is not a reliable predictor of left ventricular filling pressure in the setting of severe mitral regurgitation. 2006;19:83-7
Ommen SR (*see* Sorajja et al). 2006;19:788-95
Onishi K (*see* Yamanaka-Funabiki et al). 2006;19:1332-7
Onishi T (*see* Tanaka et al). 2006;19:756-62
Ono M (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Onozuka H (*see* Goto et al). 2006;19:857-64
Onuk T (*see* Dogan et al). 2006;19:243-8
Oppizzi M (*see* Agricola et al). 2006;19:457-63
Ortiz Oficialdegui P (*see* Perez de Isla et al). 2006;19:1338-44
Oryszak S (*see* Shin et al). 2006;19:796-801
Ostfeld RJ (*see* Ahmed et al). 2006;19:938 (Case rep.)
Ostfeld RJ (*see* Vennalaganti et al). 2006;19:1286-92 (Case rep.)
Otani K (*see* Okuda et al). 2006;19:781-7

Otani K, Masuda K, Asanuma T, Ishikura F, Beppu S. Corrected quantification method to determine myocardial blood flow using real-time myocardial contrast echocardiography. 2006; 19:973-81
Otani S (*see* Nakai et al). 2006;19:880-5
Otani S (*see* Takeuchi et al). 2006;19:1077-84
Otani S (*see* Takeuchi et al). 2006;19:294-9
Otsuka R (*see* Engel et al). 2006;19:932-7
Otsuka R (*see* Hirata et al). 2006;19:165-71
Otsuka R (*see* Pulerwitz et al). 2006;19:540-5
Otto CM (*see* Jung et al). 2006;19:802-10
Ozdemir K (*see* Can et al). 2006;19:1074.e1-e2 (Case rep.)
Ozeke O (*see* Tufekcioglu et al). 2006;19:1320-5
Ozeke O, Elhassan HHM, Aras D, Tufekcioglu O, Maden O, Selcuk MT, Topaloglu S, Deveci B, Yildiz A, Ulupinar H, Sahin O, Ergun K, Celenk MK, Golbasi Z. Hemodynamic change in transmitral gradient during Valsalva maneuver and its relation with left atrial function, functional capacity, and pulmonary artery pressure in patients with mitral stenosis in sinus rhythm. 2006;19: 1245-50
Ozer O (*see* Uslu et al). 2006;19:341-6
Ozturk S (*see* Cay et al). 2006;19:1286-92 (Case rep.)

P

Pac A (*see* Cay et al). 2006;19:1286-92 (Case rep.)
Pacchioni E (*see* Mele et al). 2006;19:1373-81
Padmanabhan S (*see* Sachdev et al). 2006;19:34-9
Pac WE (*see* Yang et al). 2006;19:1051-5
Pai RG (*see* Varadarajan et al). 2006;19:1458-61
Paimany B (*see* Premkumar et al). 2006;19:107 (Case rep.)
Palac RT (*see* Olson et al). 2006;19:83-7
Palagi C (*see* Di Bello et al). 2006;19:1063-71
Palmieri EA (*see* Palmieri et al). 2006;19:491-8
Palmieri V, Capaldo B, Russo C, Iaccarino M, Di Minno G, Riccardi G, Celentano A. Left ventricular chamber and myocardial systolic function reserve in patients with type 1 diabetes mellitus: Insight from traditional and Doppler tissue imaging echocardiography. 2006;19:848-56
Palmieri V, Innocenti F, Agresti C, Caldi F, Masotti G, Pini R. Traditional and color M-Mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: Relations to contractility reserve. 2006;19:483-90
Palmieri V, Russo C, Palmieri EA, Arezzi E, Pezzullo S, Minichiello S, Martino S, Migliaresi P, Celentano A. Isolated left ventricular diastolic dysfunction: Implications for exercise left ventricular performance in patients without congestive heart failure. 2006; 19:491-8
Palta M (*see* Gepner et al). 2006;19:223-8
Pandian NG (*see* Maddukuri et al). 2006;19:1026-32
Panzica PJ (*see* Subramaniam et al). 2006;19:429-33
Papadopoulos CE (*see* Karvounis et al). 2006;19:1150-7
Paraskevaidis IA, Tsiapras D, Karavolias G, Adamopoulos S, Dodouras T, Cokkinos P, Th Kremastinos D. The effect of carvedilol therapy on myocardial functional reserve in patients with advanced heart failure caused by nonischemic dilated cardiomyopathy. 2006;19:529-35
Paraskevaidis IA, Tsiapras D, Kyrzopoulos S, Cokkinos P, Iliodromitis EK, Parissis J, Kremastinos DT. The role of left ventricular long-axis contraction in patients with asymptomatic aortic regurgitation. 2006;19:249-54
Paraskevaidis S (*see* Karvounis et al). 2006;19:1150-7
Parharidis GE (*see* Karvounis et al). 2006;19:1150-7
Parharidis GE (*see* Zaglavara et al). 2006;19:300-6
Parissis J (*see* Paraskevaidis et al). 2006;19:249-54
Park C-S (*see* Youn et al). 2006;19:373-8
Park S (*see* Choi et al). 2006;19:1211-9
Park Y-B (*see* Kim et al). 2006;19:386-90
Park Y-B (*see* Sohn et al). 2006;19:777-80
Parness IA (*see* Katz et al). 2006;19:1446-8
Parolini M (*see* Moreo et al). 2006;19:938 (Case rep.)
Pasquinelli P (*see* Mumoli et al). 2006;19:1286-92 (Case rep.)
Patel A (*see* Woods and Patel). 2006;19:215-22
Patel AR (*see* Maddukuri et al). 2006;19:1026-32
Pauliks LB (*see* Dyer et al). 2006;19:21-7
Pedone C, Biagini E, Galema TW, Vletter WB, ten Cate FJ. Myocardial perfusion after percutaneous transluminal septal myocardial ablation as assessed by myocardial contrast echocardiography in patients with hypertrophic obstructive cardiomyopathy. 2006;19:982-6
Peels KH (*see* Jansen et al). 2006;19:185-91
Pehrsson K (*see* Hedberg et al). 2006;19:88-94
Peighambari M (*see* Sadeghpour et al). 2006;19:1073.e1-e5 (Case rep.)
Pellerin D (*see* Sharma et al). 2006;19:266-73
Pellerin D (*see* Sharma et al). 2006;19:354 (Case rep.)
Pellerin M (*see* Gisbert et al). 2006;19:140-6
Pellikka PA (*see* Bergeron et al). 2006;19:772-6
Pellikka PA (*see* Yang et al). 2006;19:1136-43
Pemberton J, Ge S, Thiele K, Jerosch-Herold M, Sahn DJ. Real-time three-dimensional color Doppler echocardiography overcomes the inaccuracies of spectral Doppler for stroke volume calculation. 2006;19:1403-10 (Review article)
Pennell DJ (*see* Lim et al). 2006;19:280-4
Perez de Isla L, Ortiz Oficialdegui P, Florit J, Angel Garcia-Fernandez M, Sanchez V, Zamorano J. Usefulness of clinical, electrocardiographic, and echocardiographic parameters to detect cardiac asynchrony in patients with left ventricular dysfunction secondary to ischemic or nonischemic heart disease. 2006;19:1338-44
Pérez-Sanz TM, Fulquet E, Neilan TG, Rollán MJ, Mar de la Torre M, Bratos JL. A case report of a round cystic tumor in the left ventricular outflow tract. 2006;19:1402.e5-e8 (Case rep.)
Perrino AC Jr., (*see* Popescu and Perrino Jr). 2006;19:1074.e5-e6 (Case rep.)
Perry R, De Pasquale CG, Chew DP, Aylward PE, Joseph MX. QRS duration alone misses cardiac dyssynchrony in a substantial proportion of patients with chronic heart failure. 2006;19: 1257-63
Peteiro J, Monserrrat L, Bouzas A, Piñon P, Mariñas J, Piñeiro M, Castro-Beiras A. Prognostic value of mitral regurgitation assessment during exercise echocardiography in patients with known or suspected coronary artery disease. 2006;19:1229-37
Peteiro JC, Monserrat L, Bouzas A, Piñon P, Mariñas J, Bouzas B, Castro-Beiras A. Risk stratification by treadmill exercise echocardiography. 2006;19:894-901
Peters GL, Davies RA, Veinot JP, Burwash IG. Cardiac actinomycosis: An unusual cause of an intracardiac mass. 2006;19: 1530.e7-e11 (Case rep.)
Peters PJ, Reinhardt S. The echocardiographic evaluation of intracardiac masses: A review. 2006;19:230-40 (Review article)
Peterson LR, Waggoner AD, Fuentes L, Schechtman KB, McGill JB, Gropler RJ, Dávila-Román VG. Alterations in left ventricular structure and function in type-1 diabetics: A focus on left atrial contribution to function. 2006;19:749-55
Peterson RE, Levi DS, Williams RJ, Lai WW, Sklansky MS, Drant S. Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum. 2006;19:1393-400
Petre L (*see* Bangalore et al). 2006;19:438-40
Pezzullo S (*see* Palmieri et al). 2006;19:491-8
Pfeffer MA (*see* Anavekar et al). 2006;19:28-33
Pfeffer MA (*see* Thune et al). 2006;19:1462-5
Picano E (*see* Agricola et al). 2006;19:457-63
Picano E, Frassi F, Agricola E, Gligorova S, Gargani L, Mottola G. Ultrasound lung comets: A clinically useful sign of extravascular lung water. 2006;19:356-63 (Review article)
Picard MH (*see* Neilan et al). 2006;19:1280-5
Piechaczek C (*see* Bara et al). 2006;19:563-8
Pignatelli RH (*see* Havemann et al). 2006;19:355 (Case rep.)
Pignatelli RH (*see* Lai et al).
Pigula FA (*see* Wald et al). 2006;19:939.e4-36 (Case rep.)
Pillay TM (*see* Zaglavara et al). 2006;19:300-6

Pinchera A (*see* Di Bello et al). 2006;19:1063-71
Piñeiro M (*see* Peteiro et al). 2006;19:1229-37
Ping C (*see* Lim et al). 2006;19:280-4
Pini R (*see* Palmieri et al). 2006;19:483-90
Piñon P (*see* Peteiro et al). 2006;19:1229-37
Piñon P (*see* Peteiro et al). 2006;19:894-901
Pinsky MR (*see* Dohi et al). 2006;19:475-82
Pisacane C, Comenale Pinto S, De Gregorio P, Pisacane F, Monda V, Caso P, Calabrò R. "Steal" collaterals: An echocardiographic diagnostic marker for anomalous origin of the left main coronary artery from the pulmonary artery in the adult. 2006;19:107 (Case rep.)
Pisacane F (*see* Pisacane et al). 2006;19:107 (Case rep.)
Pisani M (*see* Agricola et al). 2006;19:457-63
Pita CG (*see* Salemi et al). 2006;19:354 (Case rep.)
Pizzarello R (*see* LaRocca et al). 2006;19:464-7 (Case rep.)
Plante E, Lachance D, Roussel Élise, Drolet M-C, Arsenault M, Couet J. Impact of anesthesia on echocardiographic evaluation of systolic and diastolic function in rats. 2006;19:1520-5
Plappert T (*see* Anavekar et al). 2006;19:28-33
Playford D (*see* Edibam et al). 2006;19:938.e1-e4 (Case rep.)
Pohlel K, Lerakis S, Arita T, Rab T, Martin RP. Intracoronary stent visualized on transesophageal echocardiogram in a case of coronary dissection complicated by aortic dissection. 2006;19:229 (Case rep.)
Pollick C (*see* Kobal et al). 2006;19:1012-6
Polonsky TS (*see* Corsi et al). 2006;19:886-93
Pommerantzeff PM (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Popescu WM, Perrino AC Jr.,. Critical cardiac decompensation during laparoscopic surgery. 2006;19:1074.e5-e6 (Case rep.)
Popovic ZB (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Popovic ZB (*see* Neilan et al). 2006;19:1280-5
Porter TR (*see* Elhendy et al). 2006;19:1482-7
Premkumar V, Paimany B, Gopal AS. Primary large B-cell cardiac lymphoma. 2006;19:107 (Case rep.)
Price J (*see* Havemann et al). 2006;19:355 (Case rep.)
Pritchard C (*see* Belham et al). 2006;19:206-10
Proietti P (*see* De Luca et al). 2006;19:172-7
Provenza G (*see* Citro et al). 2006;19:1402.e5-e8 (Case rep.)
Pu M (*see* Yang et al). 2006;19:1051-5
Pucci A (*see* Di Bello et al). 2006;19:1063-71
Pulerwitz T, Hirata K, Abe Y, Otsuka R, Herz S, Okajima K, Jin Z, Di Tullio MR, Homma S. Feasibility of using a real-time 3-dimensional technique for contrast dobutamine stress echocardiography. 2006;19:540-5

Q

Qin JX (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Qureshi A (*see* Banchs et al). 2006;19:1401.e1-e3 (Case rep.)
Qureshi A (*see* Jassal et al). 2006;19:229 (Case rep.)

R

Rab T (*see* Pohlel et al). 2006;19:229 (Case rep.)
Raffoul H (*see* Cohen et al). 2006;19:580 (Letter)
Raisinghani AB (*see* Malouf et al). 2006;19:464-7
Ramasubbu K, Coselli J, Zoghbi WA. Unusual complication of aortic root reconstruction with sparing of the aortic valve: Left ventricular outflow tract to right atrial fistula. 2006;19:464-7 (Case rep.)
Rao YN (*see* Sachdev et al). 2006;19:34-9
Rauch A (*see* Weidemann et al). 2006;19:839-47
Recupero A (*see* de Gregorio et al). 2006;19:1190.e1-e4 (Case rep.)
Redington A (*see* Takahashi et al). 2006;19:1502-10
Redington AN (*see* Nii et al). 2006;19:1431-7
Redington AN (*see* Nii et al). 2006;19:314-21
Reho I (*see* Enseleit et al). 2006;19:1190.e5-e8 (Case rep.)
Reichek N (*see* Mihalatos et al). 2006;19:1158-64
Reichek N (*see* Mihalatos et al). 2006;19:76-82
Reinhardt S (*see* Peters and Reinhardt). 2006;19:230-40 (Review article)
Reisner SA (*see* Dabbah et al). 2006;19:733-43
Ressner M, Brodin L-A, Jansson T, Hoff L, Ask P, Janerot-Sjoberg B. Effects of ultrasound contrast agents on Doppler tissue velocity estimation. 2006;19:154-64
Rianne de Bruin-Bon HACM (*see* Hardziyenka et al). 2006;19:1272-9
Ribeiro de Oliveira A Jr., (*see* Freire et al). 2006;19:1251-6
Ribeiro de Oliveira Longo J (*see* Freire et al). 2006;19:1251-6
Riccardi G (*see* Palmieri et al). 2006;19:848-56
Richter H (*see* Cardello et al). 2006;19:1074.e3-e4 (Case rep.)
Rigolin VH (*see* Kean et al). 2006;19:202-5
Riley MF (*see* Subramaniam et al). 2006;19:429-33
Rim S-J (*see* Choi et al). 2006;19:1211-9
Rim S-J (*see* Ha et al). 2006;19:1314-9
Ringqvist I (*see* Hedberg et al). 2006;19:88-94
Rinkevich D (*see* Kalvaitis et al). 2006;19:1488-93
Robbers-Visser D (*see* van den Bosch et al). 2006;19:1-6
Roberson DA (*see* Cui and Roberson). 2006;19:1438-45
Roberson DA, Javois AJ, Cui W, Madronero LF, Cuneo BF, Muangmingsuk S. Double atrial septum with persistent interatrial space: Echocardiographic features of a rare atrial septal malformation. 2006;19:1175-81
Robinson N (*see* Mihalatos et al). 2006;19:1158-64
Rochitte CE (*see* Salemi et al). 2006;19:354 (Case rep.)
Rodrigues ACT (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Rodriguez CJ (*see* LaRocca et al). 2006;19:464-7 (Case rep.)
Rogowski J (*see* Fijalkowski et al). 2006;19:1359-64
Rollán MJ (*see* Pérez-Sanz et al). 2006;19:1402.e5-e8 (Case rep.)
Roman KS (*see* Nii et al). 2006;19:1431-7
Roman KS (*see* Nii et al). 2006;19:314-21
Roman KS (*see* Nii et al). 2006;19:391-402
Roman KS (*see* Takahashi et al). 2006;19:1502-10
Roman KS, Nii M, Macgowan CK, Barrea C, Coles J, Smallhorn JF. The impact of patch augmentation on left atrioventricular valve dynamics in patients with atrioventricular septal defects: Early and midterm follow-up. 2006;19:1382-92
Roman M (*see* Gerhard-Herman et al). 2006;19:955-72 (American Society of Echocardiography Report)
Roman MJ, Naqvi TZ, Gardin JM, Gerhard-Herman M, Jaff M, Mohler E. Clinical application of noninvasive vascular ultrasound in cardiovascular risk stratification: A report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology. 2006;19:943-54 (American Society of Echocardiography Report)
Rome J (*see* Mercer-Rosa et al). 2006;19:1511-9
Roos-Hesselink JW (*see* van den Bosch et al). 2006;19:1-6
Roos-Hesselink JW (*see* van den Bosch et al). 2006;19:7-13
Roos-Hesselink JW (*see* van den Bosch et al). 2006;19:815-21
Rosenthal DN (*see* Friedberg et al). 2006;19:559-62
Rothenburger M (*see* Bruch et al). 2006;19:522-8
Roumina S (*see* Lind et al). 2006;19:255-65
Roussel Élise (*see* Plante et al). 2006;19:1520-5
Rovner A (*see* Waggoner et al). 2006;19:307-13
Rovner A, de las Fuentes L, Waggoner AD, Memon N, Chohan R, Dávila-Román VG. Characterization of left ventricular diastolic function in hypertension by use of Doppler tissue imaging and color M-mode techniques. 2006;19:872-9
Roy TNS (*see* Sankar et al). 2006;19:241 (Letter)
Ruhparwar A (*see* Bara et al). 2006;19:563-8
Russo C (*see* Palmieri et al). 2006;19:491-8
Russo C (*see* Palmieri et al). 2006;19:848-56
Russo MG (*see* Caputo et al). 2006;19:836.e1-e4 (Case rep.)
Rutkowski B (*see* Fijalkowski et al). 2006;19:1359-64
Ruíz JF (*see* Muñóz-Castellanos et al). 2006;19:1182-9
Rychik J (*see* Lai et al).
Rychik J (*see* Mercer-Rosa et al). 2006;19:1511-9
Rychik J (*see* Seliem et al). 2006;19:121-9
Rynkiewicz A (*see* Fijalkowski et al). 2006;19:1359-64

S

Sachdev V, Aletras AH, Padmanabhan S, Sidenko S, Rao YN, Brenneman CL, Shizukuda Y, Lie GR, Vincent PS, Waclawiw MA, Arai AE. Myocardial strain decreases with increasing transmurality of infarction: A Doppler echocardiographic and magnetic resonance correlation study. 2006;19:34-9

Sadeghpour A, Peighambari M, Dalirrooyfard M, Esmailzadeh M, Maleki M, Noohi F, Ojaghi M, Samiei N. Aorta-to-left ventricle tunnel associated with noncompaction left ventricle. 2006;19:1073.e1-e5 (Case rep.)

Safadi T (*see* Shiran et al). 2006;19:578.e1-e4 (Case rep.)

Sagberg E (*see* Malm et al). 2006;19:1494-501

Sagberg E (*see* Malm et al). 2006;19:40-7

Saglam M (*see* Karakaya et al). 2006;19:434-7

Sahin O (*see* Ozeke et al). 2006;19:1245-50

Sahn DJ (*see* Pemberton et al). 2006;19:1403-10 (Review article)

Saito Y (*see* Movahed et al). 2006;19:911-3

Sajeev CG (*see* Sunil Roy et al). 2006;19:1402.e1-e4 (Case rep.)

Salemi VMC, Rochitte CE, Lemos P, Benvenuti LA, Pita CG, Mady C. Long-term survival of a patient with isolated noncompaction of the ventricular myocardium. 2006;19:354 (Case rep.)

Samiei N (*see* Sadeghpour et al). 2006;19:1073.e1-e5 (Case rep.)

Sanchez V (*see* Perez de Isla et al). 2006;19:1338-44

Sankar V, Roy TNS, Venugopal KN. Angle correction for proximal isovelocity surface area method: Is it spheric cap or lune?. 2006;19:241 (Letter)

Santini F (*see* Di Bello et al). 2006;19:1063-71

Santoro G (*see* Caputo et al). 2006;19:836.e1-e4 (Case rep.)

Santos Assreuy Diniz S (*see* Freire et al). 2006;19:1251-6

Sardella G (*see* De Luca et al). 2006;19:172-7

Sarubbi B (*see* Caputo et al). 2006;19:836.e1-e4 (Case rep.)

Sasaki K (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)

Savage R (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)

Sawhney N (*see* Waggoner et al). 2006;19:307-13

Sayah S (*see* Meimoun et al). 2006;19:1220-8

Schechtman KB (*see* Peterson et al). 2006;19:749-55

Scheld HH (*see* Bruch et al). 2006;19:522-8

Schiller NB (*see* Main et al). 2006;19:1038-44

Schneck SL (*see* Lester et al). 2006;19:919-23

Schneider C (*see* Bahlmann et al). 2006;19:578.e5-e8 (Case rep.)

Schuele S (*see* Ulrich et al). 2006;19:1165-9

Schultz AH (*see* Seliem et al). 2006;19:121-9

Scognamiglio G (*see* Dialetto et al). 2006;19:836.e5-e8 (Case rep.)

Sears-Rogan P (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)

Seeberger A (*see* Yumi Hayashi et al). 2006;19:1092-101

Segar DS (*see* Moore and Segar). 2006;19:1286-92 (Case rep.)

Seghatol FF (*see* Kean et al). 2006;19:202-5

Selcuk MT (*see* Ozeke et al). 2006;19:1245-50

Seliem MA (*see* Mercer-Rosa et al). 2006;19:1511-9

Seliem MA, Fedec A, Cohen MS, Ewing S, Farrell PE Jr., Rychik J, Schultz AH, Gaynor JW, Spray TL. Real-time 3-dimensional echocardiographic imaging of congenital heart disease using matrix-array technology: Freehand real-time scanning adds instant morphologic details not well delineated by conventional 2-dimensional imaging. 2006;19:121-9

Senior R (*see* Dwivedi et al). 2006;19:546-51

Senior R (*see* Lim et al). 2006;19:280-4

Seo H-S (*see* Choi et al). 2006;19:1211-9

Seow S-C, Koh L-P, Yeo T-C. Hemodynamic significance of mitral stenosis: Use of a simple, novel index by 2-dimensional echocardiography. 2006;19:102-6

Severyn D (*see* Dohi et al). 2006;19:475-82

Shaffer EM (*see* Dyer et al). 2006;19:21-7

Shah AS, Kukar A, Chaudhry FA, Sherrid MV. Unusual anomalous single papillary muscle causing symptomatic mid-left ventricular cavity obstruction: Octopus papillary muscle. 2006;19:939 (Case rep.)

Shandas R (*see* Dyer et al). 2006;19:21-7

Shandas R (*see* Dyer et al). 2006;19:403-12

Shanewise J (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)

Sharma R, Pellerin D, Gaze DC, Mehta RL, Gregson H, Streather CP, Collinson PO, Brecker SJD. Mitral peak Doppler e-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease. 2006;19:266-73

Sharma R, Smith J, Elliott PM, Mckenna WJ, Pellerin D. Left ventricular outflow tract obstruction caused by accessory mitral valve tissue. 2006;19:354 (Case rep.)

Sharma S (*see* Varadarajan et al). 2006;19:1458-61

Shawky Abdelaziz E (*see* Fahmy Elnoamany et al). 2006;19:1449-57

Shea S (*see* Toulany et al). 2006;19:14-20

Sheehan FH (*see* Jung et al). 2006;19:802-10

Shen K (*see* Mihalatos et al). 2006;19:1158-64

Shernan S (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)

Sherrid MV (*see* Shah et al). 2006;19:939 (Case rep.)

Sheu S-H (*see* Su et al). 2006;19:274-9

Shim W-H (*see* Ha et al). 2006;19:1314-9

Shimbo D (*see* LaRocca et al). 2006;19:464-7 (Case rep.)

Shimizu M (*see* Nii et al). 2006;19:314-21

Shin M-S, Fukuda S, Song J-M, Tran H, Oryszak S, Thomas JD, Shiota T. Relationship between left atrial and left ventricular function in hypertrophic cardiomyopathy: A real-time 3-dimensional echocardiographic study. 2006;19:796-801

Shiota T (*see* Asada-Kamiguchi et al). 2006;19:1112-8

Shiota T (*see* Shin et al). 2006;19:796-801

Shiota T (*see* Song et al). 2006;19:1129-35

Shirali GS (*see* Lai et al).

Shiran A, Zisman D, Karkabi B, Safadi T, Aravot D, Bitterman H, Lewis BS. Behçet's aortitis mimicking aortic valve endocarditis with subaortic complications. 2006;19:578.e1-e4 (Case rep.)

Shizukuda Y (*see* Sachdev et al). 2006;19:34-9

Sichrovsky T (*see* Bangalore et al). 2006;19:438-40

Siclari F (*see* Di Valentino et al). 2006;19:939 (Case rep.)

Siddiqui TS, Dawn B, Stoddard MF. Hypotension during dobutamine stress transesophageal echocardiography: Relationship with provoked left ventricular obstruction. 2006;19:1144-9

Sidenko S (*see* Sachdev et al). 2006;19:34-9

Sidhu RS (*see* Nair et al). 2006;19:924-31

Siegel RJ (*see* Kobal et al). 2006;19:1012-6

Sieswerda GT (*see* Klein et al). 2006;19:763-71

Sieswerda GTJ (*see* Dijkmans et al). 2006;19:285-93

Sila CA (*see* Ulrich et al). 2006;19:1165-9

Silbiger JJ (*see* Webber et al). 2006;19:1530.e1-e3 (Case rep.)

Silver M (*see* Lavi et al). 2006;19:1530.e5-e6 (Case rep.)

Silverman NH (*see* Friedberg and Silverman). 2006;19:1326-31

Silverstein JR, Tasset MR, Dowling RD, Alshaher MM. Traumatic intramyocardial left ventricular dissection: A case report. 2006;19:1529.e5-e8 (Case rep.)

Silvilairat S (*see* Bos et al). 2006;19:1033-7

Simoons ML (*see* van den Bosch et al). 2006;19:1-6

Simoons ML (*see* van den Bosch et al). 2006;19:7-13

Simoons ML (*see* van den Bosch et al). 2006;19:815-21

Simpson RA (*see* Lester et al). 2006;19:919-23

Sindermann J (*see* Bruch et al). 2006;19:522-8

Singh CP (*see* Martinez et al). 2006;19:1401.e5-e7 (Case rep.)

Siu SC (*see* Basic et al). 2006;19:1203-10

Siu SC (*see* Nair et al). 2006;19:924-31

Skali H (*see* Anavekar et al). 2006;19:28-33

Skali H (*see* Thune et al). 2006;19:1462-5

Skjarpe T (*see* Malm et al). 2006;19:1494-501

Skjarpe T (*see* Malm et al). 2006;19:365-72

Skjarpe T (*see* Malm et al). 2006;19:40-7

Sklansky MS (*see* Peterson et al). 2006;19:1393-400

Sklenar J (*see* Micari et al). 2006;19:379-85

Skyba DM (*see* Basic et al). 2006;19:1203-10

Sloggett C (*see* Basic et al). 2006;19:1203-10

Sloggett CE (*see* Nair et al). 2006;19:924-31

Smallhorn JF (*see* Nii et al). 2006;19:391-402
Smallhorn JF (*see* Roman et al). 2006;19:1382-92
Smallhorn JF (*see* Takahashi et al). 2006;19:1502-10
Smedira NG (*see* Song et al). 2006;19:1129-35
Smith J (*see* Sharma et al). 2006;19:354 (Case rep.)
Smulevitz B (*see* Vidovich and Smulevitz). 2005;19:1302 (Letter to the editor)
Snyder EM (*see* Kjaergaard et al). 2006;19:515-21
Sogaard P (*see* Kjaergaard et al). 2006;19:1301 (Letter)
Sogaard P (*see* Kjaergaard et al). 2006;19:1365-72
Sogaard P (*see* Kjaergaard et al). 2006;19:322-8
Sohn D-W (*see* Kim et al). 2006;19:386-90
Sohn D-W, Kim Y-J, Kim H-K, Kim K-Y, Koo B-K, Zo J-H, Kim H-S, Oh B-H, Park Y-B, Choi Y-S. Assessment of coronary vasodilatation in response to nitroglycerin with transthoracic Doppler echocardiography. 2006;19:777-80
Solomon SD (*see* Anavekar et al). 2006;19:28-33
Solomon SD (*see* Thune et al). 2006;19:1462-5
Song J-K (*see* Kang et al). 2006;19:178-84
Song J-M (*see* Kang et al). 2006;19:178-84
Song J-M (*see* Shin et al). 2006;19:796-801
Song J-M, Fukuda S, Lever HM, Daimon M, Agler DA, Smedira NG, Thomas JD, Shiota T. Asymmetry of systolic anterior motion of the mitral valve in patients with hypertrophic obstructive cardiomyopathy: A real-time three-dimensional echocardiographic study. 2006;19:1129-35
Sorajja P, Nishimura RA, Ommen SR, Ackerman MJ, Tajik AJ, Gersh BJ. Use of echocardiography in patients with hypertrophic cardiomyopathy: Clinical implications of massive hypertrophy. 2006;19:788-95
Soukhomovskaia O (*see* Mele et al). 2006;19:1373-81
Soulière V (*see* Gisbert et al). 2006;19:140-6
Spencer KT (*see* Mor-Avi et al). 2006;19:48-54
Spencer KT (*see* Takeuchi et al). 2006;19:294-9
Spevack DM (*see* Ahmed et al). 2006;19:938 (Case rep.)
Spevack DM (*see* Vennalaganti et al). 2006;19:1286-92 (Case rep.)
Spicer RL (*see* Markham et al). 2006;19:865-71
Spray TL (*see* Seliem et al). 2006;19:121-9
Springer RF (*see* Gilman et al). 2006;19:1526-8
Srivastava S (*see* Katz et al). 2006;19:1446-8
St John Sutton M (*see* Anavekar et al). 2006;19:28-33
Steen PA (*see* Malm et al). 2006;19:1494-501
Stein JH (*see* Gepner et al). 2006;19:1170-4
Stein JH (*see* Gepner et al). 2006;19:223-8
Stein JH (*see* Tzou et al). 2006;19:1286-92
Steinberg JS (*see* Bangalore et al). 2006;19:438-40
Stewart A (*see* LaRocca et al). 2006;19:464-7 (Case rep.)
Stoddard MF (*see* Banchs et al). 2006;19:1401.e1-e3 (Case rep.)
Stoddard MF (*see* Siddiqui et al). 2006;19:1144-9
Stöllberger C, Finsterer, *Wien, Austria J.* Hypertrabeculation is not distinct from noncompaction in neuromuscular and non-neuromuscular disorders. 2006;19:109-10 (Letter)
Stoylen A (*see* Malm et al). 2006;19:40-7
Streather CP (*see* Sharma et al). 2006;19:266-73
Strotmann JM (*see* Weidemann et al). 2006;19:839-47
Styliadis IH (*see* Karvounis et al). 2006;19:1150-7
Stypmann J (*see* Bruch et al). 2006;19:95-101
Su H-M, Lin T-H, Voon W-C, Lee K-T, Chu C-S, Yen H-W, Lai W-T, Sheu S-H. Single-beat differentiation among left ventricular filling patterns by pulsed wave Doppler echocardiography. 2006;19:274-9
Subramaniam B, Riley MF, Panzica PJ, Manning WJ. Transesophageal echocardiographic assessment of right atrial appendage anatomy and function: Comparison with the left atrial appendage and implications for local thrombus formation. 2006;19:429-33
Sueda T (*see* Okamoto et al). 2006;19:211-4
Sugeng L (*see* Corsi et al). 2006;19:886-93
Sugeng L (*see* Mor-Avi et al). 2006;19:48-54
Sugeng L, Coon P, Weinert L, Jolly N, Lammertin G, Bednarz JE, Thiele K, Lang RM. Use of real-time 3-dimensional transthoracic echocardiography in the evaluation of mitral valve disease. 2006;19:413-21
Sugioka K (*see* Engel et al). 2006;19:932-7
Sun B (*see* Yang et al). 2006;19:1051-5
Sunil Roy TN, Sajeev CG, Francis J, Krishnan MN, Venugopal K. Three major coronary artery-to-left ventricular fistula: An unusual cause of a diastolic murmur at the apex. 2006;19:1402.e1-e4 (Case rep.)
Sutherland GR (*see* Weidemann et al). 2006;19:839-47
Suwa M (*see* Ito et al). 2006;19:987-93

T

Taguchi H (*see* Okura et al). 2006;19:552-8
Tajik AJ (*see* Sorajja et al). 2006;19:788-95
Takagi A (*see* Takagi et al). 2006;19:1001-11
Takagi T (*see* Katayama et al). 2006;19:335-40
Takagi T, Takagi A, Hozumi T, Yoshikawa J. Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: Comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography. 2006;19:1001-11
Takahashi K, Guerra V, Roman KS, Nii M, Redington A, Smallhorn JF. Three-dimensional echocardiography improves the understanding of the mechanisms and site of left atrioventricular valve regurgitation in atrioventricular septal defect. 2006;19:1502-10
Takamoto S (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Takenaka K (*see* Uno et al). 2006;19:1401.e5-e7 (Case rep.)
Takeuchi M (*see* Nakai et al). 2006;19:880-5
Takeuchi M, Nakai H, Kokumai M, Nishikage T, Otani S, Lang RM. Age-related changes in left ventricular twist assessed by two-dimensional speckle-tracking imaging. 2006;19:1077-84
Takeuchi M, Otani S, Weinert L, Spencer KT, Lang RM. Comparison of contrast-enhanced real-time live 3-dimensional dobutamine stress echocardiography with contrast 2-dimensional echocardiography for detecting stress-induced wall-motion abnormalities. 2006;19:294-9
Takeuchi M, Yoshitani H, Miyazaki C, Yoshikawa J. Relationship between the number of coronary risk factors and coronary atherosclerosis assessed by high-frequency transthoracic echocardiography. 2006;19:1056-62
Talini E (*see* Di Bello et al). 2006;19:1063-71
Tamita K (*see* Katayama et al). 2006;19:335-40
Tan HL (*see* Hardziyenka et al). 2006;19:1272-9
Tanabe M (*see* Yamanaka-Funabiki et al). 2006;19:1332-7
Tanaka H, Kawai H, Tatsumi K, Kataoka T, Onishi T, Nose T, Mizoguchi T, Yokoyama M. Improved regional myocardial diastolic function assessed by strain rate imaging in patients with coronary artery disease undergoing percutaneous coronary intervention. 2006;19:756-62
Tarelli G (*see* Moreo et al). 2006;19:938 (Case rep.)
Tasset MR (*see* Silverstein et al). 2006;19:1529.e5-e8 (Case rep.)
Tatsumi K (*see* Tanaka et al). 2006;19:756-62
Tcheuffa JC (*see* Meimoun et al). 2006;19:1220-8
ten Cate FJ (*see* Nemes et al). 2006;19:108 (Case rep.)
ten Cate FJ (*see* Nemes et al). 2006;19:355 (Case rep.)
ten Cate FJ (*see* Pedone et al). 2006;19:982-6
Ten Harkel D-J (*see* van den Bosch et al). 2006;19:7-13
Ten Harkel D-J (*see* van den Bosch et al). 2006;19:815-21
Texler M (*see* Edibam et al). 2006;19:938.e1-e4 (Case rep.)
Th Kremastinos D (*see* Paraskevaidis et al). 2006;19:529-35
Theres H (*see* Eddicks et al). 2006;19:1073.e7-e9 (Case rep.)
Thiele K (*see* Pemberton et al). 2006;19:1403-10 (Review article)
Thiele K (*see* Sugeng et al). 2006;19:413-21
Thomas C (*see* McCulloch et al). 2006;19:744-8
Thomas JD (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Thomas JD (*see* Neilan et al). 2006;19:1280-5
Thomas JD (*see* Shin et al). 2006;19:796-801
Thomas JD (*see* Song et al). 2006;19:1129-35
Thune JJ, Køber L, Pfeffer MA, Skali H, Anavekar NS, Bourgoun M,

Ghali JK, Arnold JMO, Velazquez EJ, Solomon SD. Comparison of regional versus global assessment of left ventricular function in patients with left ventricular dysfunction, heart failure, or both after myocardial infarction: The valsartan in acute myocardial infarction echocardiographic study. 2006;19:1462-5
Toda I (*see* Okura et al). 2006;19:552-8
Tokai K (*see* Hirata et al). 2006;19:165-71
Tolstrup K (*see* Kobal et al). 2006;19:1012-6
Tomita J (*see* Akiyama et al). 2006;19:55-63
Tonari S (*see* Ito et al). 2006;19:987-93
Tong KL (*see* Kalvaitis et al). 2006;19:1488-93
Ton-Nu T-T (*see* Neilan et al). 2006;19:1280-5
Toole RS (*see* Mihalatos et al). 2006;19:1158-64
Topaloglu S (*see* Ozeke et al). 2006;19:1245-50
Topaloglu S (*see* Tufekcioglu et al). 2006;19:1320-5
Torp H (*see* Malm et al). 2006;19:365-72
Torp H (*see* Malm et al). 2006;19:40-7
Tosoratti E, Dall'Armellina E, Badano LP, Frassani R, Mazzaro E, Zakja E, Livi U, Fioretti PM. Heart failure and severe pulmonary hypertension caused by distal detachment of the valve conduit 16 years after the Cabrol composite graft procedure. 2006;19:1190.e5-e8 (Case rep.)
Toulany A, Shea S, Warren AE. Doppler tissue, strain, and strain rate imaging in pediatric patients with alström syndrome: Are there regional functional abnormalities?. 2006;19:14-20
Tousignant CP (*see* David et al). 2006;19:329-34
Toyoda T (*see* Akiyama et al). 2006;19:55-63
Toyota E (*see* Kume et al). 2006;19:914-8
Toyota E (*see* Watanabe et al). 2006;19:71-5
Tran H (*see* Shin et al). 2006;19:796-801
Tremblay TM (*see* Main et al). 2006;19:1038-44
Tribouilloy C (*see* Meimoun et al). 2006;19:1220-8
Troianos CA (*see* Mathew et al). 2006;19:1303-13 (Letter to the editor)
Tsai T-N, Tzeng B-H, Hong G-J, Chu K-M, Cheng S-M, Yang S-P. Perforated periaortic valve abscess presents as acute pericarditis and patient died in transesophageal echocardiographic study. 2006;19:579.e1-e3 (Case rep.)
Tseng J-C (*see* Gin et al). 2006;19:441-9
Tsiapras D (*see* Paraskevaidis et al). 2006;19:249-54
Tsiapras D (*see* Paraskevaidis et al). 2006;19:529-35
Tsujino H (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Tsutsui H (*see* Goto et al). 2006;19:857-64
Tsutsui JM (*see* Elhendy et al). 2006;19:1482-7
Tufekcioglu O (*see* Cay et al). 2006;19:1286-92 (Case rep.)
Tufekcioglu O (*see* Ozeke et al). 2006;19:1245-50
Tufekcioglu O, Aras D, Ozeke O, Maden O, Topaloglu S. Comparison of regional systolic myocardial velocities in patients with isolated left ventricular noncompaction and patients with idiopathic dilated cardiomyopathy. 2006;19:1320-5
Tuna F (*see* Cay et al). 2006;19:1286-92 (Case rep.)
Turkmen M (*see* Karakaya et al). 2006;19:434-7
Turkyilmaz E (*see* Karakaya et al). 2006;19:434-7
Tyler R (*see* Lester et al). 2006;19:919-23
Tzeng B-H (*see* Tsai et al). 2006;19:579.e1-e3 (Case rep.)
Tzou WS (*see* Gepner et al). 2006;19:223-8
Tzou WS, Korcarz CE, Aeschlimann SE, Stein JH. Use of hand-held ultrasound by a nonsonographer clinician to measure carotid intima-media thickness. 2006;19:1286-92

U

Ueda Y (*see* Katayama et al). 2006;19:335-40
Uehara H (*see* Hirano et al). 2006;19:536-9
Ujino K (*see* Hillis et al). 2006;19:450-6
Ulrich JN, Hesse B, Schuele S, Vlassak I, Sila CA, Jaber WA. Single-vessel versus multivessel territory acute ischemic stroke: Value of transesophageal echocardiography in the differentiation of embolic stroke. 2006;19:1165-9
Ulupinar H (*see* Cay et al). 2006;19:1286-92 (Case rep.)
Ulupinar H (*see* Ozeke et al). 2006;19:1245-50
Uno K, Takenaka K, Asada K, Ebihara A, Sasaki K, Komuro T, Nagai R, Motomura N, Ono M, Takamoto S. Diagnosis of Subacute Cardiac Rupture by Contrast Echocardiography. 2006;19:1401.e5-e7 (Case rep.)
Urheim S (*see* Wang et al). 2006;19:1102-11
Uslu N, Gorgulu S, Alper AT, Eren M, Nurkalem Z, Yildirim A, Ozer O. Erectile dysfunction as a generalized vascular dysfunction. 2006;19:341-6
Uzun M (*see* Genc et al). 2006;19:940 (Letter)

V

Valdes-Cruz L (*see* Dyer et al). 2006;19:403-12
Valdes-Cruz LM (*see* Dyer et al). 2006;19:21-7
Valente C (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
van Campen L (*see* Klein et al). 2006;19:763-71
van Dantzig JM (*see* Jansen et al). 2006;19:185-91
van den Bosch AE, Robbers-Visser D, Krenning BJ, Voormolen MM, McGhie JS, Helbing WA, Roos-Hesselink JW, Simoons ML, Meijboom FJ. Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease. 2006;19:1-6
van den Bosch AE, Ten Harkel D-J, McGhie JS, Roos-Hesselink JW, Simoons ML, Bogers AJJC, Meijboom FJ. Characterization of atrial septal defect assessed by real-time 3-dimensional echocardiography. 2006;19:815-21
van den Bosch AE, Ten Harkel D-J, McGhie JS, Roos-Hesselink JW, Simoons ML, Bogers AJJC, Meijboom FJ. Feasibility and accuracy of real-time 3-dimensional echocardiographic assessment of ventricular septal defects. 2006;19:7-13
van Hemel NM (*see* Jansen et al). 2006;19:185-91
Varadarajan P, Sharma S, Heywood JT, Pai RG. High prevalence of clinically silent severe mitral regurgitation in patients with heart failure: Role for echocardiography. 2006;19:1458-61
Vassilikos V (*see* Karvounis et al). 2006;19:1150-7
Veinot JP (*see* Peters et al). 2006;19:1530.e7-e11 (Case rep.)
Velazquez EJ (*see* Thune et al). 2006;19:1462-5
Vennalaganti PR, Ostfeld RJ, Malhotra D, Doddamani S, Gordon GM, Spevack DM. Pseudodyskinesis-a novel mechanism for abnormal motion of the heart's diaphragmatic wall. 2006;19:1286-92 (Case rep.)
Venugopal K (*see* Sunil Roy et al). 2006;19:1402.e1-e4 (Case rep.)
Venugopal KN (*see* Sankar et al). 2006;19:241 (Letter)
Veronesi F (*see* Corsi et al). 2006;19:886-93
Verzuri MS (*see* Mumoli et al). 2006;19:1286-92 (Case rep.)
Vidovich MI, Smulevitz B. Haunted liver. 2005;19:1302 (Letter to the editor)
Vieira MLC (*see* Cordovil et al). 2006;19:464-7 (Case rep.)
Vieira MLC (*see* Maddukuri et al). 2006;19:1026-32
Vincent PS (*see* Sachdev et al). 2006;19:34-9
Visser CA (*see* Dijkmans et al). 2006;19:285-93
Visser CA (*see* Klein et al). 2006;19:763-71
Visser FC (*see* Dijkmans et al). 2006;19:285-93
Visser FC (*see* Klein et al). 2006;19:763-71
Vitali Serdoz L (*see* Bahlmann et al). 2006;19:578.e5-e8 (Case rep.)
Vlassak I (*see* Ulrich et al). 2006;19:1165-9
Vletter WB (*see* Geleijnse et al). 2006;19:1076 (Letter)
Vletter WB (*see* Pedone et al). 2006;19:982-6
Voelker W (*see* Weidemann et al). 2006;19:839-47
Voigt J-U. Response to: "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet." 2006;19:837 (Letter)
Voon W-C (*see* Su et al). 2006;19:274-9
Voormolen MM (*see* van den Bosch et al). 2006;19:1-6
Vragel S (*see* Bangalore et al). 2006;19:438-40
Vyas H, Cabalka AK. Extreme redundancy of the valve of the fossa ovalis with right heart hypoplasia in a neonate with trisomy 18. 2006;19:1286-92 (Case rep.)

W

Wacker C (*see* Weidemann et al). 2006;19:839-47
Waclawiw MA (*see* Sachdev et al). 2006;19:34-9

Wada N (*see* Kume et al). 2006;19:914-8
Wada N (*see* Watanabe et al). 2006;19:71-5
Waggoner AD (*see* Bierig et al). 2006;19:471-4 (AMERICAN SOCIETY OF ECHOCARDIOGRAPHY POSITION PAPER)
Waggoner AD (*see* Dong et al). 2006;19:1017-25
Waggoner AD (*see* Peterson et al). 2006;19:749-55
Waggoner AD (*see* Rovner et al). 2006;19:872-9
Waggoner AD, Rovner A, de las Fuentes L, Faddis MN, Gleva MJ, Sawhney N, Dávila-Román VG. Clinical outcomes after cardiac resynchronization therapy: importance of left ventricular diastolic function and origin of heart failure. 2006;19:307-13
Waiss M-P (*see* Jung et al). 2006;19:802-10
Wald RM, Juraszek AL, Pigula FA, Geva T. Echocardiographic diagnosis and management of bilateral coronary ostial atresia in a patient with pulmonary atresia and intact ventricular septum. 2006;19:939.e4-36 (Case rep.)
Waldenström A (*see* Lindqvist et al). 2006;19:1345-9
Waltman TJ (*see* Malouf et al). 2006;19:464-7
Wandt B (*see* Nilsson et al). 2006;19:499-506
Wang J, Urheim S, Korinek J, Abraham TP, McMahon EM, Belohlavek M. Analysis of postsystolic myocardial thickening work in selective myocardial layers during progressive myocardial ischemia. 2006;19:1102-11
Wang W-C (*see* Gin et al). 2006;19:441-9
Wang W-C (*see* Hsiao et al). 2006;19:902-10
Warnecke G (*see* Bara et al). 2006;19:563-8
Warren AE (*see* Toulany et al). 2006;19:14-20
Watanabe H (*see* Hirata et al). 2006;19:165-71
Watanabe N (*see* Akiyama et al). 2006;19:55-63
Watanabe N (*see* Kume et al). 2006;19:914-8
Watanabe N, Ogasawara Y, Yamaura Y, Yamamoto K, Wada N, Kawamoto T, Toyota E, Akasaka T, Yoshida K. Geometric differences of the mitral valve tenting between anterior and inferior myocardial infarction with significant ischemic mitral regurgitation: Quantitation by novel software system with transthoracic real-time three-dimensional echocardiography. 2006;19:71-5
Webber GV, Fallon J, Silbiger JJ. Hematoma of a congenitally bicuspid aortic valve in a patient with polycythemia vera and the antiphospholipid antibody syndrome. 2006;19:1530.e1-e3 (Case rep.)
Weber S (*see* Meune et al). 2006;19:1466-70
Wei K (*see* Kalvaitis et al). 2006;19:1488-93
Weidemann F, Wacker C, Rauch A, Bauer WR, Bijnens B, Sutherland GR, Ertl G, Voelker W, Fidler F, Strotmann JM. Sequential changes of myocardial function during acute myocardial infarction, in the early and chronic phase after coronary intervention described by ultrasonic strain rate imaging. 2006;19:839-47
Weinberger J (*see* LaRocca et al). 2006;19:464-7 (Case rep.)
Weinert L (*see* Corsi et al). 2006;19:886-93
Weinert L (*see* Mor-Avi et al). 2006;19:48-54
Weinert L (*see* Sugeng et al). 2006;19:413-21
Weinert L (*see* Takeuchi et al). 2006;19:294-9
Weinert LC (*see* Jacobs et al). 2006;19:464-7 (Case rep.)
Weiss RJ (*see* Mor-Avi et al). 2006;19:48-54
Wencker D (*see* Mena et al). 2006;19:1295-300 (Review article)
Wichter T (*see* Bruch et al). 2006;19:522-8
Wichter T (*see* Bruch et al). 2006;19:95-101
Williams RJ (*see* Peterson et al). 2006;19:1393-400
Winter S (*see* Nesser et al). 2006;19:1286-92 (Case rep.)
Wiseth R (*see* Malm et al). 2006;19:365-72
Wittram C (*see* Jassal et al). 2006;19:229 (Case rep.)
Wong BL (*see* Markham et al). 2006;19:865-71
Wood LA (*see* Lester et al). 2006;19:919-23
Wood MJ (*see* Neilan et al). 2006;19:1280-5
Woods TD, Patel A. A critical review of patent foramen ovale detection using saline contrast echocardiography: When bubbles lie. 2006;19:215-22
Wu EB (*see* Zhang et al). 2006;19:422-8
Wu JC (*see* Leeper et al). 2006;19:1191.e5-e7 (Case rep.)
Wu L (*see* Zhang et al). 2006;19:422-8
Wyman RA (*see* Gepner et al). 2006;19:1170-4

X

Xie F (*see* Elhendy et al). 2006;19:1482-7

Y

Yagi T (*see* Katayama et al). 2006;19:335-40
Yamabe K (*see* Katayama et al). 2006;19:335-40
Yamada S (*see* Goto et al). 2006;19:857-64
Yamagishi H (*see* Hirata et al). 2006;19:165-71
Yamamoto K (*see* Watanabe et al). 2006;19:71-5
Yamamuro A (*see* Katayama et al). 2006;19:335-40
Yamanaka-Funabiki K, Onishi K, Tanabe M, Dohi K, Ito M, Ohte N, Nobori T, Nakano T. Single beat determination of regional myocardial strain measurements in patients with atrial fibrillation. 2006;19:1332-7
Yamasaki S (*see* Okamoto et al). 2006;19:211-4
Yamaura Y (*see* Watanabe et al). 2006;19:71-5
Yang H, Davidson WR Jr., Chambers CE, Pae WE, Sun B, Campbell DB, Pu M. Preoperative pulmonary hypertension is associated with postoperative left ventricular dysfunction in chronic organic mitral regurgitation: An echocardiographic and hemodynamic study. 2006;19:1051-5
Yang HS, Pellikka PA, McCully RB, Oh JK, Kukuzke JA, Khandheria BK, Chandrasekaran K. Role of biplane and biplane echocardiographically guided 3-dimensional echocardiography during dobutamine stress echocardiography. 2006;19:1136-43
Yang S-H (*see* Gin et al). 2006;19:441-9
Yang S-H (*see* Hsiao et al). 2006;19:822-8
Yang S-H (*see* Hsiao et al). 2006;19:902-10
Yang S-P (*see* Tsai et al). 2006;19:579.e1-e3 (Case rep.)
Yavas O (*see* Can et al). 2006;19:1074.e1-e2 (Case rep.)
Yen H-W (*see* Su et al). 2006;19:274-9
Yeo T-C (*see* Seow et al). 2006;19:102-6
Yeon S (*see* Han and Yeon). 2006;19:1529.e5-e8 (Case rep.)
Yildirim A (*see* Uslu et al). 2006;19:341-6
Yildiz A (*see* Ozeke et al). 2006;19:1245-50
Yip GW (*see* Zhang et al). 2006;19:422-8
Yoerger DM (*see* Neilan et al). 2006;19:1280-5
Yokoyama M (*see* Tanaka et al). 2006;19:756-62
Yoon DA (*see* Cardello et al). 2006;19:1074.e3-e4 (Case rep.)
Yoshida K (*see* Akiyama et al). 2006;19:55-63
Yoshida K (*see* Kume et al). 2006;19:914-8
Yoshida K (*see* Watanabe et al). 2006;19:71-5
Yoshida N (*see* Okamoto et al). 2006;19:211-4
Yoshikawa J (*see* Hirata et al). 2006;19:165-71
Yoshikawa J (*see* Okura et al). 2006;19:552-8
Yoshikawa J (*see* Takagi et al). 2006;19:1001-11
Yoshikawa J (*see* Takeuchi et al). 2006;19:1056-62
Yoshitani H (*see* Takeuchi et al). 2006;19:1056-62
Yoshiyama M (*see* Hirata et al). 2006;19:165-71
Youn H-J, Ihm S-H, Lee J-M, Park C-S, Cho E-J, Jung H-O, Jeon H-K, Oh Y-S, Chung W-S, Kim J-H, Choi K-B, Hong S-J. Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: Study using transthoracic Doppler echocardiography. 2006;19:373-8
Young CE (*see* Olson et al). 2006;19:83-7
Yu C-M (*see* Zhang et al). 2006;19:422-8
Yu EHC (*see* Basic et al). 2006;19:1203-10
Yu EHC (*see* Nair et al). 2006;19:924-31
Yumi Hayashi S, Lind B, Seeberger A, Mazza do Nascimento M, Lindholm B, Brodin L. Analysis of mitral annulus motion measurements derived from M-mode, anatomic M-mode, tissue Doppler displacement, and 2-dimensional strain imaging. 2006;19:1092-101

Z

Zaglavara T, Karvounis HI, Haaverstad R, Pillay TM, Hamilton JRL, Hasan A, Parharidis GE, Louridas GE, Dark JH, Kenny A. Dobutamine stress echocardiography is highly accurate for the prediction of contractile reserve in the early postoperative period, but may underestimate late recovery in contractile reserve after revascularization of the hibernating myocardium. 2006;19:300-6
Zakja E (*see* Tosoratti et al). 2006;19:1190.e5-e8 (Case rep.)
Zamorano J (*see* Perez de Isla et al). 2006;19:1338-44
Zarbatany D (*see* Naqvi et al). 2006;19:147-53
Zelinger A (*see* Lavi et al). 2006;19:1530.e5-e6 (Case rep.)
Zetts AD (*see* Asada-Kamiguchi et al). 2006;19:1112-8
Zhang Q, Kum LCC, Lee P-W, Lam Y-Y, Wu EB, Lin H, Yip GW, Wu L, Yu C-M. Effect of age and heart rate on atrial mechanical function assessed by Doppler tissue imaging in healthy individuals. 2006;19:422-8
Zisman D (*see* Shiran et al). 2006;19:578.e1-e4 (Case rep.)
Zo J-H (*see* Sohn et al). 2006;19:777-80
Zoghbi GJ (*see* Aqel et al). 2006;19:1072.e1-e3 (Case rep.)
Zoghbi WA (*see* McCulloch et al). 2006;19:744-8
Zoghbi WA (*see* Ramasubbu et al). 2006;19:464-7 (Case rep.)

Index to Subjects*

A

Ablation; *see* Catheter ablation

Abscess
Perforated periaortic valve abscess presents as acute pericarditis and patient died in transesophageal echocardiographic study (Tsai et al). 2006;19:579.e1-3 (Case rep.)

Abstracts
Abstracts. 2006;19:581-670

Acoustics
Quantification of regional left ventricular wall motion from real-time 3-dimensional echocardiography in patients with poor acoustic windows: effects of contrast enhancement tested against cardiac magnetic resonance (Corsi et al). 2006;19:886-93

Actinomycosis
Cardiac actinomycosis: an unusual cause of an intracardiac mass (Peters et al). 2006;19:1530.e7-11 (Case rep.)

Adenosine
Quantitative adenosine real-time myocardial contrast echocardiography for detection of angiographically significant coronary artery disease (Malm et al). 2006;19:365-72
Transthoracic coronary flow velocity reserve assessment: comparison between adenosine and dobutamine (Meimoun et al). 2006;19:1220-8

Adenosine triphosphate
Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11

Adrenalectomy
Reduced myocardial contractility assessed by tissue Doppler echocardiography is associated with increased risk during adrenal surgery of patients with pheochromocytoma: report of a preliminary study (Meune et al). 2006;19:1466-70

Age factors
Age-related changes in left ventricular twist assessed by two-dimensional speckle-tracking imaging (Takeuchi et al). 2006;19:1077-84
Effect of age and heart rate on atrial mechanical function assessed by Doppler tissue imaging in healthy individuals (Zhang et al). 2006;19:422-8
Effect of aging on twist-displacement loop by 2-dimensional speckle tracking imaging (Nakai et al). 2006;19:880-5

Aged
Atrial and ventricular echocardiographic correlates of the extent of pulmonary embolism in the elderly (Chung et al). 2006;19:347-53
Mitral annulus motion as a predictor of mortality in a community-based sample of 75-year-old men and women (Hedberg et al). 2006;19:88-94
Stress echocardiography in octogenarians: transesophageal atrial pacing is accurate, safe, and well tolerated (Kobal et al). 2006;19:1012-6

Alström syndrome
Doppler tissue, strain, and strain rate imaging in pediatric patients with Alström syndrome: are there regional functional abnormalities? (Toulany et al). 2006;19:14-20

American Society of Echocardiography
American Society of Echocardiography minimum standards for the cardiac sonographer: a position paper (Bierig et al). 2006;19:471-4
American Society of Echocardiography/Society of Cardiovascular Anesthesiologists recommendations and guidelines for continuous quality improvements in perioperative echocardiography (Mathew et al). 2006;19:1303-13
Clinical application of noninvasive vascular ultrasound in cardiovascular risk stratification: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Roman et al). 2006;19:943-54 (ASE report)
Guidelines for noninvasive vascular laboratory testing: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Gerhard-Herman et al). 2006;19:955-72 (ASE report)

Amyloidosis
Isolated left atrial amyloidosis: acute premitral stenosis secondary to spontaneous intramural left atrial hemorrhagic dissection (Edibam et al). 2006;19:938.e1-4 (Case rep.)

Anaphylaxis
Midventricular hypokinesis as a cardiac manifestation of anaphylaxis: a case report (Han and Yeon). 2006;19:1529.e9-11 (Case rep.)

Anesthesia
Impact of anesthesia on echocardiographic evaluation of systolic and diastolic function in rats (Plante et al). 2006;19:1520-5
Transesophageal echocardiographic acquisition of mitral annulus motion during monitoring of left ventricular function in anesthetized patients (Nilsson et al). 2006;19:499-506

Aneurysm, dissecting
Pulmonary artery dissection: a rare complication of pulmonary balloon valvuloplasty diagnosed 11 years after the procedure (Janus et al). 2006;19:1191.e5-8 (Case rep.)
Real-time 3-dimensional echocardiographic evaluation of aortic dissection (Nemes et al). 2006;19:108.e1-3 (Case rep.)
An unusual cause of pulmonary edema: acute rupture of noncoronary sinus of Valsalva aneurysm into the left atrium (Moreo et al). 2006;19:938.e9-11 (Case rep.)

Angioplasty
Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: a doppler echocardiographic study (Luca et al). 2006;19:172-7

Angioplasty, balloon
Improved regional myocardial diastolic function assessed by strain rate imaging in patients undergoing percutaneous coronary intervention (Tanaka et al). 2006;19:756-62
Sequential changes of myocardial function during acute myocardial infarction, in the early and chronic phase after coronary intervention described by ultrasonic strain rate imaging (Weidemann et al). 2006;19:839-47
Transesophageal echocardiographic diagnosis of *Aspergillus fumigatus* aortitis after percutaneous coronary intervention (Khumri et al). 2006;19:1072.e9-11 (Case rep.)

Angioplasty, transluminal, percutaneous coronary
Noninvasive diagnosis of restenosis by transthoracic doppler echocardiography after percutaneous coronary intervention: comparison with exercise Tl-SPECT (Hirata et al). 2006;19:165-71

Anthracyclines
The tei index identifies a differential effect on left and right

Indexes to the Seventeenth Annual Scientific Sessions abstracts are located in the May issue.

* January, pp 1-112; February, pp 121-242; March, pp 243-364; April, pp 365-470; May, pp. 471-732; June, pp. 733-838; July, pp. 839-942; August, pp. 943-1076; September, pp. 1077-1202; October, pp. 1203-1302; November, pp. 1303-1412; December, pp 1413-1580.

ventricular function with low-dose anthracycline chemotherapy (Belham et al). 2006;19:206-10

Antiphospholipid syndrome

Hematoma of a congenitally bicuspid aortic valve in a patient with polycythemia vera and the antiphospholipid antibody syndrome (Webber et al). 2006;19:1530.e1-3 (Case rep.)

Spontaneous retraction of an intramyocardial dissecting hemorrhage and multiple left ventricular thrombus formation in subacute myocardial infarction and antiphospholipid syndrome: a case report with long-term follow-up (Bahlmann et al). 2006;19:578.e5-8 (Case rep.)

Aorta

Aorta-to-left ventricle tunnel associated with noncompaction left ventricle (Sadeghpour et al). 2006;19:1073.e1-5 (Case rep.)

Common origin of the innominate and carotid arteries: prevalence, nomenclature, and surgical implications (Katz et al). 2006;19:1446-8

Impact of type 2 diabetes mellitus on aortic elastic properties in normotensive diabetes: Doppler tissue imaging study (Badran and Elnoaman). 2006;19:1471-81

Intracoronary stent visualized on transesophageal echocardiogram in a case of coronary dissection complicated by aortic dissection (Pohlel et al). 2006;19:229.e1-3 (Case rep.)

Unusual complication of aortic root reconstruction with sparing of the aortic valve: left ventricular outflow tract to right atrial fistula (Ramasubbu et al). 2006;19:469.e5-9 (Case rep.)

Aorta, thoracic

Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11

Stent graft in treatment of penetrating thoracic aortic ulcer (Khamaisi et al). 2006;19:835.e4-6 (Case rep.)

Thoracic aorta evaluation in patients with Takayasu's arteritis by transesophageal echocardiography (Lira-Filho et al). 2006;19:829-34

Aortic aneurysm

Real-time 3-dimensional echocardiographic evaluation of aortic dissection (Nemes et al). 2006;19:108.e1-3 (Case rep.)

Aortic valve

Acquired aortic cusp function after chronic left ventricular assist device support (Banchs et al). 2006;19:1401.e1-3 (Case rep.)

Behçet's aortitis mimicking aortic valve endocarditis with subaortic complications (Shiran et al). 2006;19:578.e1-4 (Case rep.)

Hematoma of a congenitally bicuspid aortic valve in a patient with polycythemia vera and the antiphospholipid antibody syndrome (Webber et al). 2006;19:1530.e1-3 (Case rep.)

Three-dimensional transesophageal echocardiography in the evaluation of aortic valve destruction by endocarditis (Nemes et al). 2006;19:355.e13-4 (Case rep.)

Aortic valve insufficiency

Impact of anesthesia on echocardiographic evaluation of systolic and diastolic function in rats (Plante et al). 2006;19:1520-5

The role of left ventricular long-axis contraction in patients with asymptomatic aortic regurgitation (Paraskevaidis et al). 2006;19:249-54

Aortic valve stenosis

Acute improvement in myocardial function assessed by myocardial strain and strain rate after aortic valve replacement for aortic stenosis (Iwahashi et al). 2006;19:1238-44

Rate of progression of valvular aortic stenosis in patients undergoing dialysis (Kume et al). 2006;19:914-8

Aortitis

Behçet's aortitis mimicking aortic valve endocarditis with subaortic complications (Shiran et al). 2006;19:578.e1-4 (Case rep.)

Transesophageal echocardiographic diagnosis of *Aspergillus fumigatus* aortitis after percutaneous coronary intervention (Khumri et al). 2006;19:1072.e9-11 (Case rep.)

Arrhythmia

Usefulness of clinical, electrocardiographic, and echocardiographic parameters to detect cardiac asynchrony in patients with left ventricular dysfunction secondary to ischemic or nonischemic heart disease (Perez de Isla). 2006;19:1338-44; Perez de Isla et al). 2006;19:1338-44

Arteries

Echocardiographic diagnosis and management of bilateral coronary ostial atresia in a patient with pulmonary atresia and intact ventricular septum (Wald et al). 2006;19:939.e1-3 (Case rep.)

Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: study using transthoracic Doppler echocardiography (Youn et al). 2006;19:373-8

Sequential transthoracic ultrasound assessment of coronary flow reserve in patient with tako-tsubo syndrome (Citro et al). 2006;19:1402.e5-8 (Case rep.)

Three major coronary artery-to-left ventricular fistula: an unusual cause of a diastolic murmur at the apex (Roy et al). 2006;19:1402.e1-4 (Case rep.)

Arteriovenous malformations

Pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia: an echocardiographic perspective (Jassal et al). 2006;19:229.e5-7

Artifacts

Continuous wave Doppler signal: a mystery (Enseleit et al). 2006;19:1191.e1-3 (Case rep.)

Ultrasound lung comets: a clinically useful sign of extravascular lung water (Picano et al). 2006;19:356-63 (Review article)

Ascites

The falciform ligament in the echocardiographic diagnosis of ascites (Cardello et al). 2006;19:1074.e3-4

Aspergillus fumigatus

Transesophageal echocardiographic diagnosis of *Aspergillus fumigatus* aortitis after percutaneous coronary intervention (Khumri et al). 2006;19:1072.e9-11 (Case rep.)

Atherosclerosis

Clinical application of noninvasive vascular ultrasound in cardiovascular risk stratification: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Roman et al). 2006;19:943-54 (ASE report)

Atherosclerosis, coronary; *see* Coronary arteriosclerosis

Athletes; *see* Sports

Atrial fibrillation

Single beat determination of regional myocardial strain measurements in patients with atrial fibrillation (Yamanaka-Funabiki et al). 2006;19:1332-7

Transesophageal echocardiographic assessment of right atrial appendage anatomy and function: comparison with the left atrial appendage and implications for local thrombus formation (Subramaniam et al). 2006;19:429-33

Atrial function

Effect of age and heart rate on atrial mechanical function assessed by Doppler tissue imaging in healthy individuals (Zhang et al). 2006;19:422-8

Atrial function, left

Effect of percutaneous mitral balloon valvuloplasty on left atrial appendage function: a Doppler tissue study (Karakaya et al). 2006;19:434-7

Hemodynamic change in transmitral gradient during Valsalva maneuver and its relation with left atrial function, functional capacity, and pulmonary artery pressure in patients with mitral stenosis in sinus rhythm (Ozeke et al). 2006;19:1245-50

Left atrial appendage function in patients with systemic embo-

lism in spite of in sinus rhythm (Okamoto et al). 2006;19: 211-4
Relationship between left atrial and left ventricular function in hypertrophic cardiomyopathy: a real-time 3-dimensional echocardiographic study (Shin et al). 2006;19:796-801

Atrial natriuretic factor
Exertional changes in circulating cardiac natriuretic peptides in patients with suggested coronary artery disease (Bergeron et al). 2006;19:772-6

Atrial septal defect; *see* Heart septal defects, atrial

Atrioventricular septal defect; *see* Heart septal defects, atrial

Automation
Automated quantification of the spatial extent of perfusion defects and viability on myocardial contrast echocardiography (Micari et al). 2006;19:379-85
Validation of a carotid intima-media thickness border detection program for use in an office setting (Gepner et al). 2006;19: 223-8

B

B-cell lymphoma; *see* Lymphoma, B-cell

B-type natriuretic peptide; *see* Natriuretic peptide, brain
Exertional changes in circulating cardiac natriuretic peptides in patients with suggested coronary artery disease (Bergeron et al). 2006;19:772-6
N-terminal pro B-type natriuretic peptide levels: correlation with echocardiographically determined left ventricular diastolic function in an ambulatory cohort (Dong et al). 2006; 19:1017-25

Backscatter
Obesity cardiomyopathy: is it a reality? An ultrasonic tissue characterization study (Di Bello et al). 2006;19:1063-71

Balloon dilatation
Effect of percutaneous mitral balloon valvuloplasty on left atrial appendage function: a Doppler tissue study (Karakaya et al). 2006;19:434-7
Pulmonary artery dissection: a rare complication of pulmonary balloon valvuloplasty diagnosed 11 years after the procedure (Janus et al). 2006;19:1191.e5-8 (Case rep.)

Balloon valvuloplasty; *see* Balloon dilatation

Behçet syndrome
Behçet's aortitis mimicking aortic valve endocarditis with subaortic complications (Shiran et al). 2006;19:578.e1-4 (Case rep.)

Biopsy
Transjugular biopsy of a right atrial mass under intracardiac echocardiography guidance (Aqel et al). 2006;19:1072.e5-8 (Case rep.)

Biplane echocardiography; *see* Echocardiography, biplane

Biventricular pacing; *see* Cardiac pacing, artificial

Blood flow velocity
Color tissue Doppler myocardial velocities consistently underestimate spectral tissue Doppler velocities: impact on calculation peak transmitral pulsed Doppler velocity/early diastolic tissue Doppler velocity (E/Ea) (McCulloch et al). 2006;19:744-8
Comparison of regional systolic myocardial velocities in patients with isolated left ventricular noncompaction and patients with idiopathic dilated cardiomyopathy (Tufekcioglu et al). 2006;19:1320-5
Coronary blood flow after cold exposure and the effect of warm-up exercise (Kim et al). 2006;19:386-90
Coronary flow velocity pattern assessed noninvasively by transthoracic color Doppler echocardiography serves as a predictor of adverse cardiac events and left ventricular remodeling in patients with acute myocardial infarction (Katayama et al). 2006;19:335-40
Corrected quantification method to determine myocardial blood flow using real-time myocardial contrast echocardiography (Otani et al). 2006;19:973-81
Effects of ultrasound contrast agents on doppler tissue velocity estimation (Ressner et al). 2006;19:154-64
Impact of preload and afterload on global and regional right ventricular function and pressure: a quantitative echocardiography study (Kjaergaard et al). 2006;19:515-21
Noninvasive diagnosis of ischemic and nonischemic cardiomyopathy using coronary flow velocity measurements of the left anterior descending coronary artery by transthoracic Doppler echocardiography (Okura et al). 2006;19:552-8
Prediction of transmural extent of infarction with contrast echocardiographically derived index of myocardial blood flow and myocardial blood volume fraction: comparison with contrast-enhanced magnetic resonance imaging (Choi et al). 2006;19:1211-9
Quantitative adenosine real-time myocardial contrast echocardiography for detection of angiographically significant coronary artery disease (Malm et al). 2006;19:365-72
Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: study using transthoracic Doppler echocardiography (Youn et al). 2006;19:373-8
Transthoracic coronary flow velocity reserve assessment: comparison between adenosine and dobutamine (Meimoun et al). 2006;19:1220-8
Tricuspid annular velocity in patients undergoing cardiac operation using transesophageal echocardiography (David et al). 2006;19:329-34

Blood pressure
Echocardiographic indices of increased left ventricular filling pressure and dilation after acute myocardial infarction (Hillis et al). 2006;19:450-6
The effect of acute hypertension on left ventricular diastolic pressures in a canine model of left ventricular dysfunction with a preserved ejection fraction and elevated left ventricular filling pressures (Lavine and Lavine). 2006;19:1350-8
Mitral peak Doppler E-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease (Sharma et al). 2006;19:266-73
Pulmonary hemodynamic effects of dipyridamole infusion in patients with normal and elevated pulmonary artery systolic pressure receiving PB127 (Main et al). 2006;19:1038-44

Blood vessel prosthesis
Heart failure and severe pulmonary hypertension caused by distal detachment of the valve conduit 16 years after the Cabrol composite graft procedure (Tosoratti et al). 2006;19: 1190.e5-8 (Case rep.)

Blood volume
Prediction of transmural extent of infarction with contrast echocardiographically derived index of myocardial blood flow and myocardial blood volume fraction: comparison with contrast-enhanced magnetic resonance imaging (Choi et al). 2006;19:1211-9

Blood flow velocity
Longitudinal isovolumic displacement of the left ventricular myocardium assessed by tissue velocity echocardiography in healthy individuals (Lind et al). 2006;19:255-65

Bosentan; *see* Sulfonamides

Brachiocephalic trunk
Common origin of the innominate and carotid arteries: prevalence, nomenclature, and surgical implications (Katz et al). 2006;19:1446-8

Bundle-branch block
Asynchrony of left ventricular systolic performance after the first acute myocardial infarction in patients with narrow QRS complexes: Doppler tissue imaging study (Elnoamany et al). 2006;19:1449-57
Effects of radial left ventricular dyssynchrony on cardiac performance using quantitative tissue Doppler radial strain imaging (Dohi et al). 2006;19:475-82
Left bundle branch block in chronic heart failure-impact on

diastolic function, filling pressures, and B-type natriuretic peptide levels (Bruch et al). 2006;19:95-101

C

Cabrol graft procedure; *see* Heart surgery
Calcification
Rate of progression of valvular aortic stenosis in patients undergoing dialysis (Kume et al). 2006;19:914-8
Carbazoles
The effect of carvedilol therapy on myocardial functional reserve in patients with advanced heart failure caused by nonischemic dilated cardiomyopathy (Paraskevaidis et al). 2006;19:529-35
Carcinoma, renal cell
Right ventricular metastasis caused by a renal cell carcinoma (Grawitz tumor): case report (Eddicks et al). 2006;19:1073.e11-5 (Case rep.)
Carcinoma, squamous cell
Right ventricular metastasis of uterine squamous cell carcinoma (Nesser et al). 2006;19:1294.e1-3 (Case rep.)
Cardiac catheterization; *see* Heart catheterization
Cardiac neoplasms; *see* Heart neoplasms
Cardiac output
Real-time three-dimensional color Doppler echocardiography overcomes the inaccuracies of spectral Doppler for stroke volume calculation (Pemberton et al). 2006;19:1403-10 (Review article)
Cardiac pacing, artificial
Effects of radial left ventricular dyssynchrony on cardiac performance using quantitative tissue Doppler radial strain imaging (Dohi et al). 2006;19:475-82
Stress echocardiography in octogenarians: transesophageal atrial pacing is accurate, safe, and well tolerated (Kobal et al). 2006;19:1012-6
Cardiac resynchronization therapy; *see also* Pacemaker, artificial
Clinical outcomes after cardiac resynchronization therapy: importance of left ventricular diastolic function and origin of heart failure (Waggoner et al). 2006;19:307-13
Improved papillary muscle function attenuates functional mitral regurgitation in patients with dilated cardiomyopathy after cardiac resynchronization therapy (Karvounis et al). 2006;19:1150-7
Cardiac rupture; *see* Heart rupture
Cardiac surgery; *see* Heart surgery
Cardiac tumor; *see* Heart neoplasms
Cardiac volume
Effect of preload reduction by hemodialysis on myocardial ultrasonic characterization, left atrial volume, and Doppler tissue imaging in patients with end-stage renal disease (Fijalkowski et al). 2006;19:1359-64
What is the best approach for the assessment of left atrial size? Comparison of various unidimensional and two-dimensional parameters with three-dimensional echocardiographically determined left atrial volume (Maddukuri et al). 2006;19:1026-32
What is the optimal clinical technique for measurement of left ventricular volume after myocardial infarction? A comparative study of 3-dimensional echocardiography, single photon emission computed tomography, and cardiac magnetic resonance imaging (Chan et al). 2006;19:192-201
Cardiomyopathies
Hypertrabeculation is not distinct from noncompaction in neuromuscular and nonneuromuscular disorders (Stöllberger and Finsterer). 2006;19:109-10 (Letter)
Obesity cardiomyopathy: is it a reality? An ultrasonic tissue characterization study (Di Bello et al). 2006;19:1063-71
Cardiomyopathy, congestive
Comparison of regional systolic myocardial velocities in patients with isolated left ventricular noncompaction and patients with idiopathic dilated cardiomyopathy (Tufekcioglu et al). 2006;19:1320-5
Cardiomyopathy, dilated
Abnormalities of diastolic function precede dilated cardiomyopathy associated with Duchenne muscular dystrophy (Markham et al). 2006;19:865-71
The effect of carvedilol therapy on myocardial functional reserve in patients with advanced heart failure caused by nonischemic dilated cardiomyopathy (Paraskevaidis et al). 2006;19:529-35
Improved papillary muscle function attenuates functional mitral regurgitation in patients with dilated cardiomyopathy after cardiac resynchronization therapy (Karvounis et al). 2006;19:1150-7
Cardiomyopathy, hypertrophic
Abnormal longitudinal myocardial functional reserve assessed by exercise tissue Doppler echocardiography in patients with hypertrophic cardiomyopathy (Ha et al). 2006;19:1314-9
Asymmetry of systolic anterior motion of the mitral valve in patients with hypertrophic obstructive cardiomyopathy: a real-time three-dimensional echocardiography study (Song et al). 2006;19:1129-35
Can transthoracic live 3-dimensional echocardiography improve the recognition of midventricular obliteration in hypertrophic obstructive cardiomyopathy? (Gregorio et al). 2006;19:1190.e1-4 (Case rep.)
Myocardial perfusion after percutaneous transluminal septal myocardial ablation as assessed by myocardial contrast echocardiography in patients with hypertrophic obstructive cardiomyopathy (Pedone et al). 2006;19:982-6
Regional postsystolic shortening in patients with hypertrophic cardiomyopathy: its incidence and characteristics assessed by strain imaging (Ito et al). 2006;19:987-93
Relationship between left atrial and left ventricular function in hypertrophic cardiomyopathy: a real-time 3-dimensional echocardiographic study (Shin et al). 2006;19:796-801
Role of left ventricular regional diastolic abnormalities for global diastolic dysfunction in patients with hypertrophic cardiomyopathy (Goto et al). 2006;19:857-64
Use of echocardiography in patients with hypertrophic cardiomyopathy: clinical implications of massive hypertrophy (Sorajja et al). 2006;19:788-95
Cardiomyopathy, restrictive
The systolic to diastolic duration ratio in children with heart failure secondary to restrictive cardiomyopathy (Friedberg and Silverman). 2006;19:1326-31
Cardiovascular diseases
Use of carotid intima-media thickness and vascular age to modify cardiovascular risk prediction (Gepner et al). 2006;19:1170-4
Cardioversion; *see* Electric countershock
Carotid artery
Common origin of the innominate and carotid arteries: prevalence, nomenclature, and surgical implications (Katz et al). 2006;19:1446-8
Use of carotid intima-media thickness and vascular age to modify cardiovascular risk prediction (Gepner et al). 2006;19:1170-4
Use of hand-held ultrasound by a nonsonographer clinician to measure carotid intima-media thickness (Tzou et al). 2006;19:1286-92
Validation of a carotid intima-media thickness border detection program for use in an office setting (Gepner et al). 2006;19:223-8
Carvedilol; *see* Carbazoles; Propanolamines
Case reports
Case reports. 2006;19:107.e1-6, 108.e1-3, 229.e1-3, 354.e1-3, 354.e5-8, 355.e9-11, 355.e13-4, 468.e1-7, 469.e1-3, 469.e5-9, 578.e1-8, 579.e1-7, 835.e1-6, 836.e1-8, 938.e1-7, 938.e9-11, 939.e1-3, 939.e5-7, 939.e9-11, 1072.e1-3, 1072.e5-11,

1073.e1-5, 1073.e7-9, 1073.e11-5, 1074.e1-6, 1190.e1-10, 1191.e1-3, 1191.e5-11, 1293.e1-8, 1294.e1-8, 1401.e1-11, 1402.e1-11, 1529.e1-3, 1530.e4-6

Catheter ablation

Myocardial perfusion after percutaneous transluminal septal myocardial ablation as assessed by myocardial contrast echocardiography in patients with hypertrophic obstructive cardiomyopathy (Pedone et al). 2006;19:982-6

Cavernous hemangioma; *see* Hemangioma, cavernous

Cerebrovascular disorders

Single-vessel versus multivessel territory acute ischemic stroke: value of transesophageal echocardiography in the differentiation of embolic stroke (Ulrich et al). 2006;19:1165-9

Chest pain

Effect of time delay on the diagnostic use of contrast echocardiography in patients presenting to the emergency department with chest pain and no S-T segment elevation (Kalvaitis et al). 2006;19:1488-93

Child

Doppler tissue, strain, and strain rate imaging in pediatric patients with Alström syndrome: are there regional functional abnormalities? (Toulany et al). 2006;19:14-20

Left ventricular Tei index in children: comparison of Doppler tissue imaging, pulsed wave Doppler, and M-mode echocardiography normal values (Cui and Roberson). 2006; 19:1438-45

Noninvasive Doppler tissue measurement of pulmonary artery compliance in children with pulmonary hypertension (Dyer et al). 2006;19:403-12

The systolic to diastolic duration ratio in children with heart failure secondary to restrictive cardiomyopathy (Friedberg and Silverman). 2006;19:1326-31

Ultrasonic strain rate and strain imaging of the right ventricle in children before and after percutaneous closure of an atrial septal defect (Eyskens et al). 2006;19:994-1000

Use of myocardial performance index in pediatric patients with idiopathic pulmonary arterial hypertension (Dyer et al). 2006;19:21-7

Clinical competence

The assessment of technical and interpretative proficiency in echocardiography (Nair et al). 2006;19:924-31

Clot retraction

Spontaneous retraction of an intramyocardial dissecting hemorrhage and multiple left ventricular thrombus formation in subacute myocardial infarction and antiphospholipid syndrome: a case report with long-term follow-up (Bahlmann et al). 2006;19:578.e5-8 (Case rep.)

Cold

Coronary blood flow after cold exposure and the effect of warm-up exercise (Kim et al). 2006;19:386-90

Collateral circulation

"Steal" collaterals: an echocardiographic diagnostic marker for anomalous origin of the left main coronary artery from the pulmonary artery in the adult (Pisacane et al). 2006;19: 107.e3-6 (Case rep.)

Community health services

The architecture of a mobile outreach echocardiography service (Gilman et al). 2006;19:1526-8

Complications, postoperative; *see* Postoperative complications

Computers

Geometric differences of the mitral valve tenting between anterior and inferior myocardial infarction with significant ischemic mitral regurgitation: quantitation by novel software system with transthoracic real-time three-dimensional echocardiography (Watanabe et al). 2006;19:71-5

Congenital heart defects; *see* Heart defects, congenital

Congestive heart failure; *see* Heart failure, congestive

Constrictive pericarditis; *see* Pericarditis, constrictive

Contractile reserve

Dobutamine stress echocardiography is highly accurate for the prediction of contractile reserve in the early postoperative period, but may underestimate late recovery in contractile reserve after revascularization of the hibernating myocardium (Zaglavara et al). 2006;19:300-6

The effect of carvedilol therapy on myocardial functional reserve in patients with advanced heart failure caused by nonischemic dilated cardiomyopathy (Paraskevaidis et al). 2006;19:529-35

Traditional and color M-mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: relations to contractility reserve (Palmieri et al). 2006;19:483-90

Contrast echocardiography; *see* Echocardiography, contrast

Contrast media

Color encoding of endocardial motion improves the interpretation of contrast-enhanced echocardiographic stress tests by less-experienced readers (Mor-avi et al). 2006;19:48-54

Contrast echocardiography: experience in a clinical echocardiography laboratory (Lester et al). 2006;19:919-23

Effects of ultrasound contrast during tissue velocity imaging on regional left ventricular velocity, strain, and strain rate measurements (Malm et al). 2006;19:40-7

Quantification of regional left ventricular wall motion from real-time 3-dimensional echocardiography in patients with poor acoustic windows: effects of contrast enhancement tested against cardiac magnetic resonance (Corsi et al). 2006;19:886-93

Cor triatriatum

Subtotal anomalous pulmonary venous connection and left cor triatriatum: a rare diagnosis in adulthood (Caputo et al). 2006;19:836.e1-4 (Case rep.)

Three-dimensional transthoracic echocardiography to evaluate cor triatriatum in the adult (Jacobs et al). 2006;19:468.e1-4 (Case rep.)

Coronary angiography

Quantitative adenosine real-time myocardial contrast echocardiography for detection of angiographically significant coronary artery disease (Malm et al). 2006;19:365-72

Coronary arteriosclerosis

Relationship between the number of coronary risk factors and coronary atherosclerosis assessed by high-frequency transthoracic echocardiography (Takeuchi et al). 2006;19: 1056-62

Coronary artery; *see* Arteries

Coronary artery bypass

Dynamic quantitative echocardiographic evaluation of mitral regurgitation in the operating department (Gisbert et al). 2006;19:140-6

The Impella Recover LP 5.0 left ventricular assist device: a bridge to coronary artery bypass grafting and cardiac transplantation (LaRocca et al). 2006;19:468.e5-7 (Case rep.)

Noninvasive diagnosis of coronary artery bypass graft disease by dobutamine stress real-time myocardial contrast perfusion imaging (Elhendy et al). 2006;19:1482-7

Coronary artery bypass grafting; *see* Coronary artery bypass

Coronary artery stenosis

Detection of residual infarct-related coronary artery stenosis and multivessel disease after thrombolysis: comparison between myocardial contrast echocardiography and single photon emission computed tomography (Dwivedi et al). 2006;19:546-51

Coronary circulation

Assessment of coronary vasodilatation in response to nitroglycerin with transthoracic Doppler echocardiography (Sohn et al). 2006;19:777-80

Coronary blood flow after cold exposure and the effect of warm-up exercise (Kim et al). 2006;19:386-90

Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow veloc-

ity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11
Impact of the coronary flow reduction at rest on myocardial perfusion and functional indices derived from myocardial contrast and strain echocardiography (Okuda et al). 2006;19:781-7
Sequential transthoracic ultrasound assessment of coronary flow reserve in patient with tako-tsubo syndrome (Citro et al). 2006;19:1402.e5-8 (Case rep.)
Transthoracic coronary flow velocity reserve assessment: comparison between adenosine and dobutamine (Meimoun et al). 2006;19:1220-8

Coronary disease
Efficacy of ultrasound-assisted stress testing using a hand-carried ultrasound device for diagnosis of coronary artery disease (Hirano et al). 2006;19:536-9
Exertional changes in circulating cardiac natriuretic peptides in patients with suggested coronary artery disease (Bergeron et al). 2006;19:772-6
Improved regional myocardial diastolic function assessed by strain rate imaging in patients undergoing percutaneous coronary intervention (Tanaka et al). 2006;19:756-62
Noninvasive diagnosis of coronary artery bypass graft disease by dobutamine stress real-time myocardial contrast perfusion imaging (Elhendy et al). 2006;19:1482-7
Prognostic value of mitral regurgitation assessment during exercise echocardiography in patients with known or suspected coronary artery disease (Peteiro et al). 2006;19:1229-37
Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested of known ischemic heart disease (Basic et al). 2006;19:1203-10
Quantitative adenosine real-time myocardial contrast echocardiography for detection of angiographically significant coronary artery disease (Malm et al). 2006;19:365-72
Risk stratification by treadmill exercise echocardiography (Peteiro et al). 2006;19:894-901
Traditional and color M-mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: relations to contractility reserve (Palmieri et al). 2006;19:483-90

Coronary restenosis
Noninvasive diagnosis of restenosis by transthoracic doppler echocardiography after percutaneous coronary intervention: comparison with exercise TI-SPECT (Hirata et al). 2006;19:165-71

Coronary stenosis
Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11
Noninvasive diagnosis of coronary artery bypass graft disease by dobutamine stress real-time myocardial contrast perfusion imaging (Elhendy et al). 2006;19:1482-7

Coronary vessel anomalies
"Steal" collaterals: an echocardiographic diagnostic marker for anomalous origin of the left main coronary artery from the pulmonary artery in the adult (Pisacane et al). 2006;19:107.e3-6 (Case rep.)

Coronary vessels
Heterogeneity of myocardial perfusion in distal coronary embolism with different particle sizes (Akiyama et al). 2006;19:55-63

Correspondence; *see* Letters to the editor

D

Death, sudden, cardiac
Syncope with a twist (Gosavi et al). 2006;19:1529.e1 (Case rep.)

Diabetes, gestational
Gestational diabetes: a condition of early diastolic abnormalities in young women (Freire et al). 2006;19:1251-6

Diabetes mellitus, insulin-dependent
Alterations in left ventricular structure and function in type-1 diabetics: a focus on left atrial contribution to function (Peterson et al). 2006;19:749-55
Left ventricular chamber and myocardial systolic function reserve in patients with type 1 diabetes mellitus: insight from traditional and Doppler tissue imaging echocardiography (Palmieri et al). 2006;19:848-56

Diabetes mellitus, non-insulin-dependent
Impact of type 2 diabetes mellitus on aortic elastic properties in normotensive diabetes: Doppler tissue imaging study (Badran and Elnoaman). 2006;19:1471-81

Diagnosis
The added value of real-time 3-dimensional echocardiography in the diagnosis of isolated cleft mitral valve in adults (Kuperstein et al). 2006;19:811-4
Detection of heart transplant rejection in adults by echocardiographic diastolic indices: a systematic review of the literature (Mena et al). 2006;19:1295-1300 (Review article)
Diagnosis of subacute cardiac rupture by contrast echocardiography (Uno et al). 2006;19:1401.e9-11 (Case rep.)
Echocardiographic diagnosis and management of bilateral coronary ostial atresia in a patient with pulmonary atresia and intact ventricular septum (Wald et al). 2006;19:939.e1-3 (Case rep.)
Efficacy of ultrasound-assisted stress testing using a hand-carried ultrasound device for diagnosis of coronary artery disease (Hirano et al). 2006;19:536-9
The falciform ligament in the echocardiographic diagnosis of ascites (Cardello et al). 2006;19:1074.e3-4
Hypertrabeculation is not distinct from noncompaction in neuromuscular and nonneuromuscular disorders (Stöllberger and Finsterer). 2006;19:109-10 (Letter)
Left ventricular outflow tract obstruction caused by accessory mitral valve tissue (Sharma et al). 2006;19:354.e5-8 (Case rep.)
Long-term survival of a patient with isolated noncompaction of the ventricular myocardium (Salemi et al). 2006;19:354.e1-3 (Case rep.)
Noninvasive diagnosis of ischemic and nonischemic cardiomyopathy using coronary flow velocity measurements of the left anterior descending coronary artery by transthoracic Doppler echocardiography (Okura et al). 2006;19:552-8
Noninvasive diagnosis of restenosis by transthoracic doppler echocardiography after percutaneous coronary intervention: comparison with exercise TI-SPECT (Hirata et al). 2006;19:165-71
Primary cardiac lymphoma: diagnosis and treatment guided by transesophageal echocardiogram perfusion imaging (Jang et al). 2006;19:1073.e7-9 (Case rep.)
Quantitative measures of right ventricular dysfunction by echocardiography in the diagnosis of acute nonmassive pulmonary embolism (Kjaergaard et al). 2006;19:1264-71
A rare complication of atrial septal occluders: diagnosis by transthoracic echocardiography (Dialetto et al). 2006;19:836.e5-8 (Case rep.)
Real-time 3-dimensional echocardiographic evaluation of aortic dissection (Nemes et al). 2006;19:108.e1-3 (Case rep.)
Sequential transthoracic ultrasound assessment of coronary flow reserve in patient with tako-tsubo syndrome (Citro et al). 2006;19:1402.e5-8 (Case rep.)
"Steal" collaterals: an echocardiographic diagnostic marker for anomalous origin of the left main coronary artery from the pulmonary artery in the adult (Pisacane et al). 2006;19:107.e3-6 (Case rep.)
Subtotal anomalous pulmonary venous connection and left cor triatriatum: a rare diagnosis in adulthood (Caputo et al). 2006;19:836.e1-4 (Case rep.)

Transesophageal echocardiographic diagnosis of *Aspergillus fumigatus* aortitis after percutaneous coronary intervention (Khumri et al). 2006;19:1072.e9-11 (Case rep.)

Diagnosis, computer-assisted

Accuracy and feasibility of online 3-dimensional echocardiography for measurement of left ventricular parameters (Jenkins et al). 2006;19:1119-28

Automated quantification of the spatial extent of perfusion defects and viability on myocardial contrast echocardiography (Micari et al). 2006;19:379-85

Dialysis

Rate of progression of valvular aortic stenosis in patients undergoing dialysis (Kume et al). 2006;19:914-8

Diaphragm

Pseudodyskinesis-a novel mechanism for abnormal motion of the heart's diaphragmatic wall (Vennalaganti et al). 2006;19:1294.e5-6 (Case rep.)

Diastole

Abnormalities of diastolic function precede dilated cardiomyopathy associated with Duchenne muscular dystrophy (Markham et al). 2006;19:865-71

Assessment of diastolic function with Doppler tissue imaging after cardiac surgery: effect of "postoperative septum" in on-pump and off-pump procedures (Malouf et al). 2006;19:464-7

Assessment of the evolution of normal fetal diastolic function during mid and late gestation by spectral Doppler tissue echocardiography (Nii et al). 2006;19:1431-7

Characterization of left ventricular diastolic function in hypertension by use of Doppler tissue imaging and color M-mode techniques (Rovner et al). 2006;19:872-9

Clinical outcomes after cardiac resynchronization therapy: importance of left ventricular diastolic function and origin of heart failure (Waggoner et al). 2006;19:307-13

Color tissue Doppler myocardial velocities consistently underestimate spectral tissue Doppler velocities: impact on calculation peak transmitral pulsed Doppler velocity/early diastolic tissue Doppler velocity (E/Ea) (McCulloch et al). 2006;19:744-8

Detection of heart transplant rejection in adults by echocardiographic diastolic indices: a systematic review of the literature (Mena et al). 2006;19:1295-1300 (Review article)

Doppler tissue, strain, and strain rate imaging in pediatric patients with Alström syndrome: are there regional functional abnormalities? (Toulany et al). 2006;19:14-20

Early mitral filling/diastolic mitral annular velocity ratio is not a reliable predictor of left ventricular filling pressure in the setting of severe mitral regurgitation (Olson et al). 2006;19:83-7

Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: a doppler echocardiographic study (Luca et al). 2006;19:172-7

Gestational diabetes: a condition of early diastolic abnormalities in young women (Freire et al). 2006;19:1251-6

Impact of anesthesia on echocardiographic evaluation of systolic and diastolic function in rats (Plante et al). 2006;19:1520-5

Improved regional myocardial diastolic function assessed by strain rate imaging in patients undergoing percutaneous coronary intervention (Tanaka et al). 2006;19:756-62

Intraventricular pressure gradients in left ventricular aneurysms determined by color M-mode Doppler method: an animal study (Asada-Kamiguchi et al). 2006;19:1112-8

Isolated ventricular diastolic dysfunction: implications for exercise left ventricular performance in patients without congestive heart failure (Palmieri et al). 2006;19:491-8

Left bundle branch block in chronic heart failure-impact on diastolic function, filling pressures, and B-type natriuretic peptide levels (Bruch et al). 2006;19:95-101

Mitral peak Doppler E-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease (Sharma et al). 2006;19:266-73

N-terminal pro B-type natriuretic peptide levels: correlation with echocardiographically determined left ventricular diastolic function in an ambulatory cohort (Dong et al). 2006;19:1017-25

Role of left ventricular regional diastolic abnormalities for global diastolic dysfunction in patients with hypertrophic cardiomyopathy (Goto et al). 2006;19:857-64

Single-beat differentiation among left ventricular filling pattern by pulsed wave Doppler echocardiography (Su et al). 2006;19:274-9

The systolic to diastolic duration ratio in children with heart failure secondary to restrictive cardiomyopathy (Friedberg and Silverman). 2006;19:1326-31

Three major coronary artery-to-left ventricular fistula: an unusual cause of a diastolic murmur at the apex (Roy et al). 2006;19:1402.e1-4 (Case rep.)

Traditional and color M-mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: relations to contractility reserve (Palmieri et al). 2006;19:483-90

Diastolic pressure; *see* Blood pressure

Dilated cardiomyopathy; *see* Cardiomyopathy, dilated

Dipyridamole

Pulmonary hemodynamic effects of dipyridamole infusion in patients with normal and elevated pulmonary artery systolic pressure receiving PB127 (Main et al). 2006;19:1038-44

Diverticulum

Left ventricular diverticulum with contractile function in an unusual site (Cay et al). 2006;19:1293.e3-6 (Case rep.)

Dobutamine

Comparison of contrast-enhanced real-time live 3-dimensional dobutamine stress echocardiography with contrast 2-dimensional echocardiography for detecting stress-induced wall-motion abnormalities (Takeuchi et al). 2006;19:294-9

Dobutamine stress echocardiography is highly accurate for the prediction of contractile reserve in the early postoperative period, but may underestimate late recovery in contractile reserve after revascularization of the hibernating myocardium (Zaglavara et al). 2006;19:300-6

Feasibility of using a real-time 3-dimensional technique for contrast dobutamine stress echocardiography (Pulerwitz et al). 2006;19:540-5

Hypotension during dobutamine stress transesophageal echocardiography: relationship with provoked left ventricular obstruction (Siddiqui et al). 2006;19:1144-9

Left ventricular chamber and myocardial systolic function reserve in patients with type 1 diabetes mellitus: insight from traditional and Doppler tissue imaging echocardiography (Palmieri et al). 2006;19:848-56

Noninvasive diagnosis of coronary artery bypass graft disease by dobutamine stress real-time myocardial contrast perfusion imaging (Elhendy et al). 2006;19:1482-7

Response to: "contrast-enhanced real-time 3-dimensional dobutamine stress echocardiography." (Geleijnse et al). 2006;19:1076 (Letter)

Role of biplane and biplane echocardiographically guided 3-dimensional echocardiography during dobutamine stress echocardiography (Yang et al). 2006;19:1136-43

Traditional and color M-mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: relations to contractility reserve (Palmieri et al). 2006;19:483-90

Transthoracic coronary flow velocity reserve assessment: comparison between adenosine and dobutamine (Meimoun et al). 2006;19:1220-8

Doppler echocardiography; *see* Echocardiography, Doppler

Doppler tissue imaging

Abnormal longitudinal myocardial functional reserve assessed

by exercise tissue Doppler echocardiography in patients with hypertrophic cardiomyopathy (Ha et al). 2006;19:1314-9

Accuracy, reproducibility, and comparability of Doppler tissue imaging by two high-end ultrasound systems (Kjaergaard et al). 2006;19:322-8

Analysis of mitral annulus motion measurements derived from M-mode, anatomic M-mode, tissue Doppler displacement, and 2-dimensional strain imaging (Hayashi et al). 2006;19:1092-1101

Assessment of diastolic function with Doppler tissue imaging after cardiac surgery: effect of "postoperative septum" in on-pump and off-pump procedures (Malouf et al). 2006;19:464-7

Assessment of the evolution of normal fetal diastolic function during mid and late gestation by spectral Doppler tissue echocardiography (Nii et al). 2006;19:1431-7

Asynchrony detection and Doppler tissue imaging (Gaibazzi). 2006;19:111 (Letter)

Asynchrony of left ventricular systolic performance after the first acute myocardial infarction in patients with narrow QRS complexes: Doppler tissue imaging study (Elnoamany et al). 2006;19:1449-57

Characterization of left ventricular diastolic function in hypertension by use of Doppler tissue imaging and color M-mode techniques (Rovner et al). 2006;19:872-9

Color tissue Doppler myocardial velocities consistenty underestimate spectral tissue Doppler velocities: impact on calculation peak transmitral pulsed Doppler velocity/early diastolic tissue Doppler velocity (E/Ea) (McCulloch et al). 2006;19:744-8

Doppler tissue, strain, and strain rate imaging in pediatric patients with Alström syndrome: are there regional functional abnormalities? (Toulany et al). 2006;19:14-20

Doppler tissue imaging in the assessment of atrioventricular conduction time: validation of a novel technique and comparison with electrophysiologic and pulsed wave Doppler-derived equivalents in an animal model (Nii et al). 2006;19:314-21

Effect of age and heart rate on atrial mechanical function assessed by Doppler tissue imaging in healthy individuals (Zhang et al). 2006;19:422-8

Effect of percutaneous mitral balloon valvuloplasty on left atrial appendage function: a Doppler tissue study (Karakaya et al). 2006;19:434-7

Effect of preload reduction by hemodialysis on myocardial ultrasonic characterization, left atrial volume, and Doppler tissue imaging in patients with end-stage renal disease (Fijalkowski et al). 2006;19:1359-64

Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: a doppler echocardiographic study (Luca et al). 2006;19:172-7

Effects of radial left ventricular dyssynchrony on cardiac performance using quantitative tissue Doppler radial strain imaging (Dohi et al). 2006;19:475-82

Effects of ultrasound contrast agents on doppler tissue velocity estimation (Ressner et al). 2006;19:154-64

Effects of ultrasound contrast during tissue velocity imaging on regional left ventricular velocity, strain, and strain rate measurements (Malm et al). 2006;19:40-7

Impact of the coronary flow reduction at rest on myocardial perfusion and functional indices derived from myocardial contrast and strain echocardiography (Okuda et al). 2006;19:781-7

Impact of type 2 diabetes mellitus on aortic elastic properties in normotensive diabetes: Doppler tissue imaging study (Badran and Elnoaman). 2006;19:1471-81

Improved regional myocardial diastolic function assessed by strain rate imaging in patients undergoing percutaneous coronary intervention (Tanaka et al). 2006;19:756-62

Left ventricular chamber and myocardial systolic function reserve in patients with type 1 diabetes mellitus: insight from traditional and Doppler tissue imaging echocardiography (Palmieri et al). 2006;19:848-56

Left ventricular systolic longitudinal function: comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals (Mondillo et al). 2006;19:1085-91

Left ventricular Tei index in children: comparison of Doppler tissue imaging, pulsed wave Doppler, and M-mode echocardiography normal values (Cui and Roberson). 2006;19:1438-45

Letters to the editor (Ogunyanklu) (Letter); (McCulloch and Dokainish) (Reply). 2006;19:1411-2

Longitudinal isovolumic displacement of the left ventricular myocardium assessed by tissue velocity echocardiography in healthy individuals (Lind et al). 2006;19:255-65

Noninvasive Doppler tissue measurement of pulmonary artery compliance in children with pulmonary hypertension (Dyer et al). 2006;19:403-12

Obesity cardiomyopathy: is it a reality? An ultrasonic tissue characterization study (Di Bello et al). 2006;19:1063-71

Optimization of pulsed wave tissue doppler to predict left ventricular reverse remodeling after cardiac resynchronization therapy (Jansen et al). 2006;19:185-91

Pulmonary embolism and right heart function: insights from myocardial Doppler tissue imaging (Hsiao et al). 2006;19:822-8

Quantitative echocardiographic analysis of the right ventricle in healthy individuals (Kjaergaard et al). 2006;19:1365-72

Reduced myocardial contractility assessed by tissue Doppler echocardiography is associated with increased risk during adrenal surgery of patients with pheochromocytoma: report of a preliminary study (Meune et al). 2006;19:1466-70

Response to: "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet." (Voigt). 2006;19:837 (Letter)

Response to comments on our article "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet" (Kjaergaard et al). 2006;19:1301 (Letter)

Right heart function in scleroderma: insights from myocardial Doppler tissue imaging (Hsiao et al). 2006;19:507-14

Right heart function in systemic lupus erythematosus: insights from myocardial Doppler tissue imaging (Gin et al). 2006;19:441-9

Single beat determination of regional myocardial strain measurements in patients with atrial fibrillation (Yamanaka-Funabiki et al). 2006;19:1332-7

Usefulness of ventricular longitudinal contractility assessed by doppler tissue imaging in the prediction of reverse remodeling in patients with severe left ventricular systolic dysfunction (Kang et al). 2006;19:178-84

Duchenne muscular dystrophy; *see* Muscular dystrophy, Duchenne

E

Echocardigraphy, M-mode

Response to: "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet." (Voigt). 2006;19:837 (Letter)

Echocardiography

Hypertrabeculation is not distinct from noncompaction in neuromuscular and nonneuromuscular disorders (Stöllberger and Finsterer). 2006;19:109-10 (Letter)

Echocardiography, biplane

Role of biplane and biplane echocardiographically guided 3-dimensional echocardiography during dobutamine stress echocardiography (Yang et al). 2006;19:1136-43

Echocardiography, child
Guidelines and standards for performance of a pediatric echocardiogram (Lai et al). 2006;19:1413-30
Echocardiography, contrast
Automated quantification of the spatial extent of perfusion defects and viability on myocardial contrast echocardiography (Micari et al). 2006;19:379-85
Comparison of contrast-enhanced real-time live 3-dimensional dobutamine stress echocardiography with contrast 2-dimensional echocardiography for detecting stress-induced wall-motion abnormalities (Takeuchi et al). 2006;19:294-9
Contrast echocardiography: experience in a clinical echocardiography laboratory (Lester et al). 2006;19:919-23
Contrast echocardiography versus gated single photon emission computed tomography for the assessment of parameters of left ventricular remodeling after acute myocardial infarction (Lim et al). 2006;19:280-4
Corrected quantification method to determine myocardial blood flow using real-time myocardial contrast echocardiography (Otani et al). 2006;19:973-81
A critical review of patent foramen ovale detection using saline contrast echocardiography: when bubbles lie (Woods and Patel). 2006;19:215-22
Detection of residual infarct-related coronary artery stenosis and multivessel disease after thrombolysis: comparison between myocardial contrast echocardiography and single photon emission computed tomography (Dwivedi et al). 2006;19:546-51
Diagnosis of subacute cardiac rupture by contrast echocardiography (Uno et al). 2006;19:1401.e9-11 (Case rep.)
Effect of time delay on the diagnostic use of contrast echocardiography in patients presenting to the emergency department with chest pain and no S-T segment elevation (Kalvaitis et al). 2006;19:1488-93
Effects of ultrasound contrast agents on doppler tissue velocity estimation (Ressner et al). 2006;19:154-64
Effects of ultrasound contrast during tissue velocity imaging on regional left ventricular velocity, strain, and strain rate measurements (Malm et al). 2006;19:40-7
Feasibility of using a real-time 3-dimensional technique for contrast dobutamine stress echocardiography (Pulerwitz et al). 2006;19:540-5
Heterogeneity of myocardial perfusion in distal coronary embolism with different particle sizes (Akiyama et al). 2006;19:55-63
Impact of the coronary flow reduction at rest on myocardial perfusion and functional indices derived from myocardial contrast and strain echocardiography (Okuda et al). 2006;19:781-7
Myocardial contrast echocardiographic estimates of infarct size predict likelihood of left ventricular remodeling after acute anterior wall myocardial infarction (Main et al). 2006;19:64-70
Myocardial perfusion after percutaneous transluminal septal myocardial ablation as assessed by myocardial contrast echocardiography in patients with hypertrophic obstructive cardiomyopathy (Pedone et al). 2006;19:982-6
Prediction of transmural extent of infarction with contrast echocardiographically derived index of myocardial blood flow and myocardial blood volume fraction: comparison with contrast-enhanced magnetic resonance imaging (Choi et al). 2006;19:1211-9
Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested of known ischemic heart disease (Basic et al). 2006;19:1203-10
Pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia: an echocardiographic perspective (Jassal et al). 2006;19:229.e5-7
Quantification of myocardial perfusion using intravenous myocardial contrast echocardiography in healthy volunteers: comparison with positron emission tomography (Dijkmans et al). 2006;19:285-93
Quantitative adenosine real-time myocardial contrast echocardiography for detection of angiographically significant coronary artery disease (Malm et al). 2006;19:365-72
Real-time simultaneous triplane contrast echocardiography gives rapid, accurate, and reproducible assessment of left ventricular volumes and ejection fraction: a comparison with magnetic resonance imaging (Malm et al). 2006;19:1494-1501
Echocardiography, Doppler
Analysis of mitral annulus motion measurements derived from M-mode, anatomic M-mode, tissue Doppler displacement, and 2-dimensional strain imaging (Hayashi et al). 2006;19:1092-1101
Angle correction for proximal isovelocity surface area method: is it spheric cap or lune? (Sankar et al). 2006;19:241 (Letter)
Assessment of coronary vasodilatation in response to nitroglycerin with transthoracic Doppler echocardiography (Sohn et al). 2006;19:777-80
Assessment of the evolution of normal fetal diastolic function during mid and late gestation by spectral Doppler tissue echocardiography (Nii et al). 2006;19:1431-7
Color tissue Doppler myocardial velocities consistently underestimate spectral tissue Doppler velocities: impact on calculation peak transmitral pulsed Doppler velocity/early diastolic tissue Doppler velocity (E/Ea) (McCulloch et al). 2006;19:744-8
Continuous wave Doppler signal: a mystery (Enseleit et al). 2006;19:1191.e1-3 (Case rep.)
Coronary flow velocity pattern assessed noninvasively by transthoracic color Doppler echocardiography serves as a predictor of adverse cardiac events and left ventricular remodeling in patients with acute myocardial infarction (Katayama et al). 2006;19:335-40
Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11
Doppler tissue imaging in the assessment of atrioventricular conduction time: validation of a novel technique and comparison with electrophysiologic and pulsed wave Doppler-derived equivalents in an animal model (Nii et al). 2006;19:314-21
Early mitral filling/diastolic mitral annular velocity ratio is not a reliable predictor of left ventricular filling pressure in the setting of severe mitral regurgitation (Olson et al). 2006;19:83-7
Echocardiographic indices of increased left ventricular filling pressure and dilation after acute myocardial infarction (Hillis et al). 2006;19:450-6
Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: a doppler echocardiographic study (Luca et al). 2006;19:172-7
Effects of ultrasound contrast agents on doppler tissue velocity estimation (Ressner et al). 2006;19:154-64
Impact of preload and afterload on global and regional right ventricular function and pressure: a quantitative echocardiography study (Kjaergaard et al). 2006;19:515-21
Improved regional myocardial diastolic function assessed by strain rate imaging in patients undergoing percutaneous coronary intervention (Tanaka et al). 2006;19:756-62
Intraventricular pressure gradients in left ventricular aneurysms determined by color M-mode Doppler method: an animal study (Asada-Kamiguchi et al). 2006;19:1112-8
Left ventricular filling hemodynamics in patients with pulmonary edema and preserved versus reduced left ventricular ejection fraction: a prospective Doppler echocardiographic study (Dabbah et al). 2006;19:733-43

Left ventricular systolic longitudinal function: comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals (Mondillo et al). 2006;19:1085-91

Left ventricular Tei index in children: comparison of Doppler tissue imaging, pulsed wave Doppler, and M-mode echocardiography normal values (Cui and Roberson). 2006;19:1438-45

Mitral peak Doppler E-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease (Sharma et al). 2006;19:266-73

Myocardial strain decreases with increasing transmurality of infarction: a Doppler echocardiographic and magnetic resonance correlation study (Sachdev et al). 2006;19:34-9

Noninvasive diagnosis of ischemic and nonischemic cardiomyopathy using coronary flow velocity measurements of the left anterior descending coronary artery by transthoracic Doppler echocardiography (Okura et al). 2006;19:552-8

Noninvasive diagnosis of restenosis by transthoracic doppler echocardiography after percutaneous coronary intervention: comparison with exercise Tl-SPECT (Hirata et al). 2006;19:165-71

A novel echocardiographic Doppler method for estimation of pulmonary arterial pressures (Friedberg et al). 2006;19:559-62

Optimization of pulsed wave tissue doppler to predict left ventricular reverse remodeling after cardiac resynchronization therapy (Jansen et al). 2006;19:185-91

Prediction of subclinical left ventricular dysfunction with strain rate imaging in patients with mild to moderate rheumatic mitral stenosis (Dogan et al). 2006;19:243-8

The prognostic value of pulmonary vascular capacitance determined by Doppler echocardiography in patients with pulmonary arterial hypertension (Mahapatra et al). 2006;19:1045-50

Pulmonary regurgitant flow and detection of dip plateau (Cohen et al). 2006;19:580 (Letter)

Real-time three-dimensional color Doppler echocardiography overcomes the inaccuracies of spectral Doppler for stroke volume calculation (Pemberton et al). 2006;19:1403-10 (Review article)

Reduced myocardial contractility assessed by tissue Doppler echocardiography is associated with increased risk during adrenal surgery of patients with pheochromocytoma: report of a preliminary study (Meune et al). 2006;19:1466-70

Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: study using transthoracic Doppler echocardiography (Youn et al). 2006;19:373-8

Response to: "Echocardiographic assessment of isolated pulmonary valve stenosis: which outpatient Doppler gradient has the most clinically validity?" (DeGroff). 2006;19:364 (Letter)

Single-beat differentiation among left ventricular filling pattern by pulsed wave Doppler echocardiography (Su et al). 2006;19:274-9

"Steal" collaterals: an echocardiographic diagnostic marker for anomalous origin of the left main coronary artery from the pulmonary artery in the adult (Pisacane et al). 2006;19:107.e3-6 (Case rep.)

Usefulness of ventricular longitudinal contractility assessed by doppler tissue imaging in the prediction of reverse remodeling in patients with severe left ventricular systolic dysfunction (Kang et al). 2006;19:178-84

Echocardiography, fetal

Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum (Peterson et al). 2006;19:1393-1400

Echocardiography, intracardiac

The echocardiographic evaluation of intracardiac masses: a review (Peters and Reinhardt). 2006;19:230-40 (Review article)

Intracardiac echocardiography: evolution, recent advances, and current applications (Kort). 2006;19:1192-201 (Review article)

Intracardiac echocardiography for percutaneous mitral valve repair in a swine model (Naqvi et al). 2006;19:147-53

Transjugular biopsy of a right atrial mass under intracardiac echocardiography guidance (Aqel et al). 2006;19:1072.e5-8 (Case rep.)

Echocardiography, intraoperative

Intraoperative assessment of mitral regurgitation: role of phenylephrine challenge (Mihalatos et al). 2006;19:1158-64

Echocardiography, M-mode

Analysis of mitral annulus motion measurements derived from M-mode, anatomic M-mode, tissue Doppler displacement, and 2-dimensional strain imaging (Hayashi et al). 2006;19:1092-1101

Characterization of left ventricular diastolic function in hypertension by use of Doppler tissue imaging and color M-mode techniques (Rovner et al). 2006;19:872-9

A critical review of patent foramen ovale detection using saline contrast echocardiography: when bubbles lie (Woods and Patel). 2006;19:215-22

Intraventricular pressure gradients in left ventricular aneurysms determined by color M-mode Doppler method: an animal study (Asada-Kamiguchi et al). 2006;19:1112-8

Left ventricular systolic longitudinal function: comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals (Mondillo et al). 2006;19:1085-91

Left ventricular Tei index in children: comparison of Doppler tissue imaging, pulsed wave Doppler, and M-mode echocardiography normal values (Cui and Roberson). 2006;19:1438-45

Response to comments on our article "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet" (Kjaergaard et al). 2006;19:1301 (Letter)

Traditional and color M-mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: relations to contractility reserve (Palmieri et al). 2006;19:483-90

Echocardiography, stress

Abnormal longitudinal myocardial functional reserve assessed by exercise tissue Doppler echocardiography in patients with hypertrophic cardiomyopathy (Ha et al). 2006;19:1314-9

Assessment of stress-induced pulmonary interstitial edema by chest ultrasound during exercise echocardiography and its correlation with left ventricular function (Agricola et al). 2006;19:457-63

Color encoding of endocardial motion improves the interpretation of contrast-enhanced echocardiographic stress tests by less-experienced readers (Mor-avi et al). 2006;19:48-54

Comparison of contrast-enhanced real-time live 3-dimensional dobutamine stress echocardiography with contrast 2-dimensional echocardiography for detecting stress-induced wall-motion abnormalities (Takeuchi et al). 2006;19:294-9

Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11

Dobutamine stress echocardiography is highly accurate for the prediction of contractile reserve in the early postoperative period, but may underestimate late recovery in contractile reserve after revascularization of the hibernating myocardium (Zaglavara et al). 2006;19:300-6

Efficacy of ultrasound-assisted stress testing using a hand-

carried ultrasound device for diagnosis of coronary artery disease (Hirano et al). 2006;19:536-9
Exertional changes in circulating cardiac natriuretic peptides in patients with suggested coronary artery disease (Bergeron et al). 2006;19:772-6
Feasibility of using a real-time 3-dimensional technique for contrast dobutamine stress echocardiography (Pulerwitz et al). 2006;19:540-5
Hypotension during dobutamine stress transesophageal echocardiography: relationship with provoked left ventricular obstruction (Siddiqui et al). 2006;19:1144-9
Left ventricular chamber and myocardial systolic function reserve in patients with type 1 diabetes mellitus: insight from traditional and Doppler tissue imaging echocardiography (Palmieri et al). 2006;19:848-56
Noninvasive diagnosis of coronary artery bypass graft disease by dobutamine stress real-time myocardial contrast perfusion imaging (Elhendy et al). 2006;19:1482-7
Prognostic value of mitral regurgitation assessment during exercise echocardiography in patients with known or suspected coronary artery disease (Peteiro et al). 2006;19:1229-37
Response to: "contrast-enhanced real-time 3-dimensional dobutamine stress echocardiography." (Geleijnse et al). 2006;19:1076 (Letter)
Risk stratification by treadmill exercise echocardiography (Peteiro et al). 2006;19:894-901
Role of biplane and biplane echocardiographically guided 3-dimensional echocardiography during dobutamine stress echocardiography (Yang et al). 2006;19:1136-43
Stress echocardiography in octogenarians: transesophageal atrial pacing is accurate, safe, and well tolerated (Kobal et al). 2006;19:1012-6
Traditional and color M-mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: relations to contractility reserve (Palmieri et al). 2006;19:483-90

Echocardiography, three-dimensional

Accuracy and feasibility of online 3-dimensional echocardiography for measurement of left ventricular parameters (Jenkins et al). 2006;19:1119-28
The added value of real-time 3-dimensional echocardiography in the diagnosis of isolated cleft mitral valve in adults (Kuperstein et al). 2006;19:811-4
Asymmetry of systolic anterior motion of the mitral valve in patients with hypertrophic obstructive cardiomyopathy: a real-time three-dimensional echocardiography study (Song et al). 2006;19:1129-35
Can transthoracic live 3-dimensional echocardiography improve the recognition of midventricular obliteration in hypertrophic obstructive cardiomyopathy? (Gregorio et al). 2006;19:1190.e1-4 (Case rep.)
Characterization of atrial septal defect assessed by real-time 3-dimensional echocardiography (van den Bosch et al). 2006;19:815-21
Comparison of contrast-enhanced real-time live 3-dimensional dobutamine stress echocardiography with contrast 2-dimensional echocardiography for detecting stress-induced wall-motion abnormalities (Takeuchi et al). 2006;19:294-9
Evaluation of midwall systolic function in left ventricular hypertrophy: a comparison of 3-dimensional versus 2-dimensional echocardiographic indices (Jung et al). 2006;19:802-10
Feasibility and accuracy of real-time 3-dimensional echocardiographic assessment of ventricular septal defects (van den Bosch et al). 2006;19:7-13
Feasibility of using a real-time 3-dimensional technique for contrast dobutamine stress echocardiography (Pulerwitz et al). 2006;19:540-5
Geometric differences of the mitral valve tenting between anterior and inferior myocardial infarction with significant ischemic mitral regurgitation: quantitation by novel software system with transthoracic real-time three-dimensional echocardiography (Watanabe et al). 2006;19:71-5
Illustration of the additional value of real-time 3-dimensional echocardiography to conventional transthoracic and transesophageal 2-dimensional echocardiography in imaging muscular ventricular septal defects: does this have any impact on individual patient treatment? (Mercer-Rosa et al). 2006;19:1511-9
Quantification of regional left ventricular wall motion from real-time 3-dimensional echocardiography in patients with poor acoustic windows: effects of contrast enhancement tested against cardiac magnetic resonance (Corsi et al). 2006;19:886-93
Quantitative echocardiographic analysis of the right ventricle in healthy individuals (Kjaergaard et al). 2006;19:1365-72
Real-time 3-dimensional echocardiographic evaluation of aortic dissection (Nemes et al). 2006;19:108.e1-3 (Case rep.)
Real-time 3-dimensional echocardiographic imaging of congenital heart disease using matrix-array technology: freehand real-time scanning adds instant morphologic details not well delineated by conventional 2 -dimensional imaging (Seliem et al). 2006;19:121-9
Real-time 3-dimensional echocardiography for prosthetic valve endocarditis: initial experience (Kort). 2006;19:130-9
Real-time three-dimensional color Doppler echocardiography overcomes the inaccuracies of spectral Doppler for stroke volume calculation (Pemberton et al). 2006;19:1403-10 (Review article)
Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease (van den Bosch et al). 2006;19:1-6
Relationship between left atrial and left ventricular function in hypertrophic cardiomyopathy: a real-time 3-dimensional echocardiographic study (Shin et al). 2006;19:796-801
Response to: "contrast-enhanced real-time 3-dimensional dobutamine stress echocardiography." (Geleijnse et al). 2006;19:1076 (Letter)
Role of biplane and biplane echocardiographically guided 3-dimensional echocardiography during dobutamine stress echocardiography (Yang et al). 2006;19:1136-43
Three-dimensional echocardiography improves the understanding of the mechanisms and site of left atrioventricular valve regurgitation in atrioventricular septal defect (Takahashi et al). 2006;19:1502-10
Three-dimensional transesophageal echocardiography in the evaluation of aortic valve destruction by endocarditis (Nemes et al). 2006;19:355.e13-4 (Case rep.)
Three-dimensional transthoracic echocardiography to evaluate cor triatriatum in the adult (Jacobs et al). 2006;19:468.e1-4 (Case rep.)
Three-dimensional tricuspid annular function provides insight into the mechanisms of tricuspid valve regurgitation in classic hypoplastic left heart syndrome (Nii et al). 2006;19:391-402
Use of real-time 3-dimensional transthoracic echocardiography in the evaluation of mitral valve disease (Sugeng et al). 2006;19:413-21
What is the best approach for the assessment of left atrial size? Comparison of various unidimensional and two-dimensional parameters with three-dimensional echocardiographically determined left atrial volume (Maddukuri et al). 2006;19:1026-32
What is the optimal clinical technique for measurement of left ventricular volume after myocardial infarction? A comparative study of 3-dimensional echocardiography, single photon emission computed tomography, and cardiac magnetic resonance imaging (Chan et al). 2006;19:192-201

Echocardiography, transesophageal

Acquired aortic cusp function after chronic left ventricular

assist device support (Banchs et al). 2006;19:1401.e1-3 (Case rep.)
Critical cardiac decompensation during laparoscopic surgery (Popescu and Perrino Jr.). 2006;19:1074.e5-6 (Case rep.)
Dynamic quantitative echocardiographic evaluation of mitral regurgitation in the operating department (Gisbert et al). 2006;19:140-6
Hypotension during dobutamine stress transesophageal echocardiography: relationship with provoked left ventricular obstruction (Siddiqui et al). 2006;19:1144-9
Illustration of the additional value of real-time 3-dimensional echocardiography to conventional transthoracic and transesophageal 2-dimensional echocardiography in imaging muscular ventricular septal defects: does this have any impact on individual patient treatment? (Mercer-Rosa et al). 2006;19:1511-9
In vivo echocardiographic imaging of transplanted human adult stem cells in the myocardium labeled with clinically applicable CliniMACS nanoparticles (Bara et al). 2006;19:563-8
Intracoronary stent visualized on transesophageal echocardiogram in a case of coronary dissection complicated by aortic dissection (Pohlel et al). 2006;19:229.e1-3 (Case rep.)
Intraoperative assessment of mitral regurgitation: role of phenylephrine challenge (Mihalatos et al). 2006;19:1158-64
Perforated periaortic valve abscess presents as acute pericarditis and patient died in transesophageal echocardiographic study (Tsai et al). 2006;19:579.e1-3 (Case rep.)
Primary cardiac lymphoma: diagnosis and treatment guided by transesophageal echocardiogram perfusion imaging (Jang et al). 2006;19:1073.e7-9 (Case rep.)
Primary large B-cell cardiac lymphoma (Premkumar et al). 2006;19:107.e1-2 (Case rep.)
Real-time 3-dimensional echocardiography for prosthetic valve endocarditis: initial experience (Kort). 2006;19:130-9
Single-vessel versus multivessel territory acute ischemic stroke: value of transesophageal echocardiography in the differentiation of embolic stroke (Ulrich et al). 2006;19:1165-9
Stress echocardiography in octogenarians: transesophageal atrial pacing is accurate, safe, and well tolerated (Kobal et al). 2006;19:1012-6
Thoracic aorta evaluation in patients with Takayasu's arteritis by transesophageal echocardiography (Lira-Filho et al). 2006;19:829-34
Three-dimensional transesophageal echocardiography in the evaluation of aortic valve destruction by endocarditis (Nemes et al). 2006;19:355.e13-4 (Case rep.)
Transesophageal echocardiographic acquisition of mitral annulus motion during monitoring of left ventricular function in anesthetized patients (Nilsson et al). 2006;19:499-506
Transesophageal echocardiographic assessment of right atrial appendage anatomy and function: comparison with the left atrial appendage and implications for local thrombus formation (Subramaniam et al). 2006;19:429-33
Transesophageal echocardiographic diagnosis of *Aspergillus fumigatus* aortitis after percutaneous coronary intervention (Khumri et al). 2006;19:1072.e9-11 (Case rep.)
Tricuspid annular velocity in patients undergoing cardiac operation using transesophageal echocardiography (David et al). 2006;19:329-34

Echocardiography, transthoracic

The architecture of a mobile outreach echocardiography service (Gilman et al). 2006;19:1526-8
Assessment of coronary vasodilatation in response to nitroglycerin with transthoracic Doppler echocardiography (Sohn et al). 2006;19:777-80
Atrial and ventricular echocardiographic correlates of the extent of pulmonary embolism in the elderly (Chung et al). 2006;19:347-53
Can transthoracic live 3-dimensional echocardiography improve the recognition of midventricular obliteration in hypertrophic obstructive cardiomyopathy? (Gregorio et al). 2006;19:i190.e1-4 (Case rep.)
Coronary flow velocity pattern assessed noninvasively by transthoracic color Doppler echocardiography serves as a predictor of adverse cardiac events and left ventricular remodeling in patients with acute myocardial infarction (Katayama et al). 2006;19:335-40
Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11
The falciform ligament in the echocardiographic diagnosis of ascites (Cardello et al). 2006;19:1074.e3-4
Feasibility of using a real-time 3-dimensional technique for contrast dobutamine stress echocardiography (Pulerwitz et al). 2006;19:540-5
Geometric differences of the mitral valve tenting between anterior and inferior myocardial infarction with significant ischemic mitral regurgitation: quantitation by novel software system with transthoracic real-time three-dimensional echocardiography (Watanabe et al). 2006;19:71-5
Illustration of the additional value of real-time 3-dimensional echocardiography to conventional transthoracic and transesophageal 2-dimensional echocardiography in imaging muscular ventricular septal defects: does this have any impact on individual patient treatment? (Mercer-Rosa et al). 2006;19:1511-9
Improved visualization of sinus venosus atrial septal defects in adults from the transthoracic approach (Coon and Lang). 2006;19:1072.e1-3 (Case rep.)
A new window of opportunity in echocardiography (Naqvi and Huynh). 2006;19:569-77
Noninvasive diagnosis of ischemic and nonischemic cardiomyopathy using coronary flow velocity measurements of the left anterior descending coronary artery by transthoracic Doppler echocardiography (Okura et al). 2006;19:552-8
Noninvasive diagnosis of restenosis by transthoracic doppler echocardiography after percutaneous coronary intervention: comparison with exercise Tl-SPECT (Hirata et al). 2006;19:165-71
Primary large B-cell cardiac lymphoma (Premkumar et al). 2006;19:107.e1-2 (Case rep.)
Quantitative measures of right ventricular dysfunction by echocardiography in the diagnosis of acute nonmassive pulmonary embolism (Kjaergaard et al). 2006;19:1264-71
A rare complication of atrial septal occluders: diagnosis by transthoracic echocardiography (Dialetto et al). 2006;19:836.e5-8 (Case rep.)
Real-time 3-dimensional echocardiography for prosthetic valve endocarditis: initial experience (Kort). 2006;19:130-9
Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease (van den Bosch et al). 2006;19:1-6
Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: study using transthoracic Doppler echocardiography (Youn et al). 2006;19:373-8
Relationship between the number of coronary risk factors and coronary atherosclerosis assessed by high-frequency transthoracic echocardiography (Takeuchi et al). 2006;19:1056-62
Relationship of mitral annular remodeling to severity of chronic mitral regurgitation (Mihalatos et al). 2006;19:76-82
Sequential transthoracic ultrasound assessment of coronary flow reserve in patient with tako-tsubo syndrome (Citro et al). 2006;19:1402.e5-8 (Case rep.)
Three-dimensional transthoracic echocardiography to evaluate cor triatriatum in the adult (Jacobs et al). 2006;19:468.e1-4 (Case rep.)
Transthoracic coronary flow velocity reserve assessment: com-

parison between adenosine and dobutamine (Meimoun et al). 2006;19:1220-8
Use of real-time 3-dimensional transthoracic echocardiography in the evaluation of mitral valve disease (Sugeng et al). 2006;19:413-21

Echocardiography, two-dimensional
Comparison of contrast-enhanced real-time live 3-dimensional dobutamine stress echocardiography with contrast 2-dimensional echocardiography for detecting stress-induced wall-motion abnormalities (Takeuchi et al). 2006;19:294-9
Evaluation of midwall systolic function in left ventricular hypertrophy: a comparison of 3-dimensional versus 2-dimensional echocardiographic indices (Jung et al). 2006;19:802-10
Hemodynamic significance of mitral stenosis: use of a simple, novel index by 2-dimensional echocardiography (Seow et al). 2006;19:102-6
Illustration of the additional value of real-time 3-dimensional echocardiography to conventional transthoracic and transesophageal 2-dimensional echocardiography in imaging muscular ventricular septal defects: does this have any impact on individual patient treatment? (Mercer-Rosa et al). 2006;19:1511-9
Real-time 3-dimensional echocardiographic imaging of congenital heart disease using matrix-array technology: freehand real-time scanning adds instant morphologic details not well delineated by conventional 2 -dimensional imaging (Seliem et al). 2006;19:121-9
"Steal" collaterals: an echocardiographic diagnostic marker for anomalous origin of the left main coronary artery from the pulmonary artery in the adult (Pisacane et al). 2006;19:107.e3-6 (Case rep.)
Twenty-five-year-old man with a giant left ventricular mass (Moore and Segar). 2006;19:1294.e7-8 (Case rep.)
What is the best approach for the assessment of left atrial size? Comparison of various unidimensional and two-dimensional parameters with three-dimensional echocardiographically determined left atrial volume (Maddukuri et al). 2006;19:1026-32

Echocardiography, utilization
American Society of Echocardiography/ Society of Cardiovascular Anesthesiologists recommendations and guidelines for continuous quality improvements in perioperative echocardiography (Mathew et al). 2006;19:1303-13
Atrial septal defect: anatomoechocardiographic correlation (Muñóz-Castellanos et al). 2006;19:1182-9
Cardiac actinomycosis: an unusual cause of an intracardiac mass (Peters et al). 2006;19:1530.e6 (Case rep.)
A case report of a round cystic tumor in the left ventricular outflow tract (Pérez-Sanz et al). 2006;19:1402.e9-11 (Case rep.)
Detection of heart transplant rejection in adults by echocardiographic diastolic indices: a systematic review of the literature (Mena et al). 2006;19:1295-1300 (Review article)
Double atrial septum with persistent interatrial space: echocardiographic features of a rare atrial septal malformation (Roberson et al). 2006;19:1175-81
Echocardiographic diagnosis and management of bilateral coronary ostial atresia in a patient with pulmonary atresia and intact ventricular septum (Wald et al). 2006;19:939.e1-3 (Case rep.)
Effects of radial left ventricular dyssynchrony on cardiac performance using quantitative tissue Doppler radial strain imaging (Dohi et al). 2006;19:475-82
Glucose-insulin-potassium echocardiography detects improved segmental myocardial function and viable tissue shortly after acute myocardial infarction (Klein et al). 2006;19:763-71
High prevalence of clinically silent severe mitral regurgitation in patients with heart failure: role for echocardiography (Varadarajan et al). 2006;19:1458-61
Improved detection of left ventricular thrombi and spontaneous echocontrast by tissue harmonic imaging in patients with myocardial infarction (Mele et al). 2006;19:1373-81
Left ventricular diverticulum with contractile function in an unusual site (Cay et al). 2006;19:1293.e3-6 (Case rep.)
Left ventricular outflow tract obstruction caused by accessory mitral valve tissue (Sharma et al). 2006;19:354.e5-8 (Case rep.)
New aspects of septal function by using 1-dimensional strain and strain rate imaging (Lindqvist et al). 2006;19:1345-9
Pulmonary artery dissection: a rare complication of pulmonary balloon valvuloplasty diagnosed 11 years after the procedure (Janus et al). 2006;19:1191.e5-8 (Case rep.)
Quantitative measures of right ventricular dysfunction by echocardiography in the diagnosis of acute nonmassive pulmonary embolism (Kjaergaard et al). 2006;19:1264-71
Risk stratification in chronic heart failure: independent and incremental prognostic value of echocardiography and brain natriuretic peptide and its N-terminal fragment (Bruch et al). 2006;19:522-8
Sequence of echocardiographic changes during development of right ventricular failure in rat (Hardziyenka et al). 2006;19:1272-9
Sequential changes of myocardial function during acute myocardial infarction, in the early and chronic phase after coronary intervention described by ultrasonic strain rate imaging (Weidemann et al). 2006;19:839-47
Thrombotic "ghosts": echocardiographic appearance of thrombi with hollow cores and implications regarding mechanism of spontaneous clot lysis (Lavi et al). 2006;19:1530.e5 (Case rep.)
An unusual cause of pulmonary edema: acute rupture of noncoronary sinus of Valsalva aneurysm into the left atrium (Moreo et al). 2006;19:938.e9-11 (Case rep.)
Use of echocardiography in patients with hypertrophic cardiomyopathy: clinical implications of massive hypertrophy (Sorajja et al). 2006;19:788-95
Usefulness of clinical, electrocardiographic, and echocardiographic parameters to detect cardiac asynchrony in patients with left ventricular dysfunction secondary to ischemic or nonischemic heart disease (Perez de Isla et al). 2006;19:1338-44

Ejection fraction; *see* Stroke volume

Elasticity
Impact of type 2 diabetes mellitus on aortic elastic properties in normotensive diabetes: Doppler tissue imaging study (Badran and Elnoaman). 2006;19:1471-81

Elderly; *see* Aged

Electric countershock
Cardioversion in patients with left ventricular thrombus is not associated with increased thromboembolic risk (Bangalore et al). 2006;19:438-40

Electrocardiography
Doppler tissue imaging in the assessment of atrioventricular conduction time: validation of a novel technique and comparison with electrophysiologic and pulsed wave Doppler-derived equivalents in an animal model (Nii et al). 2006;19:314-21
Efficacy of ultrasound-assisted stress testing using a hand-carried ultrasound device for diagnosis of coronary artery disease (Hirano et al). 2006;19:536-9
QRS duration alone misses cardiac dyssynchrony in a substantial proportion of patients with chronic heart failure (Perry et al). 2006;19:1257-63
Usefulness of clinical, electrocardiographic, and echocardiographic parameters to detect cardiac asynchrony in patients with left ventricular dysfunction secondary to ischemic or nonischemic heart disease (Perez de Isla et al). 2006;19:1338-44

Embolism
Heterogeneity of myocardial perfusion in distal coronary em-

bolism with different particle sizes (Akiyama et al). 2006;19:55-63
Left atrial appendage function in patients with systemic embolism in spite of in sinus rhythm (Okamoto et al). 2006;19:211-4
Single-vessel versus multivessel territory acute ischemic stroke: value of transesophageal echocardiography in the differentiation of embolic stroke (Ulrich et al). 2006;19:1165-9

Embolism, paradoxical
Transit thrombus entrapped in patent foramen ovale resolved without clinical embolic events (Can et al). 2006;19:1074.e1-2 (Case rep.)

Emergency medical service
Effect of time delay on the diagnostic use of contrast echocardiography in patients presenting to the emergency department with chest pain and no S-T segment elevation (Kalvaitis et al). 2006;19:1488-93

End stage renal disease; *see* Kidney failure, chronic

Endocarditis
Behçet's aortitis mimicking aortic valve endocarditis with subaortic complications (Shiran et al). 2006;19:578.e1-4 (Case rep.)
Real-time 3-dimensional echocardiography for prosthetic valve endocarditis: initial experience (Kort). 2006;19:130-9

Endocarditis, bacterial
Biventricular mural vegetations in a patient without valvular pathology (Ahmed et al). 2006;19:938.e5-7 (Case rep.)
Three-dimensional transesophageal echocardiography in the evaluation of aortic valve destruction by endocarditis (Nemes et al). 2006;19:355.e13-4 (Case rep.)

Endocardium
Color encoding of endocardial motion improves the interpretation of contrast-enhanced echocardiographic stress tests by less-experienced readers (Mor-avi et al). 2006;19:48-54

Endothelium, vascular
Erectile dysfunction as a generalized vascular dysfunction (Uslu et al). 2006;19:341-6

Erectile dysfunction; *see* Impotence

Estrus synchronization
Asynchrony detection and Doppler tissue imaging (Gaibazzi). 2006;19:111 (Letter)

Exercise
Coronary blood flow after cold exposure and the effect of warm-up exercise (Kim et al). 2006;19:386-90
Myocardial adaptation to short-term high-intensity exercise in highly trained athletes (Neilan et al). 2006;19:1280-5

Exercise test
Abnormal longitudinal myocardial functional reserve assessed by exercise tissue Doppler echocardiography in patients with hypertrophic cardiomyopathy (Ha et al). 2006;19:1314-9
Assessment of stress-induced pulmonary interstitial edema by chest ultrasound during exercise echocardiography and its correlation with left ventricular function (Agricola et al). 2006;19:457-63
Efficacy of ultrasound-assisted stress testing using a hand-carried ultrasound device for diagnosis of coronary artery disease (Hirano et al). 2006;19:536-9
Exertional changes in circulating cardiac natriuretic peptides in patients with suggested coronary artery disease (Bergeron et al). 2006;19:772-6
Isolated ventricular diastolic dysfunction: implications for exercise left ventricular performance in patients without congestive heart failure (Palmieri et al). 2006;19:491-8
Prognostic value of mitral regurgitation assessment during exercise echocardiography in patients with known or suspected coronary artery disease (Peteiro et al). 2006;19:1229-37
Risk stratification by treadmill exercise echocardiography (Peteiro et al). 2006;19:894-901

Extravascular lung water
Ultrasound lung comets: a clinically useful sign of extravascular lung water (Picano et al). 2006;19:356-63 (Review article)

F

Females
Mitral annulus motion as a predictor of mortality in a community-based sample of 75-year-old men and women (Hedberg et al). 2006;19:88-94

Fetal echocardiography; *see* Echocardiography, fetal

Fetus
Assessment of the evolution of normal fetal diastolic function during mid and late gestation by spectral Doppler tissue echocardiography (Nii et al). 2006;19:1431-7
Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum (Peterson et al). 2006;19:1393-1400

Fibrinolysis
Thrombotic "ghosts": echocardiographic appearance of thrombi with hollow cores and implications regarding mechanism of spontaneous clot lysis (Lavi et al). 2006;19:1530.e5 (Case rep.)

Fibrosis
Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: study using transthoracic Doppler echocardiography (Youn et al). 2006;19:373-8

Fistula
Right coronary cameral fistula resulting from surgery of double chamber right ventricle (Leeper et al). 2006;19:1191.e9-11 (Case rep.)
Three major coronary artery-to-left ventricular fistula: an unusual cause of a diastolic murmur at the apex (Roy et al). 2006;19:1402.e1-4 (Case rep.)
Unusual complication of aortic root reconstruction with sparing of the aortic valve: left ventricular outflow tract to right atrial fistula (Ramasubbu et al). 2006;19:469.e5-9 (Case rep.)

Foramen ovale; *see* Heart septum

G

Gestation; *see* Pregnancy

Gestational diabetes; *see* Diabetes, gestational

Glucose
Glucose-insulin-potassium echocardiography detects improved segmental myocardial function and viable tissue shortly after acute myocardial infarction (Klein et al). 2006;19:763-71

Graft rejection
Detection of heart transplant rejection in adults by echocardiographic diastolic indices: a systematic review of the literature (Mena et al). 2006;19:1295-1300 (Review article)

Grawitz tumor; *see* Carcinoma, renal cell

Guidelines
American Society of Echocardiography/ Society of Cardiovascular Anesthesiologists recommendations and guidelines for continuous quality improvements in perioperative echocardiography (Mathew et al). 2006;19:1303-13
Guidelines and standards for performance of a pediatric echocardiogram (Lai et al). 2006;19:1413-30
Guidelines for noninvasive vascular laboratory testing: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Gerhard-Herman et al). 2006;19:955-72 (ASE report)

H

Health personnel
American Society of Echocardiography minimum standards for the cardiac sonographer: a position paper (Bierig et al). 2006;19:471-4

Hear defects, congenital
Echocardiographic diagnosis and management of bilateral cor-

onary ostial atresia in a patient with pulmonary atresia and intact ventricular septum (Wald et al). 2006;19:939.e1-3 (Case rep.)

Heart aneurysm

Intraventricular pressure gradients in left ventricular aneurysms determined by color M-mode Doppler method: an animal study (Asada-Kamiguchi et al). 2006;19:1112-8

Heart-assist devices

Acquired aortic cusp function after chronic left ventricular assist device support (Banchs et al). 2006;19:1401.e1-3 (Case rep.)

The Impella Recover LP 5.0 left ventricular assist device: a bridge to coronary artery bypass grafting and cardiac transplantation (LaRocca et al). 2006;19:468.e5-7 (Case rep.)

Rapid ventricular remodeling with left ventricular unloading postventricular assist device placement: new insights with strain imaging (Havemann et al). 2006;19:355.e9-11 (Case rep.)

Heart atrium

Alterations in left ventricular structure and function in type-1 diabetics: a focus on left atrial contribution to function (Peterson et al). 2006;19:749-55

Double atrial septum with persistent interatrial space: echocardiographic features of a rare atrial septal malformation (Roberson et al). 2006;19:1175-81

Effect of preload reduction by hemodialysis on myocardial ultrasonic characterization, left atrial volume, and Doppler tissue imaging in patients with end-stage renal disease (Fijalkowski et al). 2006;19:1359-64

Extreme redundancy of the valve of the fossa ovalis with right heart hypoplasia in a neonate with trisomy 18 (Vyas and Cabalka). 2006;19:1293.e7-8 (Case rep.)

Isolated left atrial amyloidosis: acute premitral stenosis secondary to spontaneous intramural left atrial hemorrhagic dissection (Edibam et al). 2006;19:938.e1-4 (Case rep.)

Left atrial appendage function in patients with systemic embolism in spite of in sinus rhythm (Okamoto et al). 2006;19:211-4

Stress echocardiography in octogenarians: transesophageal atrial pacing is accurate, safe, and well tolerated (Kobal et al). 2006;19:1012-6

Syncope with a twist (Gosavi et al). 2006;19:1529.e1 (Case rep.)

Transesophageal echocardiographic assessment of right atrial appendage anatomy and function: comparison with the left atrial appendage and implications for local thrombus formation (Subramaniam et al). 2006;19:429-33

Transit thrombus entrapped in patent foramen ovale resolved without clinical embolic events (Can et al). 2006;19:1074.e1-2 (Case rep.)

Transjugular biopsy of a right atrial mass under intracardiac echocardiography guidance (Aqel et al). 2006;19:1072.e5-8 (Case rep.)

An unusual cause of pulmonary edema: acute rupture of noncoronary sinus of Valsalva aneurysm into the left atrium (Moreo et al). 2006;19:938.e9-11 (Case rep.)

Unusual complication of aortic root reconstruction with sparing of the aortic valve: left ventricular outflow tract to right atrial fistula (Ramasubbu et al). 2006;19:469.e5-9 (Case rep.)

What is the best approach for the assessment of left atrial size? Comparison of various unidimensional and two-dimensional parameters with three-dimensional echocardiographically determined left atrial volume (Maddukuri et al). 2006;19:1026-32

Heart catheterization

Response to: "Echocardiographic assessment of isolated pulmonary valve stenosis: which outpatient Doppler gradient has the most clinically validity?" (DeGroff). 2006;19:364 (Letter)

Transcatheter closure of a postinfarction ventricular septal rupture (Martinez et al). 2006;19:1401.e5-e7 (Case rep.)

Use of myocardial performance index in pediatric patients with idiopathic pulmonary arterial hypertension (Dyer et al). 2006;19:21-7

Heart conduction system

Doppler tissue imaging in the assessment of atrioventricular conduction time: validation of a novel technique and comparison with electrophysiologic and pulsed wave Doppler-derived equivalents in an animal model (Nii et al). 2006;19:314-21

Heart defects, congenital

Aorta-to-left ventricle tunnel associated with noncompaction left ventricle (Sadeghpour et al). 2006;19:1073.e1-5 (Case rep.)

Hematoma of a congenitally bicuspid aortic valve in a patient with polycythemia vera and the antiphospholipid antibody syndrome (Webber et al). 2006;19:1530.e4 (Case rep.)

Hypertrabeculation is not distinct from noncompaction in neuromuscular and nonneuromuscular disorders (Stöllberger and Finsterer). 2006;19:109-10 (Letter)

Left ventricular diverticulum with contractile function in an unusual site (Cay et al). 2006;19:1293.e3-6 (Case rep.)

Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease (van den Bosch et al). 2006;19:1-6

Heart diseases

Cardiac actinomycosis: an unusual cause of an intracardiac mass (Peters et al). 2006;19:1530.e6 (Case rep.)

Effects of ultrasound contrast agents on doppler tissue velocity estimation (Ressner et al). 2006;19:154-64

Real-time 3-dimensional echocardiographic imaging of congenital heart disease using matrix-array technology: freehand real-time scanning adds instant morphologic details not well delineated by conventional 2 -dimensional imaging (Seliem et al). 2006;19:121-9

Usefulness of clinical, electrocardiographic, and echocardiographic parameters to detect cardiac asynchrony in patients with left ventricular dysfunction secondary to ischemic or nonischemic heart disease (Perez de Isla et al). 2006;19:1338-44

Heart failure, congestive

Acquired aortic cusp function after chronic left ventricular assist device support (Banchs et al). 2006;19:1401.e1-3 (Case rep.)

Clinical outcomes after cardiac resynchronization therapy: importance of left ventricular diastolic function and origin of heart failure (Waggoner et al). 2006;19:307-13

Comparison of regional versus global assessment of left ventricular function in patients with left ventricular dysfunction, heart failure, or both after myocardial infarction: the Valsartan in Acute Myocardial Infarction Echocardiographic study (Thune et al). 2006;19:1462-5

Critical cardiac decompensation during laparoscopic surgery (Popescu and Perrino Jr.). 2006;19:1074.e5-6 (Case rep.)

The effect of acute hypertension on left ventricular diastolic pressures in a canine model of left ventricular dysfunction with a preserved ejection fraction and elevated left ventricular filling pressures (Lavine and Lavine). 2006;19:1350-8

Heart failure and severe pulmonary hypertension caused by distal detachment of the valve conduit 16 years after the Cabrol composite graft procedure (Tosoratti et al). 2006;19:1190.e5-8 (Case rep.)

High prevalence of clinically silent severe mitral regurgitation in patients with heart failure: role for echocardiography (Varadarajan et al). 2006;19:1458-61

Isolated ventricular diastolic dysfunction: implications for exercise left ventricular performance in patients without congestive heart failure (Palmieri et al). 2006;19:491-8

Left bundle branch block in chronic heart failure-impact on diastolic function, filling pressures, and B-type natriuretic peptide levels (Bruch et al). 2006;19:95-101

QRS duration alone misses cardiac dyssynchrony in a substan-

tial proportion of patients with chronic heart failure (Perry et al). 2006;19:1257-63
Risk stratification in chronic heart failure: independent and incremental prognostic value of echocardiography and brain natriuretic peptide and its N-terminal fragment (Bruch et al). 2006;19:522-8

Heart function tests

Analysis of postsystolic myocardial thickening work in selective myocardial layers during progressive myocardial ischemia (Wang et al). 2006;19:1102-11
Left atrial appendage function in patients with systemic embolism in spite of in sinus rhythm (Okamoto et al). 2006;19:211-4
Preservation of ventricular function in amateur athletes after completion of a marathon (Kean et al). 2006;19:202-5
Real-time 3-dimensional echocardiographic imaging of congenital heart disease using matrix-array technology: freehand real-time scanning adds instant morphologic details not well delineated by conventional 2 -dimensional imaging (Seliem et al). 2006;19:121-9
Response to: "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet." (Voigt). 2006;19:837 (Letter)
Risk assessment in patients with depressed left ventricular function after myocardial infarction using the myocardial performance index-Survival and Ventricular Enlargement (SAVE) Experience (Anavekar et al). 2006;19:28-33
The systolic to diastolic duration ratio in children with heart failure secondary to restrictive cardiomyopathy (Friedberg and Silverman). 2006;19:1326-31
The tei index identifies a differential effect on left and right ventricular function with low-dose anthracycline chemotherapy (Belham et al). 2006;19:206-10
Use of myocardial performance index in pediatric patients with idiopathic pulmonary arterial hypertension (Dyer et al). 2006;19:21-7
Usefulness of ventricular longitudinal contractility assessed by doppler tissue imaging in the prediction of reverse remodeling in patients with severe left ventricular systolic dysfunction (Kang et al). 2006;19:178-84

Heart injuries

Traumatic intramyocardial left ventricular dissection: a case report (Silverstein et al). 2006;19:1529.e2 (Case rep.)

Heart murmurs

Three major coronary artery-to-left ventricular fistula: an unusual cause of a diastolic murmur at the apex (Roy et al). 2006;19:1402.e1-4 (Case rep.)

Heart neoplasms

A case report of a round cystic tumor in the left ventricular outflow tract (Pérez-Sanz et al). 2006;19:1402.e9-11 (Case rep.)
The echocardiographic evaluation of intracardiac masses: a review (Peters and Reinhardt). 2006;19:230-40 (Review article)
Floating heart metastasis (Mumoli et al). 2006;19:1293.e1-2 (Case rep.)
Primary cardiac lymphoma: diagnosis and treatment guided by transesophageal echocardiogram perfusion imaging (Jang et al). 2006;19:1073.e7-9 (Case rep.)
Primary large B-cell cardiac lymphoma (Premkumar et al). 2006;19:107.e1-2 (Case rep.)
Smooth muscle neoplasm presenting as intrapericardial myxoma (Horton et al). 2006;19:835.e1-3 (Case rep.)
Transjugular biopsy of a right atrial mass under intracardiac echocardiography guidance (Aqel et al). 2006;19:1072.e5-8 (Case rep.)

Heart rate

Doppler tissue imaging in the assessment of atrioventricular conduction time: validation of a novel technique and comparison with electrophysiologic and pulsed wave Doppler-derived equivalents in an animal model (Nii et al). 2006;19:314-21
Effect of age and heart rate on atrial mechanical function assessed by Doppler tissue imaging in healthy individuals (Zhang et al). 2006;19:422-8
Single-beat differentiation among left ventricular filling pattern by pulsed wave Doppler echocardiography (Su et al). 2006;19:274-9

Heart rupture

Diagnosis of subacute cardiac rupture by contrast echocardiography (Uno et al). 2006;19:1401.e9-11 (Case rep.)
Papillary muscle rupture after blunt chest trauma (Cordovil et al). 2006;19:469.e1-3 (Case rep.)

Heart rupture, traumatic; *see* Heart injuries

Heart septal defects

The impact of patch augmentation on left atrioventricular valve dynamics in patients with atrioventricular septal defects: early and midterm follow-up (Roman et al). 2006;19:1382-92
Three-dimensional echocardiography improves the understanding of the mechanisms and site of left atrioventricular valve regurgitation in atrioventricular septal defect (Takahashi et al). 2006;19:1502-10

Heart septal defects, atrial

Atrial septal defect: anatomoechocardiographic correlation (Muñóz-Castellanos et al). 2006;19:1182-9
Characterization of atrial septal defect assessed by real-time 3-dimensional echocardiography (van den Bosch et al). 2006;19:815-21
Double atrial septum with persistent interatrial space: echocardiographic features of a rare atrial septal malformation (Roberson et al). 2006;19:1175-81
Improved visualization of sinus venosus atrial septal defects in adults from the transthoracic approach (Coon and Lang). 2006;19:1072.e1-3 (Case rep.)
A rare complication of atrial septal occluders: diagnosis by transthoracic echocardiography (Dialetto et al). 2006;19:836.e5-8 (Case rep.)
Subtotal anomalous pulmonary venous connection and left cor triatriatum: a rare diagnosis in adulthood (Caputo et al). 2006;19:836.e1-4 (Case rep.)
Three-dimensional echocardiography improves the understanding of the mechanisms and site of left atrioventricular valve regurgitation in atrioventricular septal defect (Takahashi et al). 2006;19:1502-10
Transit thrombus entrapped in patent foramen ovale resolved without clinical embolic events (Can et al). 2006;19:1074.e1-2 (Case rep.)
Ultrasonic strain rate and strain imaging of the right ventricle in children before and after percutaneous closure of an atrial septal defect (Eyskens et al). 2006;19:994-1000

Heart septal defects, ventricular

Feasibility and accuracy of real-time 3-dimensional echocardiographic assessment of ventricular septal defects (van den Bosch et al). 2006;19:7-13
Illustration of the additional value of real-time 3-dimensional echocardiography to conventional transthoracic and transesophageal 2-dimensional echocardiography in imaging muscular ventricular septal defects: does this have any impact on individual patient treatment? (Mercer-Rosa et al). 2006;19:1511-9
Transcatheter closure of a postinfarction ventricular septal rupture (Martinez et al). 2006;19:1401.e5-7 (Case rep.)

Heart septum

Assessment of diastolic function with Doppler tissue imaging after cardiac surgery: effect of "postoperative septum" in on-pump and off-pump procedures (Malouf et al). 2006;19:464-7
A critical review of patent foramen ovale detection using saline contrast echocardiography: when bubbles lie (Woods and Patel). 2006;19:215-22
Echocardiographic diagnosis and management of bilateral cor-

onary ostial atresia in a patient with pulmonary atresia and intact ventricular septum (Wald et al). 2006;19:939.e1-3 (Case rep.)
Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum (Peterson et al). 2006;19:1393-1400
Extreme redundancy of the valve of the fossa ovalis with right heart hypoplasia in a neonate with trisomy 18 (Vyas and Cabalka). 2006;19:1293.e7-8 (Case rep.)
New aspects of septal function by using 1-dimensional strain and strain rate imaging (Lindqvist et al). 2006;19:1345-9

Heart surgery
Assessment of diastolic function with Doppler tissue imaging after cardiac surgery: effect of "postoperative septum" in on-pump and off-pump procedures (Malouf et al). 2006;19:464-7
Heart failure and severe pulmonary hypertension caused by distal detachment of the valve conduit 16 years after the Cabrol composite graft procedure (Tosoratti et al). 2006;19:1190.e5-8 (Case rep.)
The impact of patch augmentation on left atrioventricular valve dynamics in patients with atrioventricular septal defects: early and midterm follow-up (Roman et al). 2006;19:1382-92
Right coronary cameral fistula resulting from surgery of double chamber right ventricle (Leeper et al). 2006;19:1191.e9-11 (Case rep.)
Tricuspid annular velocity in patients undergoing cardiac operation using transesophageal echocardiography (David et al). 2006;19:329-34
Ultrasonic strain rate and strain imaging of the right ventricle in children before and after percutaneous closure of an atrial septal defect (Eyskens et al). 2006;19:994-1000

Heart transplantation
Detection of heart transplant rejection in adults by echocardiographic diastolic indices: a systematic review of the literature (Mena et al). 2006;19:1295-1300
The Impella Recover LP 5.0 left ventricular assist device: a bridge to coronary artery bypass grafting and cardiac transplantation (LaRocca et al). 2006;19:468.e5-7 (Case rep.)

Heart valve prosthesis
Acute improvement in myocardial function assessed by myocardial strain and strain rate after aortic valve replacement for aortic stenosis (Iwahashi et al). 2006;19:1238-44
A rare complication of atrial septal occluders: diagnosis by transthoracic echocardiography (Dialetto et al). 2006;19:836.e5-8 (Case rep.)
Real-time 3-dimensional echocardiography for prosthetic valve endocarditis: initial experience (Kort). 2006;19:130-9

Heart valves
Heart failure and severe pulmonary hypertension caused by distal detachment of the valve conduit 16 years after the Cabrol composite graft procedure (Tosoratti et al). 2006;19:1190.e5-8 (Case rep.)

Heart ventricle
Age-related changes in left ventricular twist assessed by two-dimensional speckle-tracking imaging (Takeuchi et al). 2006;19:1077-84
Alterations in left ventricular structure and function in type-1 diabetics: a focus on left atrial contribution to function (Peterson et al). 2006;19:749-55
Aorta-to-left ventricle tunnel associated with noncompaction left ventricle (Sadeghpour et al). 2006;19:1073.e1-5 (Case rep.)
Biventricular mural vegetations in a patient without valvular pathology (Ahmed et al). 2006;19:938.e5-7 (Case rep.)
Cardioversion in patients with left ventricular thrombus is not associated with increased thromboembolic risk (Bangalore et al). 2006;19:438-40
A case report of a round cystic tumor in the left ventricular outflow tract (Pérez-Sanz et al). 2006;19:1402.e9-11 (Case rep.)
Diagnosis of subacute cardiac rupture by contrast echocardiography (Uno et al). 2006;19:1401.e9-11 (Case rep.)
Echocardiographic diagnosis and management of bilateral coronary ostial atresia in a patient with pulmonary atresia and intact ventricular septum (Wald et al). 2006;19:939.e1-3 (Case rep.)
Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum (Peterson et al). 2006;19:1393-1400
Improved detection of left ventricular thrombi and spontaneous echocontrast by tissue harmonic imaging in patients with myocardial infarction (Mele et al). 2006;19:1373-81
Intraventricular pressure gradients in left ventricular aneurysms determined by color M-mode Doppler method: an animal study (Asada-Kamiguchi et al). 2006;19:1112-8
Left ventricular diverticulum with contractile function in an unusual site (Cay et al). 2006;19:1293.e3-6 (Case rep.)
Midventricular hypokinesis as a cardiac manifestation of anaphylaxis: a case report (Han and Yeon). 2006;19:1529.e3 (Case rep.)
Preservation of ventricular function in amateur athletes after completion of a marathon (Kean et al). 2006;19:202-5
Pseudodyskinesis–a novel mechanism for abnormal motion of the heart's diaphragmatic wall (Vennalaganti et al). 2006;19:1294.e5-6 (Case rep.)
Quantitative echocardiographic analysis of the right ventricle in healthy individuals (Kjaergaard et al). 2006;19:1365-72
Rapid-growing left intraventricular cardiac hemangioma (Di Valentino et al). 2006;19:939.e5-7 (Case rep.)
Right coronary cameral fistula resulting from surgery of double chamber right ventricle (Leeper et al). 2006;19:1191.e9-11 (Case rep.)
Right ventricular function in asymptomatic individuals with a systemic right ventricle (Bos et al). 2006;19:1033-7
Right ventricular metastasis caused by a renal cell carcinoma (Grawitz tumor): case report (Eddicks et al). 2006;19:1073.e11-5 (Case rep.)
Right ventricular metastasis of uterine squamous cell carcinoma (Nesser et al). 2006;19:1294.e1-3 (Case rep.)
The tei index identifies a differential effect on left and right ventricular function with low-dose anthracycline chemotherapy (Belham et al). 2006;19:206-10
Three major coronary artery-to-left ventricular fistula: an unusual cause of a diastolic murmur at the apex (Roy et al). 2006;19:1402.e1-4 (Case rep.)
Traumatic intramyocardial left ventricular dissection: a case report (Silverstein et al). 2006;19:1529.e2 (Case rep.)
Twenty-five-year-old man with a giant left ventricular mass (Moore and Segar). 2006;19:1294.e7-8 (Case rep.)
Unusual complication of aortic root reconstruction with sparing of the aortic valve: left ventricular outflow tract to right atrial fistula (Ramasubbu et al). 2006;19:469.e5-9 (Case rep.)
What is the optimal clinical technique for measurement of left ventricular volume after myocardial infarction? A comparative study of 3-dimensional echocardiography, single photon emission computed tomography, and cardiac magnetic resonance imaging (Chan et al). 2006;19:192-201

Hemangioma, cavernous
A case report of a round cystic tumor in the left ventricular outflow tract (Pérez-Sanz et al). 2006;19:1402.e9-11 (Case rep.)
Rapid-growing left intraventricular cardiac hemangioma (Di Valentino et al). 2006;19:939.e5-7 (Case rep.)

Hematoma
Hematoma of a congenitally bicuspid aortic valve in a patient with polycythemia vera and the antiphospholipid antibody syndrome (Webber et al). 2006;19:1530.e1-3 (Case rep.)
Traumatic intramyocardial left ventricular dissection: a case report (Silverstein et al). 2006;19:1529.e5-8 (Case rep.)

Hemodialyis
Effect of preload reduction by hemodialysis on myocardial

ultrasonic characterization, left atrial volume, and Doppler tissue imaging in patients with end-stage renal disease (Fijalkowski et al). 2006;19:1359-64

Hemodynamics

Hemodynamic change in transmitral gradient during Valsalva maneuver and its relation with left atrial function, functional capacity, and pulmonary artery pressure in patients with mitral stenosis in sinus rhythm (Ozeke et al). 2006;19:1245-50

Hemodynamic significance of mitral stenosis: use of a simple, novel index by 2-dimensional echocardiography (Seow et al). 2006;19:102-6

Left ventricular filling hemodynamics in patients with pulmonary edema and preserved versus reduced left ventricular ejection fraction: a prospective Doppler echocardiographic study (Dabbah et al). 2006;19:733-43

Preoperative pulmonary hypertension is associated with postoperative left ventricular dysfunction in chronic organic mitral regurgitation: an echocardiographic and hemodynamic study (Yang et al). 2006;19:1051-5

Pulmonary hemodynamic effects of dipyridamole infusion in patients with normal and elevated pulmonary artery systolic pressure receiving PB127 (Main et al). 2006;19:1038-44

Hemorrhage

Isolated left atrial amyloidosis: acute premitral stenosis secondary to spontaneous intramural left atrial hemorrhagic dissection (Edibam et al). 2006;19:938.e1-4 (Case rep.)

Spontaneous retraction of an intramyocardial dissecting hemorrhage and multiple left ventricular thrombus formation in subacute myocardial infarction and antiphospholipid syndrome: a case report with long-term follow-up (Bahlmann et al). 2006;19:578.e5-8 (Case rep.)

Hereditary hemorrhagic telagiectasia; *see* Telangiectasia, hereditary hemorrhagic

Hypertension

Characterization of left ventricular diastolic function in hypertension by use of Doppler tissue imaging and color M-mode techniques (Rovner et al). 2006;19:872-9

The effect of acute hypertension on left ventricular diastolic pressures in a canine model of left ventricular dysfunction with a preserved ejection fraction and elevated left ventricular filling pressures (Lavine and Lavine). 2006;19:1350-8

Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: study using transthoracic Doppler echocardiography (Youn et al). 2006;19:373-8

Hypertension, pulmonary

Heart failure and severe pulmonary hypertension caused by distal detachment of the valve conduit 16 years after the Cabrol composite graft procedure (Tosoratti et al). 2006;19:1190.e5-8 (Case rep.)

Noninvasive Doppler tissue measurement of pulmonary artery compliance in children with pulmonary hypertension (Dyer et al). 2006;19:403-12

Preoperative pulmonary hypertension is associated with postoperative left ventricular dysfunction in chronic organic mitral regurgitation: an echocardiographic and hemodynamic study (Yang et al). 2006;19:1051-5

The prognostic value of pulmonary vascular capacitance determined by Doppler echocardiography in patients with pulmonary arterial hypertension (Mahapatra et al). 2006;19:1045-50

Pulmonary embolism and right heart function: insights from myocardial Doppler tissue imaging (Hsiao et al). 2006;19:822-8

Use of myocardial performance index in pediatric patients with idiopathic pulmonary arterial hypertension (Dyer et al). 2006;19:21-7

Hypertrabeculation; *see* Heart defects, congenital

Hypertrophic cardiomyopathy; *see* Cardiomyopathy, hypertrophic

Hypertrophic obstructive cardiomyopathy; *see* Cardiomyopathy, hypertrophic

Hypertrophy, left ventricular

Evaluation of midwall systolic function in left ventricular hypertrophy: a comparison of 3-dimensional versus 2-dimensional echocardiographic indices (Jung et al). 2006;19:802-10

Hypokinesia

Midventricular hypokinesis as a cardiac manifestation of anaphylaxis: a case report (Han and Yeon). 2006;19:1529.e9-11 (Case rep.)

Hypoplasia

Extreme redundancy of the valve of the fossa ovalis with right heart hypoplasia in a neonate with trisomy 18 (Vyas and Cabalka). 2006;19:1293.e7-8 (Case rep.)

Hypoplastic left heart syndrome

Three-dimensional tricuspid annular function provides insight into the mechanisms of tricuspid valve regurgitation in classic hypoplastic left heart syndrome (Nii et al). 2006;19:391-402

Hypotension

Hypotension during dobutamine stress transesophageal echocardiography: relationship with provoked left ventricular obstruction (Siddiqui et al). 2006;19:1144-9

I

Image enhancement

Contrast echocardiography: experience in a clinical echocardiography laboratory (Lester et al). 2006;19:919-23

Improved visualization of sinus venosus atrial septal defects in adults from the transthoracic approach (Coon and Lang). 2006;19:1072.e1-3 (Case rep.)

Intracardiac echocardiography: evolution, recent advances, and current applications (Kort). 2006;19:1192-201 (Review article)

Long-term survival of a patient with isolated noncompaction of the ventricular myocardium (Salemi et al). 2006;19:354.e1-3 (Case rep.)

Impotence

Erectile dysfunction as a generalized vascular dysfunction (Uslu et al). 2006;19:341-6

Infant, newborn

Extreme redundancy of the valve of the fossa ovalis with right heart hypoplasia in a neonate with trisomy 18 (Vyas and Cabalka). 2006;19:1293.e7-8 (Case rep.)

Infarction, myocardial; *see* Myocardial infarction

Innominate artery; *see* Brachiocephalic trunk

Insufflation

Critical cardiac decompensation during laparoscopic surgery (Popescu and Perrino Jr.). 2006;19:1074.e5-6 (Case rep.)

Insulin

Glucose-insulin-potassium echocardiography detects improved segmental myocardial function and viable tissue shortly after acute myocardial infarction (Klein et al). 2006;19:763-71

Internet

Accuracy and feasibility of online 3-dimensional echocardiography for measurement of left ventricular parameters (Jenkins et al). 2006;19:1119-28

Intracardiac echocardiography; *see* Echocardiography, intracardiac

Intraoperative echocardiography; *see* Echocardiography, intraoperative

Ischemia

Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested of known ischemic heart disease (Basic et al). 2006;19:1203-10

Single-vessel versus multivessel territory acute ischemic stroke: value of transesophageal echocardiography in the differentiation of embolic stroke (Ulrich et al). 2006;19:1165-9

Usefulness of clinical, electrocardiographic, and echocardiographic parameters to detect cardiac asynchrony in patients

with left ventricular dysfunction secondary to ischemic or nonischemic heart disease (Perez de Isla et al). 2006;19:1338-44

Ischemic stroke; *see* Cerebrovascular disorders

K

Kidney failure, chronic
Effect of preload reduction by hemodialysis on myocardial ultrasonic characterization, left atrial volume, and Doppler tissue imaging in patients with end-stage renal disease (Fijalkowski et al). 2006;19:1359-64
Mitral peak Doppler E-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease (Sharma et al). 2006;19:266-73

L

Laparoscopy
Critical cardiac decompensation during laparoscopic surgery (Popescu and Perrino Jr.). 2006;19:1074.e5-6 (Case rep.)

Left atrial function; *see* Atrial function, left

Left ventricular assist device; *see* Heart-assist devices

Left ventricular dysfunction; *see* Ventricular dysfunction, left

Left ventricular function; *see* Ventricular function, left

Left ventricular hypertrophy; *see* Hypertrophy, left ventricular

Left ventricular remodeling; *see* Ventricular remodeling

Letters to the editor
Angle correction for proximal isovelocity surface area method: is it spheric cap or lune? (Sankar et al). 2006;19:241 (Letter)
Asynchrony detection and Doppler tissue imaging (Gaibazzi). 2006;19:111 (Letter)
Haunted liver (Vidovich and Smulevitz). 2006;19:1302 (Letter)
Hypertrabeculation is not distinct from noncompaction in neuromuscular and nonneuromuscular disorders (Stöllberger and Finsterer). 2006;19:109-10 (Letter)
Letters to the editor (Ogunyanklu) (Letter); (McCulloch and Dokainish) (Reply). 2006;19:1411-2
Pulmonary regurgitant flow and detection of dip plateau (Cohen et al). 2006;19:580 (Letter)
Reply to "Angle correction for proximal isovelocity surface area method: is it spheric cap or lune? (Genc et al). 2006;19:940 (Letter)
Response to: "contrast-enhanced real-time 3-dimensional dobutamine stress echocardiography." (Geleijnse et al). 2006;19:1076 (Letter)
Response to: "Echocardiographic assessment of isolated pulmonary valve stenosis: which outpatient Doppler gradient has the most clinically validity?" (DeGroff). 2006;19:364 (Letter)
Response to: "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet." (Voigt). 2006;19:837 (Letter)
Response to comments on our article "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet" (Kjaergaard et al). 2006;19:1301 (Letter)

Levosimendan; *see* Simendan

Ligaments
The falciform ligament in the echocardiographic diagnosis of ascites (Cardello et al). 2006;19:1074.e3-4

Liver
Haunted liver (Vidovich and Smulevitz). 2006;19:1302 (Letter)

Lung
Ultrasound lung comets: a clinically useful sign of extravascular lung water (Picano et al). 2006;19:356-63 (Review article)

Lupus erythematosus, systemic
Right heart function in systemic lupus erythematosus: insights from myocardial Doppler tissue imaging (Gin et al). 2006;19:441-9

Lymphoma
Primary cardiac lymphoma: diagnosis and treatment guided by transesophageal echocardiogram perfusion imaging (Jang et al). 2006;19:1073.e7-9 (Case rep.)

Lymphoma, B-cell
Primary large B-cell cardiac lymphoma (Premkumar et al). 2006;19:107.e1-2 (Case rep.)

M

M-mode echocardiography; *see* Echocardiography, M-mode

Magnetic resonance imaging
Myocardial strain decreases with increasing transmurality of infarction: a Doppler echocardiographic and magnetic resonance correlation study (Sachdev et al). 2006;19:34-9
Prediction of transmural extent of infarction with contrast echocardiographically derived index of myocardial blood flow and myocardial blood volume fraction: comparison with contrast-enhanced magnetic resonance imaging (Choi et al). 2006;19:1211-9
Quantification of regional left ventricular wall motion from real-time 3-dimensional echocardiography in patients with poor acoustic windows: effects of contrast enhancement tested against cardiac magnetic resonance (Corsi et al). 2006;19:886-93
Real-time simultaneous triplane contrast echocardiography gives rapid, accurate, and reproducible assessment of left ventricular volumes and ejection fraction: a comparison with magnetic resonance imaging (Malm et al). 2006;19:1494-1501
Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease (van den Bosch et al). 2006;19:1-6
Right ventricular metastasis caused by a renal cell carcinoma (Grawitz tumor): case report (Eddicks et al). 2006;19:1073.e11-5 (Case rep.)
What is the optimal clinical technique for measurement of left ventricular volume after myocardial infarction? A comparative study of 3-dimensional echocardiography, single photon emission computed tomography, and cardiac magnetic resonance imaging (Chan et al). 2006;19:192-201

Metastasis; *see* Neoplasm metastasis

Microcirculation
Myocardial perfusion after percutaneous transluminal septal myocardial ablation as assessed by myocardial contrast echocardiography in patients with hypertrophic obstructive cardiomyopathy (Pedone et al). 2006;19:982-6

Mitral regurgitation; *see* Mitral valve insufficiency

Mitral valve
Analysis of mitral annulus motion measurements derived from M-mode, anatomic M-mode, tissue Doppler displacement, and 2-dimensional strain imaging (Hayashi et al). 2006;19:1092-1101
Angle correction for proximal isovelocity surface area method: is it spheric cap or lune? (Sankar et al). 2006;19:241 (Letter)
Asymmetry of systolic anterior motion of the mitral valve in patients with hypertrophic obstructive cardiomyopathy: a real-time three-dimensional echocardiography study (Song et al). 2006;19:1129-35
The impact of patch augmentation on left atrioventricular valve dynamics in patients with atrioventricular septal defects: early and midterm follow-up (Roman et al). 2006;19:1382-92
Intracardiac echocardiography for percutaneous mitral valve repair in a swine model (Naqvi et al). 2006;19:147-53
Left ventricular outflow tract obstruction caused by accessory mitral valve tissue (Sharma et al). 2006;19:354.e5-8 (Case rep.)
Left ventricular systolic longitudinal function: comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals (Mondillo et al). 2006;19:1085-91
Mitral annulus motion as a predictor of mortality in a community-based sample of 75-year-old men and women (Hedberg et al). 2006;19:88-94

Mitral peak Doppler E-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease (Sharma et al). 2006;19:266-73
A rare complication of atrial septal occluders: diagnosis by transthoracic echocardiography (Dialetto et al). 2006;19:836.e5-8 (Case rep.)
Response to: "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet." (Voigt). 2006;19:837 (Letter)
Response to comments on our article "Measurement of cardiac time intervals by Doppler tissue M-mode imaging of the anterior mitral leaflet" (Kjaergaard et al). 2006;19:1301 (Letter)
Transesophageal echocardiographic acquisition of mitral annulus motion during monitoring of left ventricular function in anesthetized patients (Nilsson et al). 2006;19:499-506
Use of real-time 3-dimensional transthoracic echocardiography in the evaluation of mitral valve disease (Sugeng et al). 2006;19:413-21

Mitral valve insufficiency
The added value of real-time 3-dimensional echocardiography in the diagnosis of isolated cleft mitral valve in adults (Kuperstein et al). 2006;19:811-4
Dynamic quantitative echocardiographic evaluation of mitral regurgitation in the operating department (Gisbert et al). 2006;19:140-6
Early mitral filling/diastolic mitral annular velocity ratio is not a reliable predictor of left ventricular filling pressure in the setting of severe mitral regurgitation (Olson et al). 2006;19:83-7
Geometric differences of the mitral valve tenting between anterior and inferior myocardial infarction with significant ischemic mitral regurgitation: quantitation by novel software system with transthoracic real-time three-dimensional echocardiography (Watanabe et al). 2006;19:71-5
High prevalence of clinically silent severe mitral regurgitation in patients with heart failure: role for echocardiography (Varadarajan et al). 2006;19:1458-61
Improved papillary muscle function attenuates functional mitral regurgitation in patients with dilated cardiomyopathy after cardiac resynchronization therapy (Karvounis et al). 2006;19:1150-7
Intraoperative assessment of mitral regurgitation: role of phenylephrine challenge (Mihalatos et al). 2006;19:1158-64
Preoperative pulmonary hypertension is associated with postoperative left ventricular dysfunction in chronic organic mitral regurgitation: an echocardiographic and hemodynamic study (Yang et al). 2006;19:1051-5
Prognostic value of mitral regurgitation assessment during exercise echocardiography in patients with known or suspected coronary artery disease (Peteiro et al). 2006;19:1229-37
Relationship of mitral annular remodeling to severity of chronic mitral regurgitation (Mihalatos et al). 2006;19:76-82
Three-dimensional echocardiography improves the understanding of the mechanisms and site of left atrioventricular valve regurgitation in atrioventricular septal defect (Takahashi et al). 2006;19:1502-10

Mitral valve stenosis
Effect of percutaneous mitral balloon valvuloplasty on left atrial appendage function: a Doppler tissue study (Karakaya et al). 2006;19:434-7
Hemodynamic change in transmitral gradient during Valsalva maneuver and its relation with left atrial function, functional capacity, and pulmonary artery pressure in patients with mitral stenosis in sinus rhythm (Ozeke et al). 2006;19:1245-50
Hemodynamic significance of mitral stenosis: use of a simple, novel index by 2-dimensional echocardiography (Seow et al). 2006;19:102-6
Increased prevalence of mitral stenosis in women (Movahed et al). 2006;19:911-3
Isolated left atrial amyloidosis: acute premitral stenosis secondary to spontaneous intramural left atrial hemorrhagic dissection (Edibam et al). 2006;19:938.e1-4 (Case rep.)
Prediction of subclinical left ventricular dysfunction with strain rate imaging in patients with mild to moderate rheumatic mitral stenosis (Dogan et al). 2006;19:243-8

Monocrotaline
Sequence of echocardiographic changes during development of right ventricular failure in rat (Hardziyenka et al). 2006;19:1272-9

Mortality
Mitral annulus motion as a predictor of mortality in a community-based sample of 75-year-old men and women (Hedberg et al). 2006;19:88-94
Mitral peak Doppler E-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease (Sharma et al). 2006;19:266-73

Movement
Effect of aging on twist-displacement loop by 2-dimensional speckle tracking imaging (Nakai et al). 2006;19:880-5
Longitudinal isovolumic displacement of the left ventricular myocardium assessed by tissue velocity echocardiography in healthy individuals (Lind et al). 2006;19:255-65

Movement disorders
Pseudodyskinesis–a novel mechanism for abnormal motion of the heart's diaphragmatic wall (Vennalaganti et al). 2006;19:1294.e5-6 (Case rep.)

Muscle, smooth, vascular
Smooth muscle neoplasm presenting as intrapericardial myxoma (Horton et al). 2006;19:835.e1-3 (Case rep.)

Muscular dystrophy, Duchenne
Abnormalities of diastolic function precede dilated cardiomyopathy associated with Duchenne muscular dystrophy (Markham et al). 2006;19:865-71

Myocardial contraction
Longitudinal isovolumic displacement of the left ventricular myocardium assessed by tissue velocity echocardiography in healthy individuals (Lind et al). 2006;19:255-65
Reduced myocardial contractility assessed by tissue Doppler echocardiography is associated with increased risk during adrenal surgery of patients with pheochromocytoma: report of a preliminary study (Meune et al). 2006;19:1466-70
The role of left ventricular long-axis contraction in patients with asymptomatic aortic regurgitation (Paraskevaidis et al). 2006;19:249-54
Usefulness of ventricular longitudinal contractility assessed by doppler tissue imaging in the prediction of reverse remodeling in patients with severe left ventricular systolic dysfunction (Kang et al). 2006;19:178-84

Myocardial diseases
Myocardial lesion formation using high-intensity focused ultrasound (Engel et al). 2006;19:932-7
Sequential transthoracic ultrasound assessment of coronary flow reserve in patient with tako-tsubo syndrome (Citro et al). 2006;19:1402.e5-8 (Case rep.)
Takotsubo cardiomyopathy: a case report (Mitchell and Crone). 2006;19:1190.e9-10 (Case rep.)

Myocardial function; *see* Heart function tests

Myocardial hypertrophy; *see* Cardiomyopathy, hypertrophy

Myocardial infarction
Asynchrony of left ventricular systolic performance after the first acute myocardial infarction in patients with narrow QRS complexes: Doppler tissue imaging study (Elnoamany et al). 2006;19:1449-57
Automated quantification of the spatial extent of perfusion defects and viability on myocardial contrast echocardiography (Micari et al). 2006;19:379-85
Comparison of regional versus global assessment of left ven-

tricular function in patients with left ventricular dysfunction, heart failure, or both after myocardial infarction: the Valsartan in Acute Myocardial Infarction Echocardiographic study (Thune et al). 2006;19:1462-5
Contrast echocardiography versus gated single photon emission computed tomography for the assessment of parameters of left ventricular remodeling after acute myocardial infarction (Lim et al). 2006;19:280-4
Coronary flow velocity pattern assessed noninvasively by transthoracic color Doppler echocardiography serves as a predictor of adverse cardiac events and left ventricular remodeling in patients with acute myocardial infarction (Katayama et al). 2006;19:335-40
Echocardiographic indices of increased left ventricular filling pressure and dilation after acute myocardial infarction (Hillis et al). 2006;19:450-6
Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: a doppler echocardiographic study (Luca et al). 2006;19:172-7
Geometric differences of the mitral valve tenting between anterior and inferior myocardial infarction with significant ischemic mitral regurgitation: quantitation by novel software system with transthoracic real-time three-dimensional echocardiography (Watanabe et al). 2006;19:71-5
Glucose-insulin-potassium echocardiography detects improved segmental myocardial function and viable tissue shortly after acute myocardial infarction (Klein et al). 2006;19:763-71
Improved detection of left ventricular thrombi and spontaneous echocontrast by tissue harmonic imaging in patients with myocardial infarction (Mele et al). 2006;19:1373-81
Intracoronary stent visualized on transesophageal echocardiogram in a case of coronary dissection complicated by aortic dissection (Pohlel et al). 2006;19:229.e1-3 (Case rep.)
Myocardial contrast echocardiographic estimates of infarct size predict likelihood of left ventricular remodeling after acute anterior wall myocardial infarction (Main et al). 2006;19:64-70
Myocardial strain decreases with increasing transmurality of infarction: a Doppler echocardiographic and magnetic resonance correlation study (Sachdev et al). 2006;19:34-9
Prediction of transmural extent of infarction with contrast echocardiographically derived index of myocardial blood flow and myocardial blood volume fraction: comparison with contrast-enhanced magnetic resonance imaging (Choi et al). 2006;19:1211-9
Risk assessment in patients with depressed left ventricular function after myocardial infarction using the myocardial performance index-Survival and Ventricular Enlargement (SAVE) Experience (Anavekar et al). 2006;19:28-33
Sequential changes of myocardial function during acute myocardial infarction, in the early and chronic phase after coronary intervention described by ultrasonic strain rate imaging (Weidemann et al). 2006;19:839-47
Spontaneous retraction of an intramyocardial dissecting hemorrhage and multiple left ventricular thrombus formation in subacute myocardial infarction and antiphospholipid syndrome: a case report with long-term follow-up (Bahlmann et al). 2006;19:578.e5-8 (Case rep.)
Transcatheter closure of a postinfarction ventricular septal rupture (Martinez et al). 2006;19:1401.e5-7 (Case rep.)
What is the optimal clinical technique for measurement of left ventricular volume after myocardial infarction? A comparative study of 3-dimensional echocardiography, single photon emission computed tomography, and cardiac magnetic resonance imaging (Chan et al). 2006;19:192-201

Myocardial ischemia

Analysis of postsystolic myocardial thickening work in selective myocardial layers during progressive myocardial ischemia (Wang et al). 2006;19:1102-11
Clinical outcomes after cardiac resynchronization therapy: importance of left ventricular diastolic function and origin of heart failure (Waggoner et al). 2006;19:307-13
Geometric differences of the mitral valve tenting between anterior and inferior myocardial infarction with significant ischemic mitral regurgitation: quantitation by novel software system with transthoracic real-time three-dimensional echocardiography (Watanabe et al). 2006;19:71-5
Noninvasive diagnosis of ischemic and nonischemic cardiomyopathy using coronary flow velocity measurements of the left anterior descending coronary artery by transthoracic Doppler echocardiography (Okura et al). 2006;19:552-8

Myocardial perfusion

Automated quantification of the spatial extent of perfusion defects and viability on myocardial contrast echocardiography (Micari et al). 2006;19:379-85
Corrected quantification method to determine myocardial blood flow using real-time myocardial contrast echocardiography (Otani et al). 2006;19:973-81
Heterogeneity of myocardial perfusion in distal coronary embolism with different particle sizes (Akiyama et al). 2006;19:55-63
Impact of the coronary flow reduction at rest on myocardial perfusion and functional indices derived from myocardial contrast and strain echocardiography (Okuda et al). 2006;19:781-7
Myocardial perfusion after percutaneous transluminal septal myocardial ablation as assessed by myocardial contrast echocardiography in patients with hypertrophic obstructive cardiomyopathy (Pedone et al). 2006;19:982-6
Noninvasive diagnosis of coronary artery bypass graft disease by dobutamine stress real-time myocardial contrast perfusion imaging (Elhendy et al). 2006;19:1482-7
Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested of known ischemic heart disease (Basic et al). 2006;19:1203-10
Quantification of myocardial perfusion using intravenous myocardial contrast echocardiography in healthy volunteers: comparison with positron emission tomography (Dijkmans et al). 2006;19:285-93

Myocardial revascularization

Dobutamine stress echocardiography is highly accurate for the prediction of contractile reserve in the early postoperative period, but may underestimate late recovery in contractile reserve after revascularization of the hibernating myocardium (Zaglavara et al). 2006;19:300-6

Myocardium

Acute improvement in myocardial function assessed by myocardial strain and strain rate after aortic valve replacement for aortic stenosis (Iwahashi et al). 2006;19:1238-44
Comparison of regional systolic myocardial velocities in patients with isolated left ventricular noncompaction and patients with idiopathic dilated cardiomyopathy (Tufekcioglu et al). 2006;19:1320-5
The effect of carvedilol therapy on myocardial functional reserve in patients with advanced heart failure caused by nonischemic dilated cardiomyopathy (Paraskevaidis et al). 2006;19:529-35
In vivo echocardiographic imaging of transplanted human adult stem cells in the myocardium labeled with clinically applicable CliniMACS nanoparticles (Bara et al). 2006;19:563-8
Left ventricular chamber and myocardial systolic function reserve in patients with type 1 diabetes mellitus: insight from traditional and Doppler tissue imaging echocardiography (Palmieri et al). 2006;19:848-56
Long-term survival of a patient with isolated noncompaction of the ventricular myocardium (Salemi et al). 2006;19:354.e1-3 (Case rep.)
Myocardial adaptation to short-term high-intensity exercise in highly trained athletes (Neilan et al). 2006;19:1280-5
Sequential changes of myocardial function during acute myo-

cardial infarction, in the early and chronic phase after coronary intervention described by ultrasonic strain rate imaging (Weidemann et al). 2006;19:839-47
Single beat determination of regional myocardial strain measurements in patients with atrial fibrillation (Yamanaka-Funabiki et al). 2006;19:1332-7

Myxoma
Smooth muscle neoplasm presenting as intrapericardial myxoma (Horton et al). 2006;19:835.e1-3 (Case rep.)

N

N-terminal pro b-type natriuretic peptide; *see* B-type natriuretic peptide; Peptide fragments

Natriuretic peptide, brain
Left bundle branch block in chronic heart failure-impact on diastolic function, filling pressures, and B-type natriuretic peptide levels (Bruch et al). 2006;19:95-101
Risk stratification in chronic heart failure: independent and incremental prognostic value of echocardiography and brain natriuretic peptide and its N-terminal fragment (Bruch et al). 2006;19:522-8

Neonate; *see* Infant, newborn

Neoplasm metastasis
Floating heart metastasis (Mumoli et al). 2006;19:1293.e1-2 (Case rep.)
Right ventricular metastasis caused by a renal cell carcinoma (Grawitz tumor): case report (Eddicks et al). 2006;19:1073.e11-5 (Case rep.)
Right ventricular metastasis of uterine squamous cell carcinoma (Nesser et al). 2006;19:1294.e1-3 (Case rep.)

Neuromuscular diseases
Hypertrabeculation is not distinct from noncompaction in neuromuscular and nonneuromuscular disorders (Stöllberger and Finsterer). 2006;19:109-10 (Letter)

Nitroglycerin
Assessment of coronary vasodilatation in response to nitroglycerin with transthoracic Doppler echocardiography (Sohn et al). 2006;19:777-80

O

Obesity
Obesity cardiomyopathy: is it a reality? An ultrasonic tissue characterization study (Di Bello et al). 2006;19:1063-71

Octogenarians; *see* Aged, 80 and over

P

Pacemaker, artificial
Optimization of pulsed wave tissue doppler to predict left ventricular reverse remodeling after cardiac resynchronization therapy (Jansen et al). 2006;19:185-91

Papillary muscles
Improved papillary muscle function attenuates functional mitral regurgitation in patients with dilated cardiomyopathy after cardiac resynchronization therapy (Karvounis et al). 2006;19:1150-7
Papillary muscle rupture after blunt chest trauma (Cordovil et al). 2006;19:469.e1-3 (Case rep.)
Unusual anomalous single papillary muscle causing symptomatic mid-left ventricular cavity obstruction: octopus papillary muscle (Shah et al). 2006;19:939.e9-11 (Case rep.)

Patent foramen ovale; *see* Heart septal defects, atrial

PB127; *see* Contrast media

Pediatric echocardiography; *see* Echocardiography, child

Peptide fragments
Risk stratification in chronic heart failure: independent and incremental prognostic value of echocardiography and brain natriuretic peptide and its N-terminal fragment (Bruch et al). 2006;19:522-8

Percutaneous coronary intervention; *see* Angioplasty, balloon

Perfusion
Primary cardiac lymphoma: diagnosis and treatment guided by transesophageal echocardiogram perfusion imaging (Jang et al). 2006;19:1073.e7-9 (Case rep.)

Pericarditis
Perforated periaortic valve abscess presents as acute pericarditis and patient died in transesophageal echocardiographic study (Tsai et al). 2006;19:579.e1-3 (Case rep.)

Pericarditis, constrictive
Pulmonary regurgitant flow and detection of dip plateau (Cohen et al). 2006;19:580 (Letter)

Pericardium
Smooth muscle neoplasm presenting as intrapericardial myxoma (Horton et al). 2006;19:835.e1-3 (Case rep.)

Perioperative care
American Society of Echocardiography/ Society of Cardiovascular Anesthesiologists recommendations and guidelines for continuous quality improvements in perioperative echocardiography (Mathew et al). 2006;19:1303-13

Phantoms, imaging
Single-beat differentiation among left ventricular filling pattern by pulsed wave Doppler echocardiography (Su et al). 2006;19:274-9

Phenylephrine
Intraoperative assessment of mitral regurgitation: role of phenylephrine challenge (Mihalatos et al). 2006;19:1158-64

Pheochromocytoma
Reduced myocardial contractility assessed by tissue Doppler echocardiography is associated with increased risk during adrenal surgery of patients with pheochromocytoma: report of a preliminary study (Meune et al). 2006;19:1466-70

Photon emission computed tomography; *see* Tomography, emission-computed, single photon

Pleural effusion
A new window of opportunity in echocardiography (Naqvi and Huynh). 2006;19:569-77

Polycythemia
Hematoma of a congenitally bicuspid aortic valve in a patient with polycythemia vera and the antiphospholipid antibody syndrome (Webber et al). 2006;19:1530.e1-3 (Case rep.)

Polycythemia vera; *see* Polycythemia

Positron-emission tomography
Quantification of myocardial perfusion using intravenous myocardial contrast echocardiography in healthy volunteers: comparison with positron emission tomography (Dijkmans et al). 2006;19:285-93

Postoperative complications
Esophageal dissection complicating transesophageal echocardiogram— the lesson to be learned: do not force the issue (El-Chami et al). 2006;19:579.e5-7 (Case rep.)

Postsystole; *see* Systole

Potassium
Glucose-insulin-potassium echocardiography detects improved segmental myocardial function and viable tissue shortly after acute myocardial infarction (Klein et al). 2006;19:763-71

Predictive value of tests
Early mitral filling/diastolic mitral annular velocity ratio is not a reliable predictor of left ventricular filling pressure in the setting of severe mitral regurgitation (Olson et al). 2006;19:83-7
Echocardiographic indices of increased left ventricular filling pressure and dilation after acute myocardial infarction (Hillis et al). 2006;19:450-6
Impact of the coronary flow reduction at rest on myocardial perfusion and functional indices derived from myocardial contrast and strain echocardiography (Okuda et al). 2006;19:781-7
Myocardial contrast echocardiographic estimates of infarct size predict likelihood of left ventricular remodeling after acute anterior wall myocardial infarction (Main et al). 2006;19:64-70

Prediction of subclinical left ventricular dysfunction with strain rate imaging in patients with mild to moderate rheumatic mitral stenosis (Dogan et al). 2006;19:243-8

Pregnancy

Assessment of the evolution of normal fetal diastolic function during mid and late gestation by spectral Doppler tissue echocardiography (Nii et al). 2006;19:1431-7

Prevalence

Increased prevalence of mitral stenosis in women (Movahed et al). 2006;19:911-3

Pro-brain natriuretic peptide; *see* Peptide fragments

Professional competence

American Society of Echocardiography minimum standards for the cardiac sonographer: a position paper (Bierig et al). 2006;19:471-4

Professional practice

The architecture of a mobile outreach echocardiography service (Gilman et al). 2006;19:1526-8

Prognosis

Effect of time delay on the diagnostic use of contrast echocardiography in patients presenting to the emergency department with chest pain and no S-T segment elevation (Kalvaitis et al). 2006;19:1488-93

Mitral annulus motion as a predictor of mortality in a community-based sample of 75-year-old men and women (Hedberg et al). 2006;19:88-94

Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested of known ischemic heart disease (Basic et al). 2006;19:1203-10

Risk assessment in patients with depressed left ventricular function after myocardial infarction using the myocardial performance index-Survival and Ventricular Enlargement (SAVE) Experience (Anavekar et al). 2006;19:28-33

Progressive systemic sclerosis; *see* Scleroderma, systemic

Propanolamines

The effect of carvedilol therapy on myocardial functional reserve in patients with advanced heart failure caused by nonischemic dilated cardiomyopathy (Paraskevaidis et al). 2006;19:529-35

Prosthesis failure

Heart failure and severe pulmonary hypertension caused by distal detachment of the valve conduit 16 years after the Cabrol composite graft procedure (Tosoratti et al). 2006;19:1190.e5-8 (Case rep.)

Proximal isovelocity surface area

Reply to "Angle correction for proximal isovelocity surface area method: is it spheric cap or lune? (Genc et al). 2006;19:940 (Letter)

Pulmonary artery

Hemodynamic change in transmitral gradient during Valsalva maneuver and its relation with left atrial function, functional capacity, and pulmonary artery pressure in patients with mitral stenosis in sinus rhythm (Ozeke et al). 2006;19:1245-50

Noninvasive Doppler tissue measurement of pulmonary artery compliance in children with pulmonary hypertension (Dyer et al). 2006;19:403-12

A novel echocardiographic Doppler method for estimation of pulmonary arterial pressures (Friedberg et al). 2006;19:559-62

The prognostic value of pulmonary vascular capacitance determined by Doppler echocardiography in patients with pulmonary arterial hypertension (Mahapatra et al). 2006;19:1045-50

Pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia: an echocardiographic perspective (Jassal et al). 2006;19:229.e5-7

Pulmonary artery dissection: a rare complication of pulmonary balloon valvuloplasty diagnosed 11 years after the procedure (Janus et al). 2006;19:1191.e5-8 (Case rep.)

Pulmonary hemodynamic effects of dipyridamole infusion in patients with normal and elevated pulmonary artery systolic pressure receiving PB127 (Main et al). 2006;19:1038-44

"Steal" collaterals: an echocardiographic diagnostic marker for anomalous origin of the left main coronary artery from the pulmonary artery in the adult (Pisacane et al). 2006;19:107.e3-6 (Case rep.)

Pulmonary atresia

Echocardiographic diagnosis and management of bilateral coronary ostial atresia in a patient with pulmonary atresia and intact ventricular septum (Wald et al). 2006;19:939.e1-3 (Case rep.)

Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum (Peterson et al). 2006;19:1393-1400

Pulmonary edema

Assessment of stress-induced pulmonary interstitial edema by chest ultrasound during exercise echocardiography and its correlation with left ventricular function (Agricola et al). 2006;19:457-63

Left ventricular filling hemodynamics in patients with pulmonary edema and preserved versus reduced left ventricular ejection fraction: a prospective Doppler echocardiographic study (Dabbah et al). 2006;19:733-43

An unusual cause of pulmonary edema: acute rupture of noncoronary sinus of Valsalva aneurysm into the left atrium (Moreo et al). 2006;19:938.e9-11 (Case rep.)

Pulmonary embolism

Atrial and ventricular echocardiographic correlates of the extent of pulmonary embolism in the elderly (Chung et al). 2006;19:347-53

Pulmonary embolism and right heart function: insights from myocardial Doppler tissue imaging (Hsiao et al). 2006;19:822-8

Quantitative measures of right ventricular dysfunction by echocardiography in the diagnosis of acute nonmassive pulmonary embolism (Kjaergaard et al). 2006;19:1264-71

Pulmonary hypertension; *see* Hypertension, pulmonary

Pulmonary stenosis

Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum (Peterson et al). 2006;19:1393-1400

Pulmonary valve insufficiency

Pulmonary regurgitant flow and detection of dip plateau (Cohen et al). 2006;19:580 (Letter)

Pulmonary valve stenosis

Response to: "Echocardiographic assessment of isolated pulmonary valve stenosis: which outpatient Doppler gradient has the most clinically validity?" (DeGroff). 2006;19:364 (Letter)

Pulmonary veins

Pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia: an echocardiographic perspective (Jassal et al). 2006;19:229.e5-7

Subtotal anomalous pulmonary venous connection and left cor triatriatum: a rare diagnosis in adulthood (Caputo et al). 2006;19:836.e1-4 (Case rep.)

Pulse

Optimization of pulsed wave tissue doppler to predict left ventricular reverse remodeling after cardiac resynchronization therapy (Jansen et al). 2006;19:185-91

Q

Quality of health care

American Society of Echocardiography/ Society of Cardiovascular Anesthesiologists recommendations and guidelines for continuous quality improvements in perioperative echocardiography (Mathew et al). 2006;19:1303-13

American Society of Echocardiography minimum standards for the cardiac sonographer: a position paper (Bierig et al). 2006;19:471-4

The assessment of technical and interpretative proficiency in echocardiography (Nair et al). 2006;19:924-31

R

Reconstructive surgical procedures
Unusual complication of aortic root reconstruction with sparing of the aortic valve: left ventricular outflow tract to right atrial fistula (Ramasubbu et al). 2006;19:469.e5-9 (Case rep.)
Rejection, transplant; *see* Graft rejection
Renal cell carcinoma; *see* Carcinoma, renal cell
Reproducibility of results
Accuracy, reproducibility, and comparability of Doppler tissue imaging by two high-end ultrasound systems (Kjaergaard et al). 2006;19:322-8
Contrast echocardiography versus gated single photon emission computed tomography for the assessment of parameters of left ventricular remodeling after acute myocardial infarction (Lim et al). 2006;19:280-4
Doppler tissue imaging in the assessment of atrioventricular conduction time: validation of a novel technique and comparison with electrophysiologic and pulsed wave Doppler-derived equivalents in an animal model (Nii et al). 2006;19:314-21
Feasibility and accuracy of real-time 3-dimensional echocardiographic assessment of ventricular septal defects (van den Bosch et al). 2006;19:7-13
Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease (van den Bosch et al). 2006;19:1-6
Response to: "Echocardiographic assessment of isolated pulmonary valve stenosis: which outpatient Doppler gradient has the most clinically validity?" (DeGroff). 2006;19:364 (Letter)
Single-beat differentiation among left ventricular filling pattern by pulsed wave Doppler echocardiography (Su et al). 2006;19:274-9
Use of myocardial performance index in pediatric patients with idiopathic pulmonary arterial hypertension (Dyer et al). 2006;19:21-7
Restrictive cardiomyopathy; *see* Cardiomyopathy, restrictive
Review articles
Review articles). 2006;19:1403-10
Rheology
Corrected quantification method to determine myocardial blood flow using real-time myocardial contrast echocardiography (Otani et al). 2006;19:973-81
Right ventricle; *see* Heart ventricle
Right ventricular dysfunction; *see* Ventricular dysfunction, right
Right ventricular function; *see* Ventricular function, right
Risk assessment
Clinical application of noninvasive vascular ultrasound in cardiovascular risk stratification: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Roman et al). 2006;19:943-54 (ASE report)
Risk stratification by treadmill exercise echocardiography (Peteiro et al). 2006;19:894-901
Risk factors
Reduced myocardial contractility assessed by tissue Doppler echocardiography is associated with increased risk during adrenal surgery of patients with pheochromocytoma: report of a preliminary study (Meune et al). 2006;19:1466-70
Relationship between the number of coronary risk factors and coronary atherosclerosis assessed by high-frequency transthoracic echocardiography (Takeuchi et al). 2006;19:1056-62
Risk assessment in patients with depressed left ventricular function after myocardial infarction using the myocardial performance index-Survival and Ventricular Enlargement (SAVE) Experience (Anavekar et al). 2006;19:28-33
Use of echocardiography in patients with hypertrophic cardiomyopathy: clinical implications of massive hypertrophy (Sorajja et al). 2006;19:788-95

S

Saline; *see* Sodium chloride
Scleroderma, systemic
Right heart function in scleroderma: insights from myocardial Doppler tissue imaging (Hsiao et al). 2006;19:507-14
Septum, heart; *see* Heart septum
Severity of illness index
Hemodynamic significance of mitral stenosis: use of a simple, novel index by 2-dimensional echocardiography (Seow et al). 2006;19:102-6
Relationship of mitral annular remodeling to severity of chronic mitral regurgitation (Mihalatos et al). 2006;19:76-82
Simendan
Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: a doppler echocardiographic study (Luca et al). 2006;19:172-7
Single photon emission computed tomography; *see* Tomography, emission-computed, single photon; Tomography, emission-computed, single-photon
Sinus of Valsalva
An unusual cause of pulmonary edema: acute rupture of noncoronary sinus of Valsalva aneurysm into the left atrium (Moreo et al). 2006;19:938.e9-11 (Case rep.)
Sinus rhythm
Hemodynamic change in transmitral gradient during Valsalva maneuver and its relation with left atrial function, functional capacity, and pulmonary artery pressure in patients with mitral stenosis in sinus rhythm (Ozeke et al). 2006;19:1245-50
Society of Cardiovascular Anesthesiologists
American Society of Echocardiography/ Society of Cardiovascular Anesthesiologists recommendations and guidelines for continuous quality improvements in perioperative echocardiography (Mathew et al). 2006;19:1303-13
Society of Vascular Medicine and Biology
Clinical application of noninvasive vascular ultrasound in cardiovascular risk stratification: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Roman et al). 2006;19:943-54 (ASE report)
Guidelines for noninvasive vascular laboratory testing: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Gerhard-Herman et al). 2006;19:955-72 (ASE report)
Sodium chloride
A critical review of patent foramen ovale detection using saline contrast echocardiography: when bubbles lie (Woods and Patel). 2006;19:215-22
SPECT; *see* Tomography, emission-computed, sinlge-photon
Sports
Myocardial adaptation to short-term high-intensity exercise in highly trained athletes (Neilan et al). 2006;19:1280-5
Squamous cell carcinoma; *see* Carcinoma, squamous cell
Staphylococcus
Biventricular mural vegetations in a patient without valvular pathology (Ahmed et al). 2006;19:938.e5-7 (Case rep.)
Stem cell transplantation
In vivo echocardiographic imaging of transplanted human adult stem cells in the myocardium labeled with clinically applicable CliniMACS nanoparticles (Bara et al). 2006;19:563-8
Stents
Intracoronary stent visualized on transesophageal echocardiogram in a case of coronary dissection complicated by aortic dissection (Pohlel et al). 2006;19:229.e1-3 (Case rep.)
Stent graft in treatment of penetrating thoracic aortic ulcer (Khamaisi et al). 2006;19:835.e4-6 (Case rep.)
Strain rate
Accuracy, reproducibility, and comparability of Doppler tissue

imaging by two high-end ultrasound systems (Kjaergaard et al). 2006;19:322-8
Acute improvement in myocardial function assessed by myocardial strain and strain rate after aortic valve replacement for aortic stenosis (Iwahashi et al). 2006;19:1238-44
Analysis of mitral annulus motion measurements derived from M-mode, anatomic M-mode, tissue Doppler displacement, and 2-dimensional strain imaging (Hayashi et al). 2006;19:1092-1101
Detection of significant stenotic lesions in the left anterior descending coronary artery using adenosine triphosphate stress strain imaging: comparison with coronary flow velocity reserve measurement using transthoracic Doppler echocardiography (Takagi et al). 2006;19:1001-11
Doppler tissue, strain, and strain rate imaging in pediatric patients with Alström syndrome: are there regional functional abnormalities? (Toulany et al). 2006;19:14-20
Effects of ultrasound contrast during tissue velocity imaging on regional left ventricular velocity, strain, and strain rate measurements (Malm et al). 2006;19:40-7
Improved regional myocardial diastolic function assessed by strain rate imaging in patients undergoing percutaneous coronary intervention (Tanaka et al). 2006;19:756-62
Myocardial strain decreases with increasing transmurality of infarction: a Doppler echocardiographic and magnetic resonance correlation study (Sachdev et al). 2006;19:34-9
New aspects of septal function by using 1-dimensional strain and strain rate imaging (Lindqvist et al). 2006;19:1345-9
Prediction of subclinical left ventricular dysfunction with strain rate imaging in patients with mild to moderate rheumatic mitral stenosis (Dogan et al). 2006;19:243-8
Rapid ventricular remodeling with left ventricular unloading postventricular assist device placement: new insights with strain imaging (Havemann et al). 2006;19:355.e9-11 (Case rep.)
Regional postsystolic shortening in patients with hypertrophic cardiomyopathy: its incidence and characteristics assessed by strain imaging (Ito et al). 2006;19:987-93
Role of left ventricular regional diastolic abnormalities for global diastolic dysfunction in patients with hypertrophic cardiomyopathy (Goto et al). 2006;19:857-64
Sequential changes of myocardial function during acute myocardial infarction, in the early and chronic phase after coronary intervention described by ultrasonic strain rate imaging (Weidemann et al). 2006;19:839-47
Ultrasonic strain rate and strain imaging of the right ventricle in children before and after percutaneous closure of an atrial septal defect (Eyskens et al). 2006;19:994-1000

Stress echocardiography; *see* Echocardiography, stress

Stress test; *see* Exercise test

Stroke; *see* Cerebrovascular disorders

Stroke volume
Comparison of regional versus global assessment of left ventricular function in patients with left ventricular dysfunction, heart failure, or both after myocardial infarction: the Valsartan in Acute Myocardial Infarction Echocardiographic study (Thune et al). 2006;19:1462-5
The effect of acute hypertension on left ventricular diastolic pressures in a canine model of left ventricular dysfunction with a preserved ejection fraction and elevated left ventricular filling pressures (Lavine and Lavine). 2006;19:1350-8
Evaluation of midwall systolic function in left ventricular hypertrophy: a comparison of 3-dimensional versus 2-dimensional echocardiographic indices (Jung et al). 2006;19:802-10
Left ventricular filling hemodynamics in patients with pulmonary edema and preserved versus reduced left ventricular ejection fraction: a prospective Doppler echocardiographic study (Dabbah et al). 2006;19:733-43
Real-time simultaneous triplane contrast echocardiography gives rapid, accurate, and reproducible assessment of left ventricular volumes and ejection fraction: a comparison with magnetic resonance imaging (Malm et al). 2006;19:1494-1501
Real-time three-dimensional color Doppler echocardiography overcomes the inaccuracies of spectral Doppler for stroke volume calculation (Pemberton et al). 2006;19:1403-10 (Review article)
Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease (van den Bosch et al). 2006;19:1-6
Severe tricuspid regurgitation shows significant impact in the relationship among peak systolic tricuspid annular velocity, tricuspid annular plane systolic excursion, and right ventricular ejection fraction (Hsiao et al). 2006;19:902-10

Sudden cardiac death; *see* Death, sudden, cardiac

Sulfonamides
Use of myocardial performance index in pediatric patients with idiopathic pulmonary arterial hypertension (Dyer et al). 2006;19:21-7

Surgery, heart; *see* Heart surgery

Syncope
Syncope with a twist (Gosavi et al). 2006;19:1529.e1-3 (Case rep.)

Systemic lupus erythematosus; *see* Lupus erythematosus, systemic

Systole
Analysis of postsystolic myocardial thickening work in selective myocardial layers during progressive myocardial ischemia (Wang et al). 2006;19:1102-11
Asymmetry of systolic anterior motion of the mitral valve in patients with hypertrophic obstructive cardiomyopathy: a real-time three-dimensional echocardiography study (Song et al). 2006;19:1129-35
Asynchrony detection and Doppler tissue imaging (Gaibazzi). 2006;19:111 (Letter)
Asynchrony of left ventricular systolic performance after the first acute myocardial infarction in patients with narrow QRS complexes: Doppler tissue imaging study (Elnoamany et al). 2006;19:1449-57
Comparison of regional systolic myocardial velocities in patients with isolated left ventricular noncompaction and patients with idiopathic dilated cardiomyopathy (Tufekcioglu et al). 2006;19:1320-5
Doppler tissue, strain, and strain rate imaging in pediatric patients with Alström syndrome: are there regional functional abnormalities? (Toulany et al). 2006;19:14-20
Effect of aging on twist-displacement loop by 2-dimensional speckle tracking imaging (Nakai et al). 2006;19:880-5
Evaluation of midwall systolic function in left ventricular hypertrophy: a comparison of 3-dimensional versus 2-dimensional echocardiographic indices (Jung et al). 2006;19:802-10
Impact of anesthesia on echocardiographic evaluation of systolic and diastolic function in rats (Plante et al). 2006;19:1520-5
Isolated ventricular diastolic dysfunction: implications for exercise left ventricular performance in patients without congestive heart failure (Palmieri et al). 2006;19:491-8
Left ventricular chamber and myocardial systolic function reserve in patients with type 1 diabetes mellitus: insight from traditional and Doppler tissue imaging echocardiography (Palmieri et al). 2006;19:848-56
Left ventricular systolic longitudinal function: comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals (Mondillo et al). 2006;19:1085-91
Myocardial strain decreases with increasing transmurality of infarction: a Doppler echocardiographic and magnetic resonance correlation study (Sachdev et al). 2006;19:34-9
Regional postsystolic shortening in patients with hypertrophic

cardiomyopathy: its incidence and characteristics assessed by strain imaging (Ito et al). 2006;19:987-93
Severe tricuspid regurgitation shows significant impact in the relationship among peak systolic tricuspid annular velocity, tricuspid annular plane systolic excursion, and right ventricular ejection fraction (Hsiao et al). 2006;19:902-10
Single beat determination of regional myocardial strain measurements in patients with atrial fibrillation (Yamanaka-Funabiki et al). 2006;19:1332-7
The systolic to diastolic duration ratio in children with heart failure secondary to restrictive cardiomyopathy (Friedberg and Silverman). 2006;19:1326-31
Usefulness of ventricular longitudinal contractility assessed by doppler tissue imaging in the prediction of reverse remodeling in patients with severe left ventricular systolic dysfunction (Kang et al). 2006;19:178-84

T

Takayasu's arteritis
Thoracic aorta evaluation in patients with Takayasu's arteritis by transesophageal echocardiography (Lira-Filho et al). 2006;19:829-34
Takotsubo cardiomyopathy; *see* Myocardial diseases
Telangiectasia, hereditary hemorrhagic
Pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia: an echocardiographic perspective (Jassal et al). 2006;19:229.e5-7
Thoracic aorta; *see* Aorta, thoracic
Thoracic injuries
Papillary muscle rupture after blunt chest trauma (Cordovil et al). 2006;19:469.e1-3 (Case rep.)
Three-dimensional echocardiography; *see* Echocardiography, three-dimensional
Thromboembolism
Cardioversion in patients with left ventricular thrombus is not associated with increased thromboembolic risk (Bangalore et al). 2006;19:438-40
Thrombolytic therapy
Detection of residual infarct-related coronary artery stenosis and multivessel disease after thrombolysis: comparison between myocardial contrast echocardiography and single photon emission computed tomography (Dwivedi et al). 2006;19:546-51
Thrombosis
Cardioversion in patients with left ventricular thrombus is not associated with increased thromboembolic risk (Bangalore et al). 2006;19:438-40
Improved detection of left ventricular thrombi and spontaneous echocontrast by tissue harmonic imaging in patients with myocardial infarction (Mele et al). 2006;19:1373-81
Spontaneous retraction of an intramyocardial dissecting hemorrhage and multiple left ventricular thrombus formation in subacute myocardial infarction and antiphospholipid syndrome: a case report with long-term follow-up (Bahlmann et al). 2006;19:578.e5-8 (Case rep.)
Syncope with a twist (Gosavi et al). 2006;19:1529.e1-3 (Case rep.)
Thrombotic "ghosts": echocardiographic appearance of thrombi with hollow cores and implications regarding mechanism of spontaneous clot lysis (Lavi et al). 2006;19:1530.e5-6 (Case rep.)
Transesophageal echocardiographic assessment of right atrial appendage anatomy and function: comparison with the left atrial appendage and implications for local thrombus formation (Subramaniam et al). 2006;19:429-33
Transit thrombus entrapped in patent foramen ovale resolved without clinical embolic events (Can et al). 2006;19:1074.e1-2 (Case rep.)
Twenty-five-year-old man with a giant left ventricular mass (Moore and Segar). 2006;19:1294.e7-8 (Case rep.)
Time
Effect of time delay on the diagnostic use of contrast echocardiography in patients presenting to the emergency department with chest pain and no S-T segment elevation (Kalvaitis et al). 2006;19:1488-93
Tissue Doppler imaging; *see* Doppler tissue imaging
Tomography, emission-computed, single-photon
Contrast echocardiography versus gated single photon emission computed tomography for the assessment of parameters of left ventricular remodeling after acute myocardial infarction (Lim et al). 2006;19:280-4
Detection of residual infarct-related coronary artery stenosis and multivessel disease after thrombolysis: comparison between myocardial contrast echocardiography and single photon emission computed tomography (Dwivedi et al). 2006;19:546-51
Noninvasive diagnosis of restenosis by transthoracic doppler echocardiography after percutaneous coronary intervention: comparison with exercise Tl-SPECT (Hirata et al). 2006;19:165-71
Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested of known ischemic heart disease (Basic et al). 2006;19:1203-10
What is the optimal clinical technique for measurement of left ventricular volume after myocardial infarction? A comparative study of 3-dimensional echocardiography, single photon emission computed tomography, and cardiac magnetic resonance imaging (Chan et al). 2006;19:192-201
Torsion
Age-related changes in left ventricular twist assessed by two-dimensional speckle-tracking imaging (Takeuchi et al). 2006;19:1077-84
Effect of aging on twist-displacement loop by 2-dimensional speckle tracking imaging (Nakai et al). 2006;19:880-5
Transesophageal echocardiography; *see* Echocardiography, transesophageal
Transposition of great vessels
Right ventricular function in asymptomatic individuals with a systemic right ventricle (Bos et al). 2006;19:1033-7
Transthoracic echocardiography; *see* Echocardiography, transthoracic
Treadmill test; *see* Exercise test
Tricuspid valve
Tricuspid annular velocity in patients undergoing cardiac operation using transesophageal echocardiography (David et al). 2006;19:329-34
Tricuspid valve insufficiency
Severe tricuspid regurgitation shows significant impact in the relationship among peak systolic tricuspid annular velocity, tricuspid annular plane systolic excursion, and right ventricular ejection fraction (Hsiao et al). 2006;19:902-10
Three-dimensional tricuspid annular function provides insight into the mechanisms of tricuspid valve regurgitation in classic hypoplastic left heart syndrome (Nii et al). 2006;19:391-402
Trisomy
Extreme redundancy of the valve of the fossa ovalis with right heart hypoplasia in a neonate with trisomy 18 (Vyas and Cabalka). 2006;19:1293.e7-8 (Case rep.)
Tunica intima
Use of carotid intima-media thickness and vascular age to modify cardiovascular risk prediction (Gepner et al). 2006;19:1170-4
Use of hand-held ultrasound by a nonsonographer clinician to measure carotid intima-media thickness (Tzou et al). 2006;19:1286-92
Validation of a carotid intima-media thickness border detection program for use in an office setting (Gepner et al). 2006;19:223-8
Two-dimensional echocardiography; *see* Echocardiography, two-dimensional

Type 1 diabetes; *see* Diabetes mellitus, insulin-dependent
Type 2 diabetes; *see* Diabetes mellitus, non-insulin-dependent

U

Ulcer
Stent graft in treatment of penetrating thoracic aortic ulcer (Khamaisi et al). 2006;19:835.e4-6 (Case rep.)
Ultrasonography
Accuracy, reproducibility, and comparability of Doppler tissue imaging by two high-end ultrasound systems (Kjaergaard et al). 2006;19:322-8
Age-related changes in left ventricular twist assessed by two-dimensional speckle-tracking imaging (Takeuchi et al). 2006; 19:1077-84
Clinical application of noninvasive vascular ultrasound in cardiovascular risk stratification: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Roman et al). 2006;19:943-54 (ASE report)
Effect of aging on twist-displacement loop by 2-dimensional speckle tracking imaging (Nakai et al). 2006;19:880-5
Effects of ultrasound contrast during tissue velocity imaging on regional left ventricular velocity, strain, and strain rate measurements (Malm et al). 2006;19:40-7
Efficacy of ultrasound-assisted stress testing using a hand-carried ultrasound device for diagnosis of coronary artery disease (Hirano et al). 2006;19:536-9
Myocardial lesion formation using high-intensity focused ultrasound (Engel et al). 2006;19:932-7
Obesity cardiomyopathy: is it a reality? An ultrasonic tissue characterization study (Di Bello et al). 2006;19:1063-71
Ultrasonic strain rate and strain imaging of the right ventricle in children before and after percutaneous closure of an atrial septal defect (Eyskens et al). 2006;19:994-1000
Ultrasound lung comets: a clinically useful sign of extravascular lung water (Picano et al). 2006;19:356-63 (Review article)
Use of hand-held ultrasound by a nonsonographer clinician to measure carotid intima-media thickness (Tzou et al). 2006; 19:1286-92
Uterine cervical neoplasms
Right ventricular metastasis of uterine squamous cell carcinoma (Nesser et al). 2006;19:1294.e1-3 (Case rep.)

V

Valsalva maneuver
Hemodynamic change in transmitral gradient during Valsalva maneuver and its relation with left atrial function, functional capacity, and pulmonary artery pressure in patients with mitral stenosis in sinus rhythm (Ozeke et al). 2006;19: 1245-50
Valve replacement; *see* Heart valve prosthesis
Valvular regurgitation; *see* Aortic valve insufficiency; Mitral valve insufficiency; Tricuspid valve insufficiency
Valvuloplasty; *see* Balloon dilatation
Vascular capacitance
The prognostic value of pulmonary vascular capacitance determined by Doppler echocardiography in patients with pulmonary arterial hypertension (Mahapatra et al). 2006;19: 1045-50
Vascular diseases
Guidelines for noninvasive vascular laboratory testing: a report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology (Gerhard-Herman et al). 2006;19:955-72 (ASE report)
Vasodilation
Assessment of coronary vasodilatation in response to nitroglycerin with transthoracic Doppler echocardiography (Sohn et al). 2006;19:777-80
Ventricular assist device; *see* Heart-assist devices
Ventricular dysfunction, left
Abnormal longitudinal myocardial functional reserve assessed by exercise tissue Doppler echocardiography in patients with hypertrophic cardiomyopathy (Ha et al). 2006;19: 1314-9
Asynchrony of left ventricular systolic performance after the first acute myocardial infarction in patients with narrow QRS complexes: Doppler tissue imaging study (Elnoamany et al). 2006;19:1449-57
Can transthoracic live 3-dimensional echocardiography improve the recognition of midventricular obliteration in hypertrophic obstructive cardiomyopathy? (Gregorio et al). 2006;19:1190.e1-4 (Case rep.)
Comparison of regional systolic myocardial velocities in patients with isolated left ventricular noncompaction and patients with idiopathic dilated cardiomyopathy (Tufekcioglu et al). 2006;19:1320-5
Comparison of regional versus global assessment of left ventricular function in patients with left ventricular dysfunction, heart failure, or both after myocardial infarction: the Valsartan in Acute Myocardial Infarction Echocardiographic study (Thune et al). 2006;19:1462-5
Echocardiographic indices of increased left ventricular filling pressure and dilation after acute myocardial infarction (Hillis et al). 2006;19:450-6
The effect of acute hypertension on left ventricular diastolic pressures in a canine model of left ventricular dysfunction with a preserved ejection fraction and elevated left ventricular filling pressures (Lavine and Lavine). 2006;19:1350-8
Effects of radial left ventricular dyssynchrony on cardiac performance using quantitative tissue Doppler radial strain imaging (Dohi et al). 2006;19:475-82
Gestational diabetes: a condition of early diastolic abnormalities in young women (Freire et al). 2006;19:1251-6
Improved regional myocardial diastolic function assessed by strain rate imaging in patients undergoing percutaneous coronary intervention (Tanaka et al). 2006;19:756-62
Isolated ventricular diastolic dysfunction: implications for exercise left ventricular performance in patients without congestive heart failure (Palmieri et al). 2006;19:491-8
Long-term survival of a patient with isolated noncompaction of the ventricular myocardium (Salemi et al). 2006;19:354.e1-3 (Case rep.)
Mitral peak Doppler E-wave to peak mitral annulus velocity ratio is an accurate estimate of left ventricular filling pressure and predicts mortality in end-stage renal disease (Sharma et al). 2006;19:266-73
Prediction of subclinical left ventricular dysfunction with strain rate imaging in patients with mild to moderate rheumatic mitral stenosis (Dogan et al). 2006;19:243-8
Preoperative pulmonary hypertension is associated with postoperative left ventricular dysfunction in chronic organic mitral regurgitation: an echocardiographic and hemodynamic study (Yang et al). 2006;19:1051-5
QRS duration alone misses cardiac dyssynchrony in a substantial proportion of patients with chronic heart failure (Perry et al). 2006;19:1257-63
Role of left ventricular regional diastolic abnormalities for global diastolic dysfunction in patients with hypertrophic cardiomyopathy (Goto et al). 2006;19:857-64
Single-beat differentiation among left ventricular filling pattern by pulsed wave Doppler echocardiography (Su et al). 2006; 19:274-9
Takotsubo cardiomyopathy: a case report (Mitchell and Crone). 2006;19:1190.e9-10 (Case rep.)
Unusual anomalous single papillary muscle causing symptomatic mid-left ventricular cavity obstruction: octopus papillary muscle (Shah et al). 2006;19:939.e9-11 (Case rep.)
Usefulness of clinical, electrocardiographic, and echocardiographic parameters to detect cardiac asynchrony in patients with left ventricular dysfunction secondary to ischemic or nonischemic heart disease (Perez de Isla et al). 2006;19: 1338-44

Ventricular dysfunction, right
Quantitative measures of right ventricular dysfunction by echocardiography in the diagnosis of acute nonmassive pulmonary embolism (Kjaergaard et al). 2006;19:1264-71
Sequence of echocardiographic changes during development of right ventricular failure in rat (Hardziyenka et al). 2006;19:1272-9
Severe tricuspid regurgitation shows significant impact in the relationship among peak systolic tricuspid annular velocity, tricuspid annular plane systolic excursion, and right ventricular ejection fraction (Hsiao et al). 2006;19:902-10
Ventricular ejection fraction; *see* Stroke volume
Ventricular function
Preservation of ventricular function in amateur athletes after completion of a marathon (Kean et al). 2006;19:202-5
The tei index identifies a differential effect on left and right ventricular function with low-dose anthracycline chemotherapy (Belham et al). 2006;19:206-10
Ventricular function, left
Accuracy and feasibility of online 3-dimensional echocardiography for measurement of left ventricular parameters (Jenkins et al). 2006;19:1119-28
Alterations in left ventricular structure and function in type-1 diabetics: a focus on left atrial contribution to function (Peterson et al). 2006;19:749-55
Assessment of stress-induced pulmonary interstitial edema by chest ultrasound during exercise echocardiography and its correlation with left ventricular function (Agricola et al). 2006;19:457-63
Asynchrony detection and Doppler tissue imaging (Gaibazzi). 2006;19:111 (Letter)
Characterization of left ventricular diastolic function in hypertension by use of Doppler tissue imaging and color M-mode techniques (Rovner et al). 2006;19:872-9
Clinical outcomes after cardiac resynchronization therapy: importance of left ventricular diastolic function and origin of heart failure (Waggoner et al). 2006;19:307-13
Color encoding of endocardial motion improves the interpretation of contrast-enhanced echocardiographic stress tests by less-experienced readers (Mor-avi et al). 2006;19:48-54
Color tissue Doppler myocardial velocities consistently underestimate spectral tissue Doppler velocities: impact on calculation peak transmitral pulsed Doppler velocity/early diastolic tissue Doppler velocity (E/Ea) (McCulloch et al). 2006;19:744-8
Early mitral filling/diastolic mitral annular velocity ratio is not a reliable predictor of left ventricular filling pressure in the setting of severe mitral regurgitation (Olson et al). 2006;19:83-7
Effects of levosimendan on left ventricular diastolic function after primary angioplasty for acute anterior myocardial infarction: a doppler echocardiographic study (Luca et al). 2006;19:172-7
Effects of ultrasound contrast during tissue velocity imaging on regional left ventricular velocity, strain, and strain rate measurements (Malm et al). 2006;19:40-7
Glucose-insulin-potassium echocardiography detects improved segmental myocardial function and viable tissue shortly after acute myocardial infarction (Klein et al). 2006;19:763-71
Left bundle branch block in chronic heart failure-impact on diastolic function, filling pressures, and B-type natriuretic peptide levels (Bruch et al). 2006;19:95-101
Left ventricular filling hemodynamics in patients with pulmonary edema and preserved versus reduced left ventricular ejection fraction: a prospective Doppler echocardiographic study (Dabbah et al). 2006;19:733-43
Left ventricular systolic longitudinal function: comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals (Mondillo et al). 2006;19:1085-91
Left ventricular Tei index in children: comparison of Doppler tissue imaging, pulsed wave Doppler, and M-mode echocardiography normal values (Cui and Roberson). 2006;19:1438-45
Letters to the editor (Ogunyanklu) (Letter); (McCulloch and Dokainish) (Reply). 2006;19:1411-2
Longitudinal isovolumic displacement of the left ventricular myocardium assessed by tissue velocity echocardiography in healthy individuals (Lind et al). 2006;19:255-65
Mitral annulus motion as a predictor of mortality in a community-based sample of 75-year-old men and women (Hedberg et al). 2006;19:88-94
N-terminal pro B-type natriuretic peptide levels: correlation with echocardiographically determined left ventricular diastolic function in an ambulatory cohort (Dong et al). 2006;19:1017-25
Optimization of pulsed wave tissue doppler to predict left ventricular reverse remodeling after cardiac resynchronization therapy (Jansen et al). 2006;19:185-91
Quantification of regional left ventricular wall motion from real-time 3-dimensional echocardiography in patients with poor acoustic windows: effects of contrast enhancement tested against cardiac magnetic resonance (Corsi et al). 2006;19:886-93
Real-time simultaneous triplane contrast echocardiography gives rapid, accurate, and reproducible assessment of left ventricular volumes and ejection fraction: a comparison with magnetic resonance imaging (Malm et al). 2006;19:1494-1501
Real-time transthoracic three-dimensional echocardiographic assessment of left ventricular volume and ejection fraction in congenital heart disease (van den Bosch et al). 2006;19:1-6
Relationship between left atrial and left ventricular function in hypertrophic cardiomyopathy: a real-time 3-dimensional echocardiographic study (Shin et al). 2006;19:796-801
Risk assessment in patients with depressed left ventricular function after myocardial infarction using the myocardial performance index-Survival and Ventricular Enlargement (SAVE) Experience (Anavekar et al). 2006;19:28-33
The role of left ventricular long-axis contraction in patients with asymptomatic aortic regurgitation (Paraskevaidis et al). 2006;19:249-54
Traditional and color M-mode parameters of left ventricular diastolic function during low-dose dobutamine stress echocardiography: relations to contractility reserve (Palmieri et al). 2006;19:483-90
Transesophageal echocardiographic acquisition of mitral annulus motion during monitoring of left ventricular function in anesthetized patients (Nilsson et al). 2006;19:499-506
Tricuspid annular velocity in patients undergoing cardiac operation using transesophageal echocardiography (David et al). 2006;19:329-34
Usefulness of ventricular longitudinal contractility assessed by doppler tissue imaging in the prediction of reverse remodeling in patients with severe left ventricular systolic dysfunction (Kang et al). 2006;19:178-84
Ventricular function, right
Impact of preload and afterload on global and regional right ventricular function and pressure: a quantitative echocardiography study (Kjaergaard et al). 2006;19:515-21
Pulmonary embolism and right heart function: insights from myocardial Doppler tissue imaging (Hsiao et al). 2006;19:822-8
Quantitative echocardiographic analysis of the right ventricle in healthy individuals (Kjaergaard et al). 2006;19:1365-72
Right heart function in systemic lupus erythematosus: insights from myocardial Doppler tissue imaging (Gin et al). 2006;19:441-9
Right ventricular function in asymptomatic individuals with a systemic right ventricle (Bos et al). 2006;19:1033-7
Ultrasonic strain rate and strain imaging of the right ventricle in

children before and after percutaneous closure of an atrial septal defect (Eyskens et al). 2006;19:994-1000

Ventricular outflow obstruction

A case report of a round cystic tumor in the left ventricular outflow tract (Pérez-Sanz et al). 2006;19:1402.e9-11 (Case rep.)

Hypotension during dobutamine stress transesophageal echocardiography: relationship with provoked left ventricular obstruction (Siddiqui et al). 2006;19:1144-9

Left ventricular outflow tract obstruction caused by accessory mitral valve tissue (Sharma et al). 2006;19:354.e5-8 (Case rep.)

Unusual anomalous single papillary muscle causing symptomatic mid-left ventricular cavity obstruction: octopus papillary muscle (Shah et al). 2006;19:939.e9-11 (Case rep.)

Ventricular remodeling

Contrast echocardiography versus gated single photon emission computed tomography for the assessment of parameters of left ventricular remodeling after acute myocardial infarction (Lim et al). 2006;19:280-4

Coronary flow velocity pattern assessed noninvasively by transthoracic color Doppler echocardiography serves as a predictor of adverse cardiac events and left ventricular remodeling in patients with acute myocardial infarction (Katayama et al). 2006;19:335-40

Myocardial contrast echocardiographic estimates of infarct size predict likelihood of left ventricular remodeling after acute anterior wall myocardial infarction (Main et al). 2006;19:64-70

Rapid ventricular remodeling with left ventricular unloading postventricular assist device placement: new insights with strain imaging (Havemann et al). 2006;19:355.e9-11 (Case rep.)

Ventricular septal defects; *see* Heart septal defects, ventricular

W

Wall motion

Color encoding of endocardial motion improves the interpretation of contrast-enhanced echocardiographic stress tests by less-experienced readers (Mor-avi et al). 2006;19:48-54

Comparison of contrast-enhanced real-time live 3-dimensional dobutamine stress echocardiography with contrast 2-dimensional echocardiography for detecting stress-induced wall-motion abnormalities (Takeuchi et al). 2006;19:294-9

Comparison of regional versus global assessment of left ventricular function in patients with left ventricular dysfunction, heart failure, or both after myocardial infarction: the Valsartan in Acute Myocardial Infarction Echocardiographic study (Thune et al). 2006;19:1462-5

Feasibility of using a real-time 3-dimensional technique for contrast dobutamine stress echocardiography (Pulerwitz et al). 2006;19:540-5

Quantification of regional left ventricular wall motion from real-time 3-dimensional echocardiography in patients with poor acoustic windows: effects of contrast enhancement tested against cardiac magnetic resonance (Corsi et al). 2006;19:886-93